



PlantGen2019

The Fifth International Scientific Conference

Current Challenges in Plant Genetics, Genomics, Bioinformatics, and Biotechnology



JUNE 24–29, 2019
NOVOSIBIRSK, RUSSIA
CONF.BIONET.NSC.RU/PLANTGEN2019/EN

Institute of Cytology and Genetics, Siberian Branch of the Russian Academy of Sciences
Novosibirsk State University

Alexey Kochetov · Elena Salina
Editors

Current Challenges in Plant Genetics, Genomics, Bioinformatics, and Biotechnology

Proceedings of the Fifth International
Scientific Conference PlantGen2019
June 24–29, 2019 Novosibirsk, Russia

conf.bionet.nsc.ru/plantgen2019/en



Novosibirsk
ICG SB RAS
2019

Current Challenges in Plant Genetics, Genomics, Bioinformatics, and Biotechnology:
Proceedings of the Fifth International Scientific Conference PlantGen2019 (June 24–29,
2019, Novosibirsk, Russia) / Eds. A. Kochetov, E. Salina / Institute of Cytology and
Genetics, Siberian Branch of the Russian Academy of Sciences. – Novosibirsk: ICG
SB RAS, 2019. – 252 pp. – ISBN 978-5-91291-048-7.

DOI 10.18699/ICG-PlantGen2019-82

Editors

Prof. *Alexey Kochetov*
Institute of Cytology and Genetics,
Siberian Branch of the Russian
Academy of Sciences
630090 Prospekt Lavrentyeva, 10
Novosibirsk, Russian Federation
e-mail: ak@bionet.nsc.ru

Prof. *Elena Salina*
Institute of Cytology and Genetics,
Siberian Branch of the Russian
Academy of Sciences
630090 Prospekt Lavrentyeva, 10
Novosibirsk, Russian Federation
e-mail: salina@bionet.nsc.ru

Additional Reviewers

Alexey Kochetov, Dr. Sci. (Biology), Professor, Corr. Member of the RAS

Elena Salina, Dr. Sci. (Biology), Professor

Irina Leonova, Dr. Sci. (Biology)

Dmitry Afonnikov, Cand. Sci. (Biology)

Contacts

E-mail: PlantGen2019@bionet.nsc.ru
Web: conf.bionet.nsc.ru/plantgen2019/en

Organizers

Institute of Cytology and Genetics SB RAS (ICG SB RAS), Novosibirsk, Russia

Novosibirsk State University (NSU), Novosibirsk, Russia

Ministry of Science and Higher Education of the Russian Federation, Moscow, Russia

The Vavilov Society of Geneticists and Breeders (VSGB), St. Petersburg, Russia

The Vavilov Society of Geneticists and Breeders (Novosibirsk Branch), Novosibirsk, Russia

EUCARPIA (European Association for Research on Plant Breeding), Wageningen, Netherlands

International Program Committee

Alexey Kochetov, Institute of Cytology and Genetics, SB RAS, Novosibirsk, Russia
(Conference Chair)

Elena Salina, Institute of Cytology and Genetics, SB RAS, Novosibirsk, Russia
(Conference Co-Chair)

Vladimir Shumny, Institute of Cytology and Genetics, SB RAS, Novosibirsk, Russia

Nikolay Kolchanov, Institute of Cytology and Genetics, SB RAS, Novosibirsk, Russia

Andreas Börner, Leibniz Institute of Plant Genetics and Crop Plant Research, Gatersleben,
Germany

Yerlan Turuspekov, Institute of Plant Biology and Biotechnology, Almaty, Kazakhstan

Konstantin V. Krutovsky, Georg August University of Göttingen, Göttingen, Germany

Olga Afanasenko, All-Russian Institute of Plant Protection, St. Petersburg, Russia

Philipp B. Gannibal, All-Russian Institute of Plant Protection, St. Petersburg, Russia

Tatjana Gavrilenko, Federal Research Center the N.I. Vavilov All-Russian Institute of Plant
Genetic Resources, St. Petersburg, Russia

Elena Khlestkina, Federal Research Center the N.I. Vavilov All-Russian Institute of Plant
Genetic Resources, St. Petersburg, Russia

Igor Tikhonovich, All-Russian Research Institute for Agricultural Microbiology,
St. Petersburg, Russia

Gennady Karlov, All-Russian Research Institute of Agricultural Biotechnology, Moscow,
Russia

Alexander Kudriavtsev, Vavilov Institute of General Genetics, RAS, Moscow, Russia

Elena Kochieva, Federal Research Center Fundamentals of Biotechnology, RAS, Moscow,
Russia

Nikolai Kashevarov, Siberian Federal Scientific Center of Agrobiotechnologies, RAS,
Krasnoobsk, Novosibirsk region, Russia

Alexander Zhuchenko, All-Russian Breeding and Technological Institute of Horticulture
and Nursery, Moscow, Russia

Vladimir Kosolapov, Federal Williams Research Center of Forage Production
and Agroecology, Lobnya, Moscow region, Russia

Nikolay Goncharov, Institute of Cytology and Genetics, SB RAS, Novosibirsk, Russia

Contents

| | |
|--|----|
| PlantGen2019 | 9 |
| Plant Genetic Resources for Breeding and Producing Functional Nutraceutical Food | |
| <hr/> | |
| Introduction in Siberia (Russia) of new vegetable species with a high biochemical value <i>Y.V. Fotev, A.I. Syso, O.M. Shevchuk</i> | 12 |
| Anthocyanins in wheat grain: genetic control, health benefit and bread-making quality <i>E.K. Khlestkina, O.Y. Shoeva, E.I. Gordeeva, Y.S. Otmakhova, N.I. Usenko, M.A. Tikhonova, M.V. Tenditnik, T.G. Amstislavskaya</i> | 15 |
| Identification of wheat varieties with high grain protein and gluten content <i>D-r.F. Askhadullin, D-l.F. Askhadullin, N.Z. Vasilova, E.V. Zuev, I.I. Khusainova, M.R. Tazutdinova, E.Z. Bagavieva</i> | 19 |
| Wheat genetic resources for breeding on adaptability <i>M.A. Yessimbekova, M.Sh. Suleimenova, K.B. Mukin</i> | 21 |
| Development of new forms of spring and winter wheat with the involvement of the gene pool of wheatgrass and soybean and confirmation of the applicability of these methods in practical breeding <i>E.P. Razmakhnin, T.M. Razmakhnina, N.I. Steepochkina, V.I. Ponomarenko, V.I. Kozlov, K.K. Musinov, A.A. Surnachev, G.V. Artemova, I.E. Likhenko, G.Y. Galycin, N.P. Goncharov</i> | 24 |
| Breeding of soft spring wheat in a changing climate <i>V.V. Novokhatin</i> | 28 |
| Spring common wheat breeding lines produced on the basis of distant hybridization: ecological strain testing in Bagan <i>B.F. Nemtsev, A.B. Nemtsev, N.P. Goncharov, S.V. Kurkova</i> | 30 |
| Creation and study of partial waxy wheat lines adapted for the Middle Volga region of Russia <i>D-l.F. Askhadullin, D-r.F. Askhadullin, N.Z. Vasilova, R.R. Vafin</i> | 34 |
| Internal stem features of spring wheat varieties as factors determining resistance to lodging <i>E.V. Ageeva, I.N. Leonova, E.A. Salina, I.E. Likhenko</i> | 36 |
| Application of ISSR markers for study of genetic polymorphism of dark grain maize varieties <i>L.S. Valiyeva, G.K. Rahimova, N.A. Nabiyeva</i> | 39 |
| Genetic structure of the ampelographic collection maintained in the Dagestan experimental station of VIR revealed by microsatellite analysis <i>M.M. Agakhanov, V.A. Volkov, E.K. Potokina</i> | 42 |
| Genetic variation of water caltrop (<i>Trapa L.</i>) in several Russian populations <i>A.E. Artyukhin, E.V. Mikhaylova, B.R. Kuluev</i> | 44 |
| Potato starch resistivity as an inheritable trait <i>L.M. Gvozdeva, I.V. Rozanova, V.K. Khlestkin, E.K. Khlestkina</i> | 46 |
| Evaluation of genetic variability of buckwheat varieties (<i>Fagopyrum esculentum</i>) using ISSR-analysis <i>A.G. Klykov, P.V. Fisenko, A.S. Chibizova, E.N. Barsukova</i> | 49 |

Plant Resistance to Pathogens and Other Biotic Stresses

| | |
|--|----|
| Resistance to stem rust and a molecular marker analysis of <i>Sr</i> genes in spring bread wheat introgression lines <i>O.A. Baranova, S.N. Sibikeev, A.E. Druzhin</i> | 52 |
| Development of the panel of SSR markers to estimate the level of genetic diversity among the Siberian stem rust population <i>V.N. Kelbin, E.S. Skolotneva, S. Vidich, M.A. Nesterov, E.M. Sergeeva, E.A. Salina</i> | 56 |
| Study of introgression lines of common wheat <i>Triticum aestivum</i> / <i>Triticum miguschovae</i> for resistance to leaf rust <i>R.O. Davoyan, I.V. Bebykina, E.R. Davoyan, D.S. Mikov, Y.S. Zubanova, D.M. Boldakov, A.N. Zinchenco</i> | 58 |
| Development and study of new spring wheat lines containing alien genetic material from <i>Th. intermedium</i> and <i>Ae. speltoides</i> <i>I.N. Leonova, V.V. Piskarev, N.I. Boiko, A.I. Stasyuk, I.G. Adonina, E.A. Salina</i> | 61 |
| Interaction of the <i>Tsn1</i> and <i>Tsc2</i> sensitivity genes with the <i>ToxA</i> and <i>ToxB</i> effector genes in <i>Triticum</i> L. species <i>N.V. Mironenko, O.A. Baranova, N.M. Kovalenko, O.P. Mitrofanova</i> | 64 |
| Molecular screening of wheat entries for resistance to the toxins Ptr ToxA and Ptr ToxB from the tan spot pathogen <i>Pyrenophora tritici-repentis</i> <i>A.M. Kokhmetova, S. Ali, M.N. Atishova</i> | 66 |
| Functional characterization of <i>Stagonospora nodorum</i> necrotrophic effectors <i>S.V. Veselova, G.F. Burkhanova, T.V. Nuzhnaya, I.V. Maksimov</i> | 69 |
| <i>Septoria</i> blight of spring wheat in West Siberia <i>E.Yu. Toropova, O.A. Kazakova, V.V. Piskarev</i> | 72 |
| Race composition of the loose smut (<i>Ustilago tritici</i>) in Western Siberia <i>E.A. Orlova, N.P. Bechtold</i> | 74 |
| Potential of ribonuclease-synthesizing plant growth promoting rhizobacteria in plant defence against viruses <i>G.F. Burkhanova, A.V. Sorokan, V.U. Alekseev, A.A. Mazitova, I.V. Maksimov</i> | 77 |
| Studying the infection rate of seeds of various genotypes of sweet sorghum (<i>Sorghum bicolor</i> L.) by fungus and bacterial microflora <i>K.M. Iskakova, B.B. Anapiyayev, E.B. Beisenbek, A.B. Akhmetova, A.M. Sagimbayeva, A.S. Omarova</i> | 80 |
| Role of genes involved in lignin biosynthesis in flax response to <i>Fusarium oxysporum</i> <i>R.O. Novakovskiy, G.S. Krasnov, T.A. Rozhmina, L.P. Kudryavtseva, A.A. Zhuchenko, E.N. Pushkova, L.V. Povkhova, P. Kezimana, A.A. Dmitriev, N.V. Melnikova</i> | 83 |

Genetic and Epigenetic Mechanisms of Plant Resistance to Abiotic Stresses

| | |
|---|----|
| Epigenetic mechanisms of the karyogenomic system of mature wheat germ derived under conditions of cold stress <i>E.A. Ivanova, G.Kh. Vafina</i> | 87 |
| Comparative characteristic of near-isogenic lines differing by the <i>Blp</i> locus in respect to abiotic stress resistance <i>A.Y. Glagoleva, O.Y. Shoeva, E.K. Khlestkina</i> | 89 |
| Creation of mutant populations of barley (<i>Hordeum vulgare</i> L.) and flax (<i>Linum ussitatissimum</i> L.) induced by the chemical mutagen of the phosphemide <i>N.A. Bome, K.P. Korolev, N.V. Tetyannikov, L.I. Weisfeld, N.N. Kolokolova</i> | 92 |
| Functional characterization of drought resistance in peach (<i>Prunus pertsica</i> (L.) Batsch) cultivars damaged with Sharka disease <i>V.A. Brailko, I.V. Mitrofanova, N.V. Mesyats, O.V. Mitrofanova, S.V. Chirkov</i> | 95 |
| Response of the antioxidant system of two varieties of potatoes to salt stress <i>O.K. Murgan, L.V. Kolomeichuk, I.S. Kovtun, M.V. Efimova</i> | 98 |

| | |
|--|-----|
| Mechanism of plant adaptation to changing illumination by rearrangements of their photosynthetic apparatus <i>V.I. Belkov, E.Yu. Garnik, Yu.M. Konstantinov</i> | 101 |
| Features of the interaction of miRNAs with genes of the rice MYB family under stress <i>A.T. Ivashchenko, A.K. Rakhmetullina, A.U. Pyrkova</i> | 104 |
| Sensitive biomarker for assessing the effects of the environment on the population of plants <i>L.A. Minasbekyan, G.S. Aidarkhanova, I.A. Avagyan</i> | 107 |
| Selenium nanocomposites as a protection agent from potato ring rot <i>O.A. Nozhkina, A.I. Perfileva, I.A. Graskova, B.G. Sukhov</i> | 110 |

Plant Biotechnology in the Postgenomic Era

| | |
|---|-----|
| Modern CRISPR/Cas9 biotechnology is a current challenge to create elite disease resistant crops cultivars <i>O.I. Kershanskaya, D.S. Nelidova, G.L. Esenbaeva, G.S. Mukiyanova, S.N. Nelidov</i> | 113 |
| Study of a region on wheat chromosome 5BS by targeted sequencing <i>M.K. Bragina, D.A. Afonnikov, G.V. Vasiliev, E.A. Salina</i> | 117 |
| Intergenomic substitutions and translocations in wheat-alien lines and their use in studying the characteristics that determine adaptation and stress resistance <i>T.T. Efremova, E.V. Chumanova</i> | 120 |
| Copy number variation of the <i>Vrn-1b</i> allele in emmer and spelt wheat <i>A. Muterko</i> | 124 |
| Evaluation of <i>in vitro</i> plant regeneration efficiency in Siberian wheat cultivars <i>D.N. Miroshnichenko, A.A. Klementyeva, E.A. Salina, S.V. Dolgov</i> | 126 |
| Improvement of sorghum grain quality using modern genetic tools <i>L.A. Elkonin, V.M. Panin, G.A. Gerashchenkov, O.A. Kenzhegulov</i> | 129 |
| New efficient gene promoters from <i>Stellaria media</i> for plant genetic engineering <i>L.N. Efremova, S.R. Strelnikova, R.A. Komakhin</i> | 133 |
| Role of the expansin and xyloglucan endotransglycosylase genes in the regulation of tobacco growth under the influence of abiotic stress factors <i>B.R. Kuluev, Kh.G. Musin, Z.A. Berezhneva, E.V. Mikhaylova, E.A. Zaikina</i> | 135 |
| Reproduction and genetic accuracy during somatic embryogenesis in <i>Larix sibirica</i> <i>I.N. Tretyakova, M.E. Park, N.V. Oreshkova, D.V. Kulagin, A.V. Konstantinov, V.E. Padutov</i> | 137 |
| Reactivation of <i>VaSTS1</i> expression in <i>Arabidopsis thaliana</i> transgenic plants by retransformation with <i>2b</i> from the <i>Cucumber Mosaic Virus</i> isolate NK <i>N.N. Nityagovsky, A.P. Tyunin, K.V. Kiselev</i> | 140 |
| Development of a new method for eradication of viruses by induction of posttranscriptional gene silencing in transgenic potato plants <i>A.M. Alexandrova, O.V. Karpova, R.M. Nargilova, R.V. Kryldakov, B.K. Iskakov</i> | 143 |
| Up-regulation of the vacuolar invertase TAI gene may contribute to the accumulation of carotenoids in tomato fruits <i>M.A. Slugina, E.A. Dzhos, A.V. Shchennikova, E.Z. Kochieva</i> | 146 |
| Induction of <i>in vitro</i> propagation in mature embryos of maize <i>B.M.H. Humood, O.I. Yudakova</i> | 150 |
| Preparation of DNA libraries for sequencing of <i>SAD</i> and <i>FAD</i> genes in flax <i>P. Kezimana, T.A. Rozhmina, G.S. Krasnov, R.O. Novakovskiy, L.V. Povkhova, E.N. Pushkova, M. Pavelek, E.V. Romanova, A.A. Dmitriev, N.V. Melnikova</i> | 153 |
| Rare species <i>Iris scariosa</i> and <i>I. pumila</i> in the Southern Urals <i>E.V. Mikhaylova, A.N. Mustafina, Y.M. Golovanov, A.V. Kryukova</i> | 156 |
| Extraction of high-molecular-weight DNA from poplar plants for Nanopore sequencing <i>E.N. Pushkova, A.D. Beniaminov, G.S. Krasnov, R.O. Novakovskiy, L.V. Povkhova, N.V. Melnikova, A.A. Dmitriev</i> | 158 |

Plant Systems Biology and Digital Technologies

| | |
|---|-----|
| Novel genomic marker for the Alm locus in barley identified based on transcriptome analysis <i>N.A. Shmakov, A.Yu. Glagoleva, G.V. Vasiliev, D.A. Afonnikov, E.K. Khlestkina</i> | 162 |
| Comparative transcriptome analysis of <i>Syringa vulgaris</i> vegetative apices <i>in vivo</i> and <i>in vitro</i> <i>A.A. Krinitsina, A.S. Speranskaya, O.A. Churikova, M.D. Logacheva, E.A. Konorov, M.S. Belenikin</i> | 166 |
| Overexpression of RPOTmp, an Arabidopsis RNA-polymerase with dual-targeting, influences plant growth and transcriptome <i>V.I. Tarasenko, A.I. Katyshev, I.V. Gorbenko, V.I. Belkov, E.Y. Garnik, Y.M. Konstantinov, M.V. Koulintchenko</i> | 169 |
| <i>NDH</i> gene disruption – possible implication for adaptation of <i>Allium</i> species to different light conditions <i>A.S. Speranskaya, M.S. Belenikin, E.A. Konorov, S.V. Kuptsov, M.I. Antipin, M.D. Logacheva, D.O. Omelchenko, A.A. Krinitsina</i> | 173 |
| Gene expression in Scots pine populations from the Chernobyl exclusion zone: transcriptomic approach <i>P.Yu. Volkova, G.T. Duarte, S.A. Geras'kin</i> | 177 |
| Use of deep sequencing to study the genetic diversity of flax pathogens <i>R.O. Novakovskiy, G.S. Krasnov, E.N. Pushkova, L.P. Kudryavtseva, T.A. Rozhmina, P. Kezimana, L.V. Povkhova, N.V. Melnikova, A.A. Dmitriev</i> | 179 |
| Computer program for construction of regression function for phenotype in agroclimatic models with interactions <i>K.N. Kozlov, S.V. Nuzhdin, M.G. Samsonova</i> | 181 |
| Bioninformatics research of CRISPR/Cas-systems in the genomes of the phytopatogenic strain <i>Agrobacterium fabrum</i> C58 <i>I.A. Portnaia, A.Yu. Borisenko, Yu.P. Dzhioev, O.G. Karnoukhova, Ju.A. Markova, L.A. Stepanenko, O.N. Reva, N.P. Peretolchina, V.I. Zlobin</i> | 184 |

Plant Breeding in the 21st Century

| | |
|--|-----|
| Main directions of the spring bread wheat breeding in Western Siberia <i>I.A. Belan, L.P. Rosseeva, N.P. Blokhina, L.F. Lozhnikova, V.V. Nemchenko, S.N. Abakumov, R.K. Kadikov, N.V. Trubacheeva, L.A. Pershina</i> | 187 |
| Improved marker for the <i>Rht-Blp</i> dwarfing allele in wheat <i>M.S. Bazhenov, L.A. Nazarova, A.G. Chernook, M.G. Divashuk</i> | 190 |
| Marker-assisted selection and yield component assessment of spring wheat plants derived from crosses between spring and winter wheat cultivars <i>A.I. Stasyuk, I.N. Leonova, E.A. Salina</i> | 193 |
| Transferability of barley EST markers used for analysis <i>T. aestivum</i> – <i>H. marinum</i> subsp. <i>gussoneanum</i> introgression lines <i>N.V. Trubacheeva, E.D. Badaeva, T.S. Osadchaya, L.A. Pershina</i> | 196 |
| Study of introgression lines of common wheat obtained with the participation of the synthetic forms <i>T. miguschovae</i> and Avrodes by allelic variants of the <i>Wx</i> genes <i>E.R. Davoyan, R.O. Davoyan, Yu.S. Zubanova, D.S. Mikov, D.M. Boldakov, A.G. Chernook, M.G. Divashuk</i> | 199 |
| <i>Non-brittle rachis 1-A (Btr1-A)</i> gene in di- and hexaploid wheat species <i>V. Vavilova, I. Konopatskaia, N.P. Goncharov</i> | 202 |
| Development of androgenic polyembryoids in the anther culture of alloplasmic introgression lines of (<i>H. vulgare</i>) – <i>T. aestivum</i> and (<i>H. marinum</i> ssp. <i>gussoneanum</i>) – <i>T. aestivum</i> <i>L.A. Pershina, N.V. Trubacheeva, T.S. Osadchaya, I.A. Belan, L.P. Rosseeva</i> | 205 |
| Challenges and opportunities of breeding and genetic improvement of durum wheat in Russia <i>N.P. Mal'chikov, M.A. Rozova, A.I. Ziborov, M.G. Myasnikova, T.V. Chakheeva</i> | 207 |

| | |
|--|---------------------|
| Study of 8x and 6x triticale with dominant <i>Vrn</i> genes <i>P.I. Steepochkin</i> | 211 |
| Breeding of potato resistant to late blight using genetic resources and DNA markers <i>J. Śliwka, M. Brylińska, E. Stefańczyk, J. Plich, P. Smyda-Dajmund, S. Sobkowiak</i> | 214 |
| Search for genomic regions associated with potato starch granules morphology of <i>Solanum tuberosum</i> L. <i>T.V. Erst, I.V. Rozanova, V.K. Khlestkin, E.K. Khlestkina</i> | 217 |
| Comprehensive evaluation of the Siberian gene pool of common bean (<i>Phaseolus vulgaris</i> L.) in the conditions of Western Siberia <i>O.E. Yakubenko, O.V. Parkina</i> | 219 |
| Peculiarities of heterosis manifested by yield and fruit quality traits in pepper F ₁ hybrids developed using classical and MAS methods <i>O.G. Babak, T.V. Nikitinskaya, N.A. Nevestenko, I.G. Puhacheva, M.M. Dobrodin, L.V. Khotyleva, A.V. Kilchevsky</i> | 221 |
| Key directions of breeding and seed production of alfalfa in European Russia <i>V.I. Cherniavskih, E.V. Dumacheva, Z.A. Borodaeva, D.V. Dumachev</i> | 224 |

GWAS for Dissecting Traits and Pinpoint Candidate Genes

| | |
|---|---------------------|
| High-throughput technologies for sunflower oil improvement <i>A.I. Chernova, E.U. Martynova</i> | 227 |
| High-throughput genotyping for association mapping of agronomically important traits in rapeseed: a brief review of the current status <i>R.F. Gubaev, S.V. Boldyrev, D.V. Goryunov, S.V. Goryunova</i> | 230 |
| Genomic analysis in soybean breeding <i>V.V. Barzanova, A.A. Novikova</i> | 232 |

Plant Cytogenetics in the Genomic and Postgenomic Era

| | |
|--|---------------------|
| Molecular cytological analysis of alien introgressions in common wheat lines with <i>T. kiharae</i> genetic material <i>O.A. Orlovskaya, N.I. Dubovets, L.A. Solovey, E.B. Bondarevich, I.N. Leonova</i> | 235 |
| Expression profile of two <i>CENH3</i> genes in different tissues of the <i>Secale cereale</i> <i>S.S. Gatzkaya, E.V. Evtushenko, A.V. Vershinin</i> | 238 |
| Chromosome constitution and activity of the centromeric histone H3 (<i>CENH3</i>) variants in octoploid triticale lines <i>E.V. Evtushenko, Yu.A. Lipikhina, P.I. Steepochkin, A.V. Vershinin</i> | 241 |
| Karyotyping of triticale-wheat hybrid lines with the sham ramification of spike <i>I.G. Adonina, S.P. Mehdiyeva, M.V. Prokopjeva, N.Kh. Aminov, E.A. Salina</i> | 244 |
| Identification of univalent chromosomes in monosomic lines of the <i>Gossypium hirsutum</i> L. cotton type with the help of cytogenetic markers <i>M.F. Sanamyan, Sh.U. Bobokhujayev</i> | 247 |
| Author index | 250 |

Features of the interaction of miRNAs with genes of the rice MYB family under stress

A.T. Ivashchenko*, A.K. Rakhmetullina, A.U. Pyrkova

Al-Farabi Kazakh National University, Almaty, Kazakhstan

DOI 10.18699/ICG-PlantGen2019-33

© Autors, 2019

* e-mail: a.iavashchenko@gmail.com

Abstract: Data on the involvement of miRNA in the plant response to stress factors are considered. The MYB family of transcription factors plays a great role in the plant development, metabolism and responses to biotic and abiotic stress. The free energy of miRNA binding, the value of free energy of interaction, the position and schemes of potential binding sites were calculated using the MirTarget program. To identify miRNAs whose targets are genes in the MYB family, a search for binding sites of 738 miRNAs in mRNAs of 124 MYB family genes of *Oryza sativa* was performed. 16 genes were identified as targets for 11 miRNAs. The results show that miRNA can regulate the expression of most MYB genes and thus affect productivity and sustainability.

Key words: miRNA; mRNA; genes; transcription factors; rice; stress.

1. Introduction

Rice is one of the main food crops. According to the Food and Agriculture Organization of the United Nations, rice is the staple food for more than half of the world's population. MYB proteins constitute one of the largest transcription factor families in the plant kingdom, members of which are key factors in regulatory networks controlling development, metabolism and responses to biotic and abiotic stresses in the plant genome (Shushi et al., 2015). In agricultural production, abiotic stresses are known as the main causes leading to lower yields (Hoang et al., 2017). The plants have developed complex mechanisms for overcoming various stresses (Jian et al., 2010). Recent evidence suggests that miRNAs are involved in the regulation of gene expression, which affects the post-transcriptional stage of gene expression and relates to biotic and abiotic stress reactions in plants (Bari et al., 2014). Stress in plants causes an increase or decrease in the expression of certain miRNAs or the synthesis of new miRNAs (Varsha et al., 2016). Several stress-regulated miRNAs have been found in plants under various biotic and abiotic stressful conditions, including nutrient deficiency (Fujii et al., 2005), drought (Zhao et al., 2007; Liu et al., 2008; Zhou et al., 2010), low temperatures (Zho et al., 2008), salinity (Sunkar et al., 2008), bacterial infection (Navarro et al., 2006), UV-B radiation (Zhou et al., 2007), and mechanical stress (Lu et al., 2005). These studies suggest that miRNA is regulated in various ways in response to stress. However, limited studies have reported an association of miRNA with the expression of plant MYB genes. In the present work, the task was set to identify, using bioinformatics approaches, miRNAs that can bind to mRNA genes of the MYB family of rice and regulate their expression.

2. Materials and methods

The object of the study was the completely sequenced *O. sativa* genome. The nucleotide sequences of the mRNA genes of the MYB family were obtained from PlantTFDB, a plant transcription factor database (planttfdb.cbi.pku.edu.cn/). The miRNA nucleotide sequences were borrowed from miRBase

(<http://www.mirbase.org/>). The free energy (ΔG) of miRNA binding, the position and schemes of potential binding sites were calculated using the MirTarget program (Ivashchenko et al., 2016). ΔG_m for miRNA was defined as the free energy of miRNA binding with its fully complementary nucleotide sequence. The miRNA binding sites with mRNA were selected with the $\Delta G/\Delta G_m$ ratio being more than 90 %. Unique features of the MirTarget program include the consideration of the nucleotide interaction in miRNA with the mRNA of target genes not only between adenine (A) and uracil (U), guanine (G) and cytosine (C), G-U, but also between A and C through a single hydrogen bond based on the fact that the distance between A and C is equal to the distance between the nucleotides G-C, A-U, G-U (Leontis et al., 2002; Kool et al., 2001).

3. Results and discussion

The study of the binding of 738 miRNA to mRNA of 124 genes of the MYB *O. sativa* family revealed that only 16 genes were targets for 11 miRNA (Table 1).

Four miRNA of the miR159c,d,e,f-3p family had the largest number of binding sites which were associated with mRNA genes (LOC_Os06g40330.1, LOC_Os01g59660.1, LOC_Os05g41166.1, LOC_Os03g38210.1, LOC_Os04g46384.1) of the MYB family with the $\Delta G/\Delta G_m$ value from 90 to 96 %. Therefore, miR159 involved in the response to stress (Varsha et al., 2016) can affect the expression of the rice genes studied. miR2102-5p was binding with mRNA genes (LOC_Os03g26130.1, LOC_Os01g64360.1, LOC_Os01g62410.1, LOC_Os03g25550.1) with the energy from -110 kJ/mole to -113 kJ/mole and the degree of complementarity ($\Delta G/\Delta G_m$) 91 % and 93 %. miR5075-3p had two binding sites in the mRNA of MYB genes. miR171d-5p, miR172d-5p; miR5827-5p had only one target gene (LOC_Os03g27090.1, LOC_Os04g42950.1, LOC_Os06g02250.1, LOC_Os06g46560.1, LOC_Os01g18240.1) with the value of $\Delta G/\Delta G_m$ from 91 to 93 %. The miRNA binding sites in the mRNA genes of the MYB *O. sativa* family were located in the protein coding region.

- Kool E.T. Hydrogen bonding, base stacking, and steric effects in DNA replication. *Annu Rev Biophys Biomol Struct.* 2001;30:1–22. DOI 10.1146/annurev.biophys.30.1.1.
- Leontis N.B., Stombaugh J., Westhof E. The non-Watson-Crick base pairs and their associated isostericity matrices. *Nucleic Acids Res.* 2002;30(16):3497–3531. DOI 10.1093/nar/gkf481.
- Liu H.H., Tian X., Li Y.J., Wu C.A., Zheng C.C. Microarray-based analysis of stress-regulated microRNAs in *Arabidopsis thaliana*. *RNA.* 2008;14(5):836–843. DOI 10.1261/rna.895308.
- Lu S., Sun Y.H., Shi R., Clark C., Li L., Chiang V.L. Novel and mechanical stress-responsive microRNAs in *Populus trichocarpa* that are absent from *Arabidopsis*. *Plant Cell.* 2005;17(8):2186–2203. DOI 10.1105/tpc.105.033456.
- Navarro L., Dunoyer P., Jay F., Arnold B., Dharmasiri N., Estelle M., Voinnet O., Jones J.D. A plant miRNA contributes to antibacterial resistance by repressing auxin signaling. *Sci.* 2006;312(5772):436–439. DOI 10.1126/science.1126088.
- Shushi S., Amit K., Viswanathan Ch., Dev M.P., Kailash C.B. Transcriptional Regulatory Network Analysis of MYB Transcription Factor Family Genes in Rice. *Front Plant Sci.* 2015;6:1157. DOI 10.3389/fpls.2015.01157.
- Sunkar R., Zhou X., Zheng Y., Zhang W., Zhu J.-K., Identification of novel and candidate miRNAs in rice by high throughput sequencing. *BMC Plant Biol.* 2008;29(8):25. DOI 10.1186/1471-2229-8-25.
- Zhao B., Liang R., Ge L., Li W., Xiao H., Lin H., Ruan K., Jin Y. Identification of drought-induced microRNAs in rice. *Biochem. Biophys. Res. Commun.* 2007;354(2):585–590. DOI 10.1016/j.bbrc.2007.01.022
- Zhou L., Liu Y., Liu Z., Kong D., Duan M., Luo L. Genome-wide identification and analysis of drought-responsive microRNAs in *Oryza sativa*. *J. Exp. Bot.* 2010;61(15):4157–4168. DOI 10.1093/jxb/erq237.
- Zhou X., Wang G., Zhang W. UV-B responsive microRNA genes in *Arabidopsis thaliana*. *Mol. Syst. Biol.* 2007;3:103. DOI 10.1038/msb4100143.
- Zhou X., Wang G., Sutoh K., Zhu J.-K., Zhang W. Identification of cold-inducible microRNAs in plants by transcriptome analysis. *Biochim. Biophys. Acta.* 2008;1779(11):780–788. DOI 10.1016/j.bbagr.2008.04.005.

Conflict of interest. The authors declare no conflict of interest.