








21st International Symposium on
HEAVY ION FUSION
Astana, Kazakhstan • July 18-22, 2016

Book of Abstracts



LASER-INDUCED-FLUORESCENCE MEASUREMENT OF THERMAL CONDUCTIVITY IN WARM DENSE MATTER GENERATED BY PULSED-POWER DISCHARGE.....	81
T. Sasaki, S. Sugimoto, Y. Sugimoto, A. Watabe, Sh. Kusano, K. Takahashi, T. Kikuchi, N. Harada	
ENERGY YIELD OF THERMONUCLEAR FUEL.....	82
Zh.B.Seksembayev, O.B.Bayakhmetov, S.K.Sakhiyev	
REFLECTIVITY OF THE DENSE XENON PLASMA.....	83
E.O. Shalenov, K.N. Dzhumagulova, T.S. Ramazanov, G. Röpke, H. Reinholz	
ION CURRENT DENSITY PROFILE OF LASER ABLATION PLASMA TRANSPORTED IN MULTICUSP MAGNETIC FIELD.....	84
K. Takahashi, T. Uchino, K. Ikegami, T. Sasaki, Takashi Kikuchi, N. Harada	
DESIGN OF VELOCITY MEASUREMENT SYSTEM TO GENERATE WARM DENSE MATTER OF INSULATOR BY FLYER IMPACT.....	85
F. Tamura, N. Takakura, Yu. Nakayama, K. Kashine, A. Tokuchi, T. Kikuchi, K. Takahashi, T. Sasaki, N. Harada, W. Jiang, Sh. Fujioka, T. Sano	
STUDY OF PEELING OF SINGLE CRYSTAL SILICON BY INTENSE PULSED ION BEAM.....	86
X. Yu, J. Shen, J. Zhang, H. Zhong, X. Cui, G. Liang, Sh. Yan, G. Zhang, X. Zhang, X. Le	
ACCELERATOR COMPLEX BASED ON DC-60 CYCLOTRON.....	87
M. Zdorovets, I. Ivanov, M. Koloberdin, S. Kozin, V. Alexandrenko, E. Sambaev, A. Kurakhmedov	
STUDY ON ABLATION PRODUCTS OF ZINC BY INTENSE PULSED ION BEAM IRRADIATION.....	88
J. Zhang, H. Zhong, Zh. Ye, J. Shen, X. Yu, G. Liang, X. Cui, X. Zhang, G. Zhang, Sh. Yan, X. Le	
X-RAY EMISSION FOR 424 MEV/U C IONS IMPACTING ON SELECTED TARGETS.....	89
X. Zhou, R. Cheng, Yu. Wang, Yu Lei, J. Ren, Sh. Liu, Y. Zhao	
THE CALCULATION OF PARAMETERS AND DESIGN OF «PLASMA FOCUS» FACILITY.....	90
A.M. Zhukeshov, B.M. Useinov, A.T.Gabdullina, A.U. Amrenova, K. Serik	
INVESTIGATION OF DIFFUSION PROCESSES AT METAL TARGET IRRADIATION BY PULSED PLASMA FLOW.....	91
A. M. Zhukeshov, Z. Mazhit	

Reflectivity of the dense xenon plasma

E.O. Shalenov^{*1}, K.N. Dzhumagulova¹, T.S. Ramazanov¹, G. Röpke², H. Reinholz²

¹IETP, Department of Physics, al-Farabi Kazakh National University, al-Farbi 71, 050040Almaty, Kazakhstan

²Institute of Physics, University of Rostock, A.-Einstein-Str. 23-24, 18059Rostock, Germany

^{*}shalenov.erik@mail.ru

The investigation of optical properties of the dense xenon plasma is important for the realization of different technological applications [1-2].

In this work we consider the dense partially ionized xenon plasma consisting of the electrons, ions and atoms. Particle densities are in the range of $n = 10^{20} + 10^{23} \text{ cm}^{-3}$ and the temperature range is from $2.5 \times 10^4 \text{ K}$ up to $5 \times 10^4 \text{ K}$.

In work [3] the effective potential of electron-atom interaction, taking into account both quantum-mechanical effect of diffraction and screening effects, is presented. The way to take into account the dynamic screening was proposed in work [4], where the statical Debye radius was replaced by the dynamic one:

$$r_o = r_D \left(1 + \frac{v^2}{v_{Th}^2}\right)^{1/2}, \quad (1)$$

here v is the relative velocity of the colliding particles, v_{Th} is the thermal velocity of the particles in the system. Then the effective potential of electron-atom interaction with dynamic screening can be rewritten as [5]:

$$\Phi_{ea}^{dyn}(r) = -\frac{e^2 \alpha}{2r^4 (1 - 4\lambda_{ea}^2 / r_o^2)} \left(e^{-Br} (1 + Br) - e^{-Ar} (1 + Ar) \right)^2, \quad (2)$$

where $A^2 = \frac{1}{2\lambda_{ea}^2} \left(1 + \sqrt{1 - 4\lambda_{ea}^2 / r_o^2} \right),$

$$B^2 = \frac{1}{2\lambda_{ea}^2} \left(1 - \sqrt{1 - 4\lambda_{ea}^2 / r_o^2} \right).$$

In the framework of these pseudopotential models for the particle interactions, the scattering phase shifts were calculated on the basis of the Calogero equation [6].

Phase shifts enable us to calculate the transport scattering cross section $Q_{ea}^T(k)$. The collision fre-

quency of electrons with atoms ν_{ea} can be obtained by the following expression:

$$\nu_{ea} = 4 \sqrt{\frac{2}{\pi}} n_a \sqrt{\frac{k_B T}{\mu_{ea}}} \int_0^\infty Q_{ea}^T(g) g^3 \text{Exp}(-g^2) dg, \quad (3)$$

here $\mu_{ea} = m_e m_a / (m_e + m_a)$ is the reduced mass of the electron-atom pair, g is dimensionless reduced velocity. The results will be compared with earlier results which were based on experiments for the transport cross section [7].

After that the dielectric function due to the electron atom interactions was calculated using the generalized Drude-Lorentz [8] equation:

$$\varepsilon(\omega) = 1 - \frac{\omega_p^2}{\omega^2 - i\nu_{ea}\omega}. \quad (4)$$

However, for the total electronic dielectric functions contributions due to electron-ion and electron-electron interactions will also be considered [8,9]. From the dielectric function the reflectivity of the semiclassical electron plasma is calculated, using the Fresnel formula, and is compared with experimental results [10].

References

- [1] L. Martinu *et al* 2000 *J. Vac. Sci. Technol. A* **18**, 2619–2645
- [2] S. Pongratz *et al* 1992 *J. Vac. Sci. Technol. A* **10**, 1897–1904
- [3] T.S. Ramazanov *et al* 2005 *Phys. Plasmas* **12**, 092702
- [4] D. Kremp *et al* 2005 *Springer*, p. 326
- [5] K.N. Dzhumagulova *et al* 2014 *Contr. Plasma Phys* **55**, 230-235
- [6] L.D. Landau *et al* 1989 *Nauka, Moscow*, p. 691
- [7] J.R. Adams *et al* 2007 *Phys. Rev. E* **76**, 035405
- [8] H. Reinholz 2005 *Ann. Phys. Fr.* **30**, 1
- [9] H. Reinholz *et al* 2012 *Phys. Rev. E* **85**, 036401
- [10] Yu.B. Zaporoghets *et al* 2016 *Contr. Plasma Phys*. in press