

E ISSN 1512-3391



ANDREA RAZMADZE MATHEMATICAL INSTITUTE
of Ivane Javakhishvili Tbilisi State University

REPORTS OF QUALITDE

Volume 3

Tbilisi, Georgia, 2024

International Workshop on the Qualitative Theory of Differential Equations – QUALITDE-2024
December 21-23, 2024

Andrea Razmadze Mathematical Institute of Ivane Javakhishvili Tbilisi State University
Tbilisi, Georgia

Program Committee: I. Kiguradze (Chairman) (Georgia), R. P. Agarwal (USA), R. Hakl (Czech Republic), N. A. Izobov (Belarus), S. Kharibegashvili (Georgia), T. Kiguradze (USA), T. Kusano (Japan), A. Ponosov (Norway), M. Tvrdý (Czech Republic)

Organizing Committee: N. Partsvania (Chairman), M. Japoshvili (Secretary), O. Jokhadze, M. Kvinikadze, Z. Sokhadze

•

REPORTS OF QUALITDE

Volume 3

Editor: Nino Partsvania

© Andrea Razmadze Mathematical Institute, Ivane Javakhishvili Tbilisi State University

Publisher: Tbilisi State University Press

On the Solvability of a Family of Nonlinear Two-Point Boundary Value Problems for Systems of Integro-Differential Equations and its Application

Svetlana Temesheva

Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan

E-mail: temeshevasvetlana@gmail.com

Perizat Abdimanapova

Almaty Technological University, Almaty, Kazakhstan

E-mails: peryzat74@gmail.com

1 Introduction

Conditions for the existence of solutions of systems of integro-differential equations and boundary-value problems for these systems are considered in [8, 9]. A general theory for these systems is developed and effective methods for solving them are proposed. In this paper, we study the x -parametric family of nonlinear boundary-value problems for integro-differential equations

$$\frac{\partial V}{\partial t} = f\left(x, t, \psi(t) + \int_0^x V(\xi, t) d\xi, V\right), \quad V \in \mathbb{R}^n, \quad (x, t) \in [0, \omega] \times [0, T], \quad (1.1)$$

$$g(x, V(x, 0), V(x, T)) = 0, \quad x \in [0, \omega], \quad (1.2)$$

where the functions $f : [0, \omega] \times [0, T] \times \mathbb{R}^{2n} \rightarrow \mathbb{R}^n$ and $g : [0, \omega] \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ are continuous.

Our study is based on the parametrization method [10, 12] proposed by Dzhumabaev. The parametrization method was developed to various boundary-value problems for some types of differential equations [2, 13, 15], such that Fredholm integro-differential equations [3–7, 17], delay differential equations [16], hyperbolic equations [1, 14] etc. The application of this method made it possible to derive the solvability conditions for the above problems.

This approach can also be applied to study the nonlinear nonlocal boundary value problem for the system of partial differential equations ($m = 1, 2, \dots$)

$$\begin{aligned} \frac{\partial^{m+1}u}{\partial t \partial x^m} &= f\left(x, t, \frac{\partial^{m-1}u}{\partial x^{m-1}}, \frac{\partial^m u}{\partial x^m}\right), \quad u \in \mathbb{R}^n, \quad (x, t) \in [0, \omega] \times (0, T), \\ \frac{\partial^k u}{\partial x^k} \Big|_{x=0} &= \psi_k(t), \quad t \in [0, T], \quad k = 0, 1, \dots, m-1, \\ g\left(x, \frac{\partial^m u(x, t)}{\partial x^m} \Big|_{t=0}, \frac{\partial^m u(x, t)}{\partial x^m} \Big|_{t=T}\right) &= 0. \end{aligned}$$

Therefore, the study of problem (1.1), (1.2) is of interest from the point of view of its application to nonlocal boundary value problems for a class of partial differential equations.

In the present paper, we propose a modified algorithm of the parametrization method for finding an isolated solution of problem (1.1), (1.2) and derive sufficient conditions for the existence of such a solution.

2 Setting of the problem and the main results

We consider the nonlinear nonlocal boundary value problem (1.1), (1.2) for the x -parametric family of integro-differential equations.

A solution of problem (1.1), (1.2) is a function $V(x, t) \in C([0, \omega] \times [0, T] \mathbb{R}^n)$, which is continuously differentiable on $[0, T]$ (at fixed $x \in [0, \omega]$) and satisfies the system of integro-differential equations (1.1) and the boundary conditions (1.2).

We take $h > 0$, $Nh = T$ ($N \in \mathbb{N}$), and make the partition $[0, \omega] \times [0, T] = \bigcup_{r=1}^N \Omega_r$, where $\Omega_r = [0, \omega] \times [(r - 1)h, rh]$, $r = \overline{1, N}$.

We will use the following notations.

- $C([0, \omega], \mathbb{R}^{n(N+1)})$ is the space of systems of functions $\lambda(x) = (\lambda_1(x), \lambda_2(x), \dots, \lambda_{N+1}(x))$ with the norm

$$\|\lambda\|_1 = \max_{x \in [0, \omega]} \max_{r=1, (N+1)} \|\lambda_r(x)\|,$$

here the functions $\lambda_r : [0, \omega] \rightarrow \mathbb{R}^n$ are continuous, $r = \overline{1, (N + 1)}$;

- $C([0, \omega] \times [0, T], \Omega_r, \mathbb{R}^{nN})$ is the space of systems of functions

$$V[x, t] = (V_1(x, t), V_2(x, t), \dots, V_N(x, t))$$

with the norm

$$\|V[\cdot]\|_2 = \max_{r=1, N} \max_{x \in [0, \omega]} \sup_{t \in [t_{r-1}, t_r]} \|V_r(x, t)\|,$$

where the functions $V_r(x, t) \in C(\Omega_r)$ have finite limits $\lim_{t \rightarrow t_r - 0} V_r(x, t)$ uniform in x , $x \in [0, \omega]$ ($r = \overline{1, N}$);

The restriction of the function $V(x, t)$ into Ω_r is denoted by $V_r(x, t)$, i.e. $V_r(x, t) = V(x, t)$, $(x, t) \in \Omega_r$, $r = \overline{1, N}$.

Let us set additional parameters

$$\lambda_r(x) = V_r(x, (r - 1)h), \quad r = \overline{1, N}$$

and

$$\lambda_{N+1}(x) = \lim_{t \rightarrow T - 0} V_N(x, t), \quad x \in [0, \omega],$$

and introduce the functions

$$\tilde{V}_r(x, t) = V_r(x, t) - \lambda_r(x) \quad \text{on } \Omega_r, \quad r = \overline{1, N}.$$

We then obtain the family of multipoint nonlinear boundary value problems for integro-differential equations with parameters

$$\frac{\partial \tilde{V}_r}{\partial t} = f\left(x, t, \psi(t) + \int_0^x \lambda_r(\xi) d\xi + \int_0^x \tilde{V}_r(\xi, t) d\xi, \lambda_r(x) + \tilde{V}_r\right), \quad (x, t) \in \overline{\Omega}_r, \quad r = \overline{1, N}, \quad (2.1)$$

$$\tilde{V}_r(x, (r - 1)h) = 0, \quad x \in [0, \omega], \quad r = \overline{1, N}, \quad (2.2)$$

$$g(x, \lambda_1(x), \lambda_{N+1}(x)) = 0, \quad x \in [0, \omega], \quad (2.3)$$

$$\lambda_r(x) + \lim_{t \rightarrow rh - 0} \tilde{V}_r(x, t) - \lambda_{r+1}(x) = 0, \quad x \in [0, \omega], \quad r = \overline{1, N}. \quad (2.4)$$

It can be easily shown that the families of problems (1.1), (1.2) and (2.1)–(2.4) are equivalent.

Suppose that for all $r = \overline{1, N + 1}$ and $x \in [0, \omega]$ the family of parameters $\lambda_r(x)$ is known. Then the functions $\tilde{V}_r(x, t)$, $(x, t) \in \Omega_r$ ($r = \overline{1, N}$), can be determined from the Cauchy problem (2.1), (2.2). For a fixed $x \in [0, \omega]$, this problem is equivalent to the family of mixed type systems of integral equations

$$\tilde{V}_r(x, t) = \int_{(r-1)h}^t f\left(x, \tau, \psi(t) + \int_0^x \lambda_r(\xi) d\xi + \int_0^x \tilde{V}_r(\xi, \tau) d\xi, \lambda_r(x) + \tilde{V}_r(x, \tau)\right) d\tau, \quad (2.5)$$

$$t \in [(r - 1)h, rh], \quad r = \overline{1, N}.$$

By substituting the values of $\lim_{t \rightarrow rh-0} \tilde{V}_r(x, t)$, found from (2.5), into (2.3) and (2.4), we obtain

$$g(x, \lambda_1(x), \lambda_{N+1}(x)) = 0,$$

$$\lambda_r(x) + \int_{(r-1)h}^{rh} f\left(x, t, \psi(t) + \int_0^x (\lambda_r(\xi) + \tilde{V}_r(\xi, \tau)) d\xi, \lambda_r(x) + \tilde{V}_r(x, t)\right) dt - \lambda_{r+1}(x) = 0.$$

This is a system of nonlinear functional equations in parameters $\lambda_r(x)$, $x \in [0, \omega]$, $r = \overline{1, N + 1}$. We rewrite this system in the form

$$Q_{1,h}\left(x, \lambda(x), \int_0^x \lambda(\xi) d\xi, \tilde{V}\right) = 0, \quad \lambda(x) \in \mathbb{R}^{n(N+1)}, \quad x \in [0, \omega]. \quad (2.6)$$

Condition 2.1. There exists $h > 0 : Nh = T$ ($N \in \mathbb{N}$), such that the family of systems of implicit nonlinear Fredholm integral equations (2.6), where $\tilde{V} = 0$, has a solution $\lambda^{(0)}(x) = (\lambda_1^{(0)}(x), \lambda_2^{(0)}(x), \dots, \lambda_{N+1}^{(0)}(x)) \in C([0, \omega], \mathbb{R}^{n(N+1)})$.

Let Condition 2.1 be met. We denote the solution of the Cauchy problem (2.1), (2.2), corresponding to $\lambda_r(x) = \lambda_r^{(0)}(x)$, by $\tilde{V}_r^{(0)}(x, t)$. Let us define the function

$$V^{(0)}(x, t) = \begin{cases} \lambda_r^{(0)}(x) + \tilde{V}_r^{(0)}(x, t) & \text{for } (x, t) \in \Omega_r, \quad r = \overline{1, N}, \\ \lambda_{N+1}^{(0)}(x) & \text{for } (x, t) \in [0, \omega] \cup \{T\}. \end{cases}$$

We choose some numbers $\rho_\lambda > 0$, $\rho_{\tilde{v}} > 0$, $\rho_V > 0$ and define the following sets:

$$S(\lambda^{(0)}(x), \rho_\lambda) = \left\{ \lambda(x) \in C([0, \omega], \mathbb{R}^{n(N+1)}) : \|\lambda - \lambda^{(0)}\|_1 < \rho_\lambda \right\},$$

$$S(\tilde{V}^{(0)}(x, [t]), \rho_{\tilde{v}}) = \left\{ \tilde{V}(x, [t]) \in C(\overline{\Omega}, \Omega_r, \mathbb{R}^{nN}) : \|(\tilde{V} - \tilde{V}^{(0)})[\cdot]\|_2 < \rho_{\tilde{v}} \right\},$$

$$S(V^{(0)}(x, t), \rho_V) = \left\{ V(x, t) \in C(\overline{\Omega}, \mathbb{R}^n) : \max_{(x,t) \in \overline{\Omega}} \|V(x, t) - V^{(0)}(x, t)\| < \rho_V \right\},$$

$$G_f(x, \rho_v) = \left\{ (x, t, u, v) \in \overline{\Omega} \times \mathbb{R}^{2n} : (x, t) \in \overline{\Omega}, \quad \|u - u^{(0)}(x, t)\| < \omega \cdot \rho_v, \quad \|v - v^{(0)}(x, t)\| < \rho_v \right\},$$

$$G_g(x, \rho_\lambda) = \left\{ (x, w_1, w_2) \in [0, \omega] \times \mathbb{R}^{2n} : \|w_1 - V^{(0)}(x, 0)\| < \rho_\lambda, \quad \|w_2 - V^{(0)}(x, T)\| < \rho_\lambda \right\}.$$

Condition 2.2. The function $f(x, t, u, v)$ has uniformly continuous partial derivatives f'_u, f'_v in $G_f(x, \rho_u, \rho_v)$ and the following inequalities hold:

$$\|f'_u(x, t, u, v)\| \leq L_1, \quad \|f'_v(x, t, u, v)\| \leq L_2 \quad \forall (x, t, u, v) \in G_f(x, \rho_u, \rho_v).$$

The function $g(x, w_1, w_2)$ has uniformly continuous partial derivatives g'_{w_1}, g'_{w_2} in $G_g(x, \rho_\lambda)$ and the following inequalities hold:

$$\|g'_{w_1}(x, w_1, w_2)\| \leq L_3, \quad \|g'_{w_2}(x, w_1, w_2)\| \leq L_4, \quad (x, w_1, w_2) \in G_g(x, \rho_\lambda).$$

Here L_i ($i = \overline{1, 4}$) are some constants.

Let Condition 2.2 be met. We take the pair $(\lambda^{(0)}(x), \tilde{V}^{(0)}(x, [t]))$ and determine the sequence $(\lambda^{(k)}(x), \tilde{V}^{(k)}(x, [t]))$, $k = 1, 2, \dots$, by the following algorithm.

Step 1.

- (i) Find $\lambda^{(1)}(x) = (\lambda_1^{(1)}(x), \lambda_2^{(1)}(x), \dots, \lambda_{N+1}^{(1)}(x)) \in C([0, \omega], \mathbb{R}^{n(N+1)})$ by solving the family of systems of implicit nonlinear Fredholm integral equations (2.6), where $\tilde{V} = \tilde{V}^{(0)}$.
- (ii) By solving the family of Cauchy problems (2.1), (2.2), where $\lambda(x) = \lambda^1(x)$, find the system of functions $\tilde{V}^{(1)}(x, [t])$.
- (iii) Define the function

$$V^{(1)}(x, t) = \begin{cases} \lambda_r^{(1)}(x) + \tilde{V}_r^{(1)}(x, t) & \text{for } (x, t) \in \Omega_r, \quad r = \overline{1, N}, \\ \lambda_{N+1}^{(1)}(x) & \text{for } (x, t) \in [0, \omega] \cup \{T\}. \end{cases}$$

Step k.

- (i) Find $\lambda^{(k)}(x) = (\lambda_1^{(k)}(x), \lambda_2^{(k)}(x), \dots, \lambda_{N+1}^{(k)}(x)) \in C([0, \omega], \mathbb{R}^{n(N+1)})$ by solving the family of systems of implicit nonlinear Fredholm integral equations (2.6), where $\tilde{V} = \tilde{V}^{(k-1)}$.
- (ii) By solving the family of Cauchy problems (2.1), (2.2), where $\lambda(x) = \lambda^2(x)$, find the system of functions $\tilde{V}^{(2)}(x, [t])$.
- (iii) Define the function

$$V^{(k)}(x, t) = \begin{cases} \lambda_r^{(k)}(x) + \tilde{V}_r^{(k)}(x, t) & \text{for } (x, t) \in \Omega_r, \quad r = \overline{1, N}, \\ \lambda_{N+1}^{(k)}(x) & \text{for } (x, t) \in [0, \omega] \cup \{T\}. \end{cases}$$

The following statement represents sufficient conditions for the existence of an isolated solution of the family of boundary value problems with parameters (2.1)–(2.4).

Theorem 2.1. *Let for some $h > 0 : Nh = T$ ($N = 1, 2, \dots$), $\rho_\lambda > 0$, $\rho_{\tilde{V}} > 0$, $\rho_V > 0$ fulfill Condition 2.1 and Condition 2.2 are met, the Jacobi matrix $\frac{\partial Q_{1,h}(x, \tilde{w}_1, \tilde{w}_2, \tilde{V})}{\partial \tilde{w}_1}$ has an inverse for $x \in [0, \omega]$ ($\tilde{w}_1 = \lambda(x)$, $\tilde{w}_2 = \int_0^x \lambda(\xi) d\xi$) and for all $(\lambda(x), \tilde{V}(x, [t])) \in S(\lambda^{(0)}(x), \rho_\lambda) \times S(\tilde{V}^{(0)}(x, [t]), \rho_{\tilde{V}})$, and let the following inequalities hold:*

- (1) $\left\| \left(\frac{\partial}{\partial \tilde{w}_1} Q_{1,h} \left(x, \lambda(x), \int_0^x \lambda(\xi) d\xi, \tilde{V} \right) \right)^{-1} \right\| \leq \gamma_1(h), \quad x \in [0, \omega], \quad \gamma_1(h) - \text{const};$
- (2) $q_1(h) = \gamma_1(h) e^{h \cdot \gamma_1(h) L_1 \omega} \frac{(L_1 \omega + L_2)^2 h^2}{1 - (L_1 \omega + L_2) h} < 1;$

$$(3) \quad \frac{\gamma_1(h)}{1 - q_1(h)} e^{h \cdot \gamma_1(h)L_1\omega} \max_{x \in [0, \omega]} \left\| Q_{1,h} \left(x, \lambda^{(0)}(x), \int_0^x \lambda^{(0)}(\xi) d\xi, \tilde{V}^{(0)} \right) \right\| < \rho_\lambda;$$

$$(4) \quad \frac{\gamma_1(h)}{1 - q_1(h)} e^{h \cdot \gamma_1(h)L_1\omega} \cdot \frac{(L_1\omega + L_2)h}{1 - (L_1\omega + L_2)h} \max_{x \in [0, \omega]} \left\| Q_{1,h} \left(x, \lambda^{(0)}(x), \int_0^x \lambda^{(0)}(\xi) d\xi, \tilde{V}^{(0)} \right) \right\| < \rho_{\tilde{V}};$$

$$(5) \quad \rho_\lambda + \rho_{\tilde{V}} < \rho_V.$$

Then for any $x \in [0, \omega]$ the sequence of pairs

$$(\lambda^{(k)}(x), \tilde{V}^{(k)}(x, [t])) \in S(\lambda^{(0)}(x), \rho_\lambda) \times S(\tilde{V}^{(0)}(x, [t]), \rho_{\tilde{V}})$$

converges to $(\lambda^*(x), \tilde{V}^*(x, [t]))$; an isolated solution of problem (2.1)–(2.4) in $S(\lambda^{(0)}(x), \rho_\lambda) \times S(\tilde{V}^{(0)}(x, [t]), \rho_{\tilde{V}})$. Moreover, the following estimates hold:

$$\begin{aligned} \|\lambda^* - \lambda^{(0)}\|_1 &\leq \frac{h \cdot \gamma_1(h)}{1 - q_1(h)} e^{h \cdot \gamma_1(h)L_1\omega} \frac{(L_1\omega + L_2)h}{1 - h(L_1\omega + L_2)} \max_{r=1, N} \tilde{K}_r, \\ \|\tilde{V}^* - \tilde{V}^{(0)}\|_2 &\leq \frac{(L_1\omega + L_2)h}{1 - (L_1\omega + L_2)h} \|\lambda^* - \lambda^{(0)}\|_1, \end{aligned}$$

where

$$\tilde{K}_r = \sup_{(x,t) \in \Omega_r} \left\| f \left(x, t, \int_0^x \lambda_r^{(0)}(\xi) d\xi, \lambda_r^{(0)}(x) \right) \right\|, \quad r = \overline{1, N}.$$

The proof of Theorem 2.1 is based on the sequential implementation of the steps of the proposed algorithm. To find the solution of the nonlinear operator equation with respect to the family of parameters for each fixed $x \in [0, \omega]$, a sharper version of the local Hadamard theorem [12, p. 41] is used.

Remark. The conditions of Theorem 2.1 are sufficient for the feasibility and convergence of the proposed algorithm.

Due to the equivalence of problems (2.1)–(2.4) and problems (1.1), (1.2), the following statement is true.

Theorem 2.2. *Let for some $h > 0$: $Nh = T$ ($N = 1, 2, \dots$), $\rho_\lambda > 0$, $\rho_{\tilde{V}} > 0$ and $\rho_V > 0$ all conditions of Theorem 2.1 are met. Then for any $x \in [0, \omega]$ the sequence of functions $V^{(k)}(x, t) \in S(V^{(0)}(x, t), \rho_V)$ converges to $V^*(x, t)$, an isolated solution of problem (1.1), (1.2) in $S(V^{(0)}(x, t), \rho_V)$ and the following estimate holds:*

$$\max_{(x,t) \in \Omega} \|V^*(x, t) - V^{(0)}(x, t)\| \leq \frac{h \cdot \gamma_1(h)}{1 - q_1(h)} e^{h \cdot \gamma_1(h)L_1\omega} \frac{h \cdot (L_1\omega + L_2)}{(1 - h(L_1\omega + L_2))^2} \cdot K,$$

where

$$K = \max_{r=1, N} \sup_{(x,t) \in \Omega_r} \left\| f \left(x, t, \int_0^x (V^{(0)}(\xi, t) - V^{(0)}(\xi, (r-1)h)) d\xi, V^{(0)}(x, t) - V^{(0)}(x, (r-1)h) \right) \right\|.$$

Theorem 2.2 is a corollary of Theorem 2.1.

Acknowledgements

This research has been funded by the Committee of Science of the Ministry of Science and Higher Education of the Republic of Kazakhstan (Grant # AP23488811).

References

- [1] P. B. Abdimanapova and S. M. Temesheva, On a solution of a nonlinear nonlocal boundary value problem for one class of hyperbolic equation. *Lobachevskii J. Math.* **44** (2023), no. 7, 2529–2541.
- [2] P. B. Abdimanapova and S. M. Temesheva, Well-posedness criteria for one family of boundary value problems. *Bull. Karaganda Univ. Math. Ser.* no. 4(112), (2023), 5–20.
- [3] A. T. Assanova, E. A. Bakirova, Zh. M. Kadirbayeva and R. E. Uteshova, A computational method for solving a problem with parameter for linear systems of integro-differential equations. *Comput. Appl. Math.* **39** (2020), no. 3, Paper no. 248, 23 pp.
- [4] A. T. Assanova, S. G. Karakenova, S. T. Mynbayeva and R. E. Uteshova, A solution to a nonlinear Fredholm integro-differential equation. *Quaest. Math.* **47** (2024), no. 1, 61–73.
- [5] A. T. Assanova, A. P. Sabalakhova and Z. M. Toleukhanova, On the unique solvability of a family of boundary value problems for integro-differential equations of mixed type. *Lobachevskii J. Math.* **42** (2021), no. 6, 1228–1238.
- [6] A. T. Assanova, S. S. Zhumatov, S. T. Mynbayeva and S. G. Karakenova, On solvability of boundary value problem for a nonlinear Fredholm integro-differential equation. *Bull. Karaganda Univ., Math. Ser.* no. 1(105) (2022), 25–34.
- [7] È. A. Bakirova, N. B. Iskakova and A. T. Assanova, A numerical method for solving a linear boundary value problem for integrodifferential equations based on spline approximation. (Russian) *Ukrain. Mat. Zh.* **71** (2019), no. 9, 1176–1191; translation in *Ukrainian Math. J.* **71** (2020), no. 9, 1341–1358
- [8] O. A. Boichuk and Ī. A. Golovatska, Boundary value problems for systems of integro-differential equations. (Ukrainian) *Nelīnīvī Koliv.* **16** (2013), no. 4, 460–474; translation in *J. Math. Sci. (N.Y.)* **203** (2014), no. 3, 306–321.
- [9] A. A. Boichuk and A. M. Samoilenko, *Generalized Inverse Operators and Fredholm Boundary-Value Problems*. VSP, Utrecht, 2004.
- [10] D. S. Dzhumabayev, Criteria for the unique solvability of a linear boundary-value problem for an ordinary differential equation. *USSR Computational Mathematics and Mathematical Physics* **29** (1989), no. 1, 34–46.
- [11] D. S. Dzhumabaev and A. T. Assanova, Indications of correct solvability of linear nonlocal boundary value problems for systems of hyperbolic equations. *Doklady NAN Ukraine* **4** (2010), 7–11.
- [12] D. S. Dzhumabaev and S. M. Temesheva, A parametrization method for solving nonlinear two-point boundary value problems. (Russian) *Zh. Vychisl. Mat. Mat. Fiz.* **47** (2007), no. 1, 39–63; translation in *Comput. Math. Math. Phys.* **47** (2007), no. 1, 37–61
- [13] D. S. Dzhumabaev and S. M. Temesheva, Necessary and sufficient conditions for the existence of an “isolated” solution of a nonlinear two-point boundary value problem. (Russian) *Nelīnīvī Koliv.* **15** (2012), no. 4, 435–446; translation in *J. Math. Sci. (N.Y.)* **194** (2013), no. 4, 341–353.

-
- [14] D. S. Dzhumabaev and S. M. Temesheva, Bounded solution on a strip to a system of nonlinear hyperbolic equations with mixed derivatives. *Bull. Karaganda Univ., Math. Ser.* no. 4(84) (2016), 35–45.
- [15] D. S. Dzhumabaev and S. M. Temesheva, Existence criteria for an isolated solution of a nonlinear boundary value problem. (Russian) *Ukrain. Mat. Zh.* **70** (2018), no. 3, 356–365; translation in *Ukrainian Math. J.* **70** (2018), no. 3, 410–421.
- [16] N. Iskakova, S. Temesheva and R. Utesheva, On a problem for a delay differential equation. *Math. Methods Appl. Sci.* **46** (2023), no. 9, 11283–11297.
- [17] Zh. Kadirbayeva, E. Bakirova and A. Tleulessova, Solving Fredholm integro-differential equations involving integral condition: a new numerical method. *Math. Slovaca* **74** (2024), no. 2, 403–416.

N. Partsvania

- Initial Value Problem on an Infinite Interval for First Order
Advanced Differential Equations 221

G. Petryna, A. Murzabaeva, Z. Khaletska

- Approximation of Stochastic Systems with Integral-Type Delay
by Stochastic Systems without Delay 227

V. A. Pokhachevskiy, V. V. Bykov

- On the Baire Classification of the Weak Oscillation Exponents
of Roots and Hypermultiple Roots 233

O. Pravdyvyi, A. Stanzhytskyi, O. Martynyuk

- Limiting Behavior of Invariant Measures of Stochastic Functional-Differential
Neutral Equations in Hilbert Space 237

A. Rontó, M. Rontó, N. Rontóová, I. Varga

- On Investigation of Non-Linear Differential Systems with Mixed
Boundary Conditions 244

J. Šremr

- On Lyapunov and Krasovskii Stability/Instability of Pendulum Equation
with Non-Constant Coefficients 249

S. Staněk

- Dirichlet Problem for Singular Fractional Differential Equations
with Given Maximal Value for Positive Solutions 257

O. Stanzhytskyi, D. Shtefan, R. Uteshova

- Asymptotic Equivalence of Infinite-Dimensional Linear Stochastic Systems 262

T. Tadumadze, T. Shavadze

- On the Optimization Problem for the Quasi-Linear Neutral
Functional-Differential Equation with the Discontinuous Initial Condition 268

R. Taranets, M. Chugunova

- On a Control-Volume Model for Fiber Coating 272

S. Temesheva, P. Abdimanapova

- On the Solvability of a Family of Nonlinear Two-Point Boundary Value Problems
for Systems of Integro-Differential Equations and its Application 275