



Organized by



# 15th Tetrahedron Symposium

CHALLENGES IN BIOORGANIC AND ORGANIC MEDICINAL CHEMISTRY

24–27 June 2014 | London, UK

## Programme Book

[www.tetrahedron-symposium.elsevier.com](http://www.tetrahedron-symposium.elsevier.com)



**15<sup>th</sup> Tetrahedron Symposium**  
CHALLENGES IN BIOORGANIC & ORGANIC MEDICINAL CHEMISTRY

24-27 JUNE 2014  
LONDON, UK

## **15<sup>th</sup> Tetrahedron Symposium**

### ***Challenges in Bioorganic and Organic Medicinal Chemistry***

**24<sup>th</sup> – 27<sup>th</sup> June 2014**

***London, United Kingdom***

Organised by:



**ELSEVIER**

**Delegates have been emailed the URL to access the abstracts.  
If you have any questions please ask at the registration desk.**

[P1.94]

**Tablets on the basis of herbal substance "Limonidin"**

G.E. Zhussupova<sup>1</sup>, K.B. Murzagulova<sup>2</sup>, A.I. Zhussupova<sup>3</sup>, A.V. Gadetskaya<sup>1</sup>, Z.A. Abilov<sup>1</sup>  
<sup>1</sup>*Al-Farabi Kazakh National University, Kazakhstan, <sup>2</sup>Romat Pharmaceutical Company, Kazakhstan*

Isolation of substances from the roots and aerial parts of local medicinal plant *Limonium gmelinii* was conducted using a simple, economically feasible and environmentally friendly technological scheme with a high yield (up to 35 % of the dried raw material weight). Vegetable substances extracted from the test plants in the form of dry extracts are characterized by hygroscopicity; their complex with  $\beta$ -cyclodextrin was obtained in order to reduce this hygroscopicity. The molecular complexation of  $\beta$ -cyclodextrin with the substance was studied using 2 methods: a) method of paste-forming; b) method of briquetting. The process of nano-encapsulation was monitored by observation of changes in shape and size of the particles. The resulting complex of the "Limonidin" substance with cyclodextrin is a light-brown powder with the patches of darker particles with faint odor, moisture content of not more than 3 %, soluble in water, 0.1 N HCl solution and aqueous solutions of ethyl alcohol (30, 50 %). Bulk density before shrinkage is 0.707 g/cm, after the shrinkage - 0.809 g/cm. The study of the complex was carried out using the methods of IR and UV spectroscopy and diffractometry. The biopharmaceutical properties of the complex were studied: release of the active substance in the amount of not less than 46.0 %, in various environments. Quality specification was designed. On the basis of this complex of the substance and  $\beta$ -cyclodextrin, two sets of granulates were developed. The tablets produced on their basis meet the requirements of pharmacopoeia on the following indicators of quality: compressibility profiles, hardness, friability, friability, disintegration, quantification of the active substance and its release.

Keywords: tablets, herbal, substance, production

[P1.95]

**Novel compound from *Limonium myrianthum* possessing antioxidant activity**

A.V. Gadetskaya<sup>1</sup>, G.E. Zhussupova<sup>1</sup>, M.K. Murzakmetova<sup>2</sup>, S.A. Ross<sup>3</sup>, A.I. Zhussupova<sup>\*1</sup>  
<sup>1</sup>*Al-Farabi Kazakh National University, Kazakhstan*, <sup>2</sup>*Institute for Human and Animal Physiology  
MES RK, Kazakhstan*, <sup>3</sup>*University of Mississippi, USA*

Plants of the *Limonium* Mill genus (*Plumbagenaceae* family) are represented by 300 species world-wide and 18 in Kazakhstan. Two species, *L. gmelinii* and *L. myrianthum*, are present in stocks exceeding 50 thousand tons in Kazakhstan. The current studies are focused on *L. myrianthum*. Specimens of *L. myrianthum* were collected in South Kazakhstan in August 2010. Finely ground, air-dried roots (300 g) were subjected to sequential extraction with *n*-hexane (0.3 L x 2; 48 h), acetone (0.3 L x 3; 48 h) and MeOH (0.3 L x 3; 48 h) at 30 °C with constant mixing. Extracts were evaporated to dryness under reduced pressure at 35-37 °C. Acetone extracts of the roots showed the highest antioxidant activities and were combined then subjected to silica gel chromatography with elution successively with 100% CH<sub>2</sub>Cl<sub>2</sub>, then CH<sub>2</sub>Cl<sub>2</sub>: MeOH mixtures (5%, 10%, 15%, 25%, 30%, 35%) and finally 100% MeOH to yield 8 fractions. Active fractions were combined and subjected to Sephadex LH-20 chromatography to yield 33 fractions of decreasing molecular mass. A yellow compound isolated from fractions 12 and 13 was identified as epigallocatechin-2-O-*p*-phenoxy (C<sub>21</sub>H<sub>18</sub>O<sub>8</sub>; mol wt 398.36). It showed potent antioxidant activity in assays of liver microsomal lipid peroxidation. At 4 µg/mL, it reduced the LPO level to 62 % of control and at 20 µg/mL, it reduced the level of LPO to 12 % of control. The antioxidant properties of epigallocatechin-2-O-*p*-phenoxy require further investigation. Two-dimensional spectra were obtained on a Bruker DRX-500 spectrometer. GC-MS spectra were obtained using an HP 6890 gas chromatograph.

Keywords: novel compound, antioxidant activity