








21st International Symposium on  
**HEAVY ION FUSION**  
Astana, Kazakhstan • July 18-22, 2016

# Book of Abstracts



 NAZARBAYEV  
UNIVERSITY  
NATIONAL  
LABORATORY ASTANA

I S T C  
  
М Н Т Ц

**БИОХИМ**  
П Р И Б О Р

**ORAL TALKS**

STOPPING OF RELATIVISTIC IONS IN MULTICOMPONENT PLASMAS Yu.V. Arkhipov, A.B. Ashikbayeva, A. Askaruly, A.E. Davletov, I.M. Tkachenko	37
SUBSYSTEM TABULATION OF THE SINGLE PASS RF DRIVER R. Burke, A. Burke, A. K. Delavar	38
INVESTIGATIONS ON THE INTERACTION BETWEEN THE LOW ENERGY HEAVY ION BEAMS AND HYDROGEN PLASMA R.Cheng, X.Zhou, Y.Wang, Y.Lei, Y.Sun, J.Ren, X.Ma, G.Xiao, Y.Zhao, R.Gavrillin, S. Savin, I.Roudskoy, A.Golubev, D.H.H.Hoffmann	39
CREATION OF NANOPOROUS $\text{SiO}_2/\text{Si}$ WITH PRECIPITATED ZN USING TRACK TECHNOLOGY A. Dauletbekova, A. Alzhanova, A. Akilbekov, F. Komarov, L. Vlasukova, M. Ibraimov, M. Zdorovets	40
TWO-BODY AND THREE-BODY MESONIC BOUND STATES IN ULTRADENSE PLASMAS OF FUSION CATALYSIS CONCERN C. Deutsch	41
COLLISIONAL AND OPTICAL PROPERTIES OF THE DENSE SEMICLASSICAL PLASMA K.N. Dzhumagulova, E.O. Shalenov, T.S. Ramazanov, G. Röpke, H. Reinholz	42
THERMODYNAMIC PROPERTIES OF NON-ISOTHERMAL PLASMA OF ICF M.T.Gabdullin, T.S.Ramazanov, T.N.Ismagambetova, Zh.A.Moldabekov, Zh.Y.Otarbay, M.B.Mamatova	43
ELECTRON-ION ENERGY PARTITION FOR ALPHA PARTICLE MOVING IN FUSION DT PLASMAS MIXED WITH HOT AU B. He, J. Wang	44
THE ECONOMICS OF AN HIF SPRFD FUSION SYSTEM IN A DESERT REGION Ch. Helsley	45
DEVELOPMENT OF AN ADVANCED WATER FILTERING SYSTEM BASED ON GRAPHENE IRRADIATED BY GAS CLUSTER AND HIGHLY CHARGED IONS Z. Insepov, A. Ainabayev, K. Dybyspayeva, Z. Ramazanova, G. Ikram, S. Kirkpatrick, M. Walsh, M. Terasawa	46
EFFECT OF TAPERED MAGNETIC FIELD ON EXPANDING LASER PRODUCED PLASMA FOR HEAVY ION INERTIAL FUSION T. Kanesue, Sh. Ikeda, M. Okamura	47
TARGET IMPLOSION UNIFORMITY IN HEAVY ION FUSION T. Karino, S. Kawata, S. Kondo, T. Iinuma, T. Kubo, H. Kato, A. I. Ogoyski	48
EFFECTS OF IRRADIATION OF STAINLESS STEEL AND TIMON COATINGS ON STAINLESS WITH LOW-ENERGY ALPHA-PARTICLES, KRYPTON AND XENON IONS Kislitsin S., Uglov V., Larionov A.	49
DEVELOPMENT AND FABRICATION OF ELECTRO-MAGNETIC SHOCK TUBE WITH HIGH RESOLUTION SPECTROMETER FOR STOPPING POWER MEASUREMENT IN ATOMIC HYDROGEN GAS K. Kondo, K. Kawachi, H. Kurita, T. Takahashi, Y. Oguri	50

## Collisional and optical properties of the dense semiclassical plasma

K.N. Dzhumagulova<sup>\*1</sup>, E.O. Shalenov<sup>\*1</sup>, T.S. Ramazanov<sup>1</sup>, G. Röpke<sup>2</sup>, H. Reinholz<sup>2</sup>

<sup>1</sup>IETP, Department of Physics, al-Farabi Kazakh National University, al-Farbi 71, 050040Almaty, Kazakhstan

<sup>2</sup>Institute of Physics, University of Rostock, A.-Einstein-Str. 23-24, 18059Rostock, Germany

\*shalenov.erik@mail.ru

Elastic and non elastic scattering of the electrons are the elementary processes that continuously attract the attention of the researchers. For developing of the applications, where plasma is a working medium, one needs adequate data on physical properties of the system. Electron scattering strongly influences the transport, optical and other properties of plasmas. A large number of various theoretical and experimental studies is devoted to the scattering of the electrons on the noble gas atoms. Various theoretical methods are used for studying of the elementary processes and plasma properties.

In this work we consider the dense partially ionized plasma of the noble gases consisting of the electrons, ions and atoms. Particle densities are in the range of  $n = 10^{20} + 10^{23} \text{ cm}^{-3}$  and the temperature range is from  $2.5 \times 10^4 \text{ K}$  up to  $5 \times 10^4 \text{ K}$ .

In work [1] the effective potential of electron-atom interaction, taking into account both quantum-mechanical effect of diffraction and screening effects, is presented. The way to take into account the dynamic screening was proposed in work [2], where the statical Debye radius was replaced by the dynamic one:

$$r_o = r_D \left(1 + \frac{v^2}{v_{Th}^2}\right)^{1/2}, \quad (1)$$

here  $v$  is the relative velocity of the colliding particles,  $v_{Th}$  is the thermal velocity of the particles in the system. Then the effective potential of electron-atom interaction with dynamic screening can be rewritten as [3]:

$$\Phi_{ea}(r) = -\frac{e^2 \alpha}{2r^4 (1 - 4\lambda_{ea}^2 / r_o^2)} \left( e^{-Br} (1 + Br) - e^{-Ar} (1 + Ar) \right)^2, \quad (2)$$

$$\text{where } A^2 = \frac{1}{2\lambda_{ea}^2} \left( 1 + \sqrt{1 - 4\lambda_{ea}^2 / r_o^2} \right),$$

$$B^2 = \frac{1}{2\lambda_{ea}^2} \left( 1 - \sqrt{1 - 4\lambda_{ea}^2 / r_o^2} \right).$$

In the framework of this pseudopotential model for the particle interactions, the scattering phase

shifts  $\delta_l$  were calculated on the basis of the Ca-logero equation:

$$\frac{d\delta_l(k, r)}{dr} = -\frac{1}{k} U(r) \times \quad (3)$$

$$\left[ \cos \delta_l(k, r) \cdot j_l(kr) - \sin \delta_l(k, r) \cdot n_l(kr) \right]^2$$

$$\delta_l(k, 0) = 0.$$

Here  $k$  is the magnitude of the wave vector of the incident particle,  $U(r) = \frac{2m}{\hbar^2} \Phi_{ea}(r)$ ,  $j_l(kr)$  and

$n_l(kr)$  are the Ricatti - Bessel functions.

Phase shifts enable us to calculate the transport scattering cross section  $Q_{ea}^T(k)$ . The collision frequency of the electrons with atoms  $\nu_{ea}$  can be obtained by the following expression:

$$\nu_{ea} = 4 \sqrt{\frac{2}{\pi}} n_a \sqrt{\frac{k_B T}{\mu_{ea}}} \int_0^\infty Q_{ea}^T(g) g^3 \text{Exp}(-g^2) dg, \quad (4)$$

here  $\mu_{ea} = m_e m_a / (m_e + m_a)$  is the reduced mass of the electron-atom pair,  $g$  is the reduced velocity.

Dielectric function due to the electron - atom interactions was calculated using the generalized Drude-Lorentz equation:

$$\varepsilon(\omega) = 1 - \frac{\omega_p^2}{\omega^2 - i\nu_{ea}\omega}. \quad (5)$$

From the dielectric function the reflectivity of the semiclassical electron plasma is calculated, using the Fresnel formula, and is compared with experimental results [4].

### References

- [1] T.S. Ramazanov *et al* 2005 *Phys.Plasmas* **12**, 092702
- [2] D. Kremp *et al* 2005 *Springer*, p. 326
- [3] K.N. Dzhumagulova *et al* 2014 *Contr. Plasma Phys* **55**, 230-235
- [4] Yu.B. Zaporoghets *et al* 2016 *Contr. Plasma Phys*. in press