# Third International Conference on Analysis and Applied Mathematics

# ICAAM 2016

# THE ABSTRACT BOOK

Institute of Mathematics and Mathematical Modeling

www.math.kz

Editor:
Allaberen Ashyralyev

07-10 September 2016

Institute of Mathematics
and Mathematical Modeling
Almaty, Kazakhstan

**УДК 517.9 (043.2)**
**ББК 22.161**
**Т 45**

Third International Conference on Analysis and Applied Mathematics (ICAAM 2016): THE ABSTRACT BOOK. (Тезисы докладов) / Ред. Allaberen Ashyralyev. – Алматы: Институт математики и математического моделирования, 2016. – 325 с.

**ISBN 978-601-280-766-0**
**Т 45**

The conference is organized biannually. Previous conferences were held in Gumushane, Turkey in 2012 and in Shymkent, Kazakhstan in 2014.

The aim of the International Conference on Analysis and Applied Mathematics (ICAAM) is to bring mathematicians working in the area of analysis and applied mathematics together to share new trends of applications of mathematics. In mathematics, the developments in the field of applied mathematics open new research areas in analysis and vice versa. That is why, we plan to found the conference series to provide a forum for researches and scientists to communicate their recent developments and to present their original results in various fields of analysis and applied mathematics.

Web page: http://icaam-online.org

УДК 517.9 (043.2)
ББК 22.161

ISBN 978-601-280-766-0
© Third International Conference on Analysis and Applied Mathematics ICAAM 2016

**Conference Sections and Minisymposiums :**

- Analysis
- Applied Mathematics
- Mathematics Education
- Other Topics (Algebra, Geometry, Topology...)
- Minisymposium:Mathematical Modelling
- Minisymposium:Spectral Theory of Differential Operators
- Minisymposium:Dynamical Systems

**Invited Speakers**

- Prof. **Alexander Ivanovich Aptekarev**, Director of the Keldysh Institute of Applied Mathematics of Russian Academy of Sciences, Moscow, Russia
- Prof. **Shavkat Arifdzhanovich Alimov**, Head of the laboratory of mathematical modeling and Chief Scientist of the Malaysian Institute of Microelectronic Systems (MIMOS) in Kuala Lumpur, Malaysia
- Prof. **Ravi Agarwal**, Texas A & M University, USA
- Prof. **Vasilii Nikolaevich Denisov**, Lomonosov Moscow State University, Faculty of Computational Mathematics and Cybernetics, Russia
- Prof. **Mokhtar Kirane**, Universitet de La Rochelle, La Rochelle, France
- Prof. **Ljubisa D.R. Kocinac**, University of Nis, Serbia
- Prof. **Vladimir Alekseevich Levin**, Head of the Department of Computational Mechanics, Mechanics and Mathematics Faculty, Lomonosov Moscow State University, Russia
- Prof. **Nedyu I. Popivanov**, The Faculty of Mathematics and Informatics, University of Sofia "St. Kliment Ohridski", Bulgaria
- Prof. **Mikhail Vladimirovich Ruzhansky**, Department of Mathematics, Imperial College London, UK
- Prof. **Alexander Leonidovich Skubachevskii**, Head of Department of "Differential Equations", Peoples Friendship University of Russia, Moscow, Russia
- Prof. **Andrei Andreevich Shkalikov**, Head of the Laboratory of operator models and spectral analysis, Lomonosov Moscow State University, Russia
- Prof. **Iskander A. Taimanov**, Chair of Geometry and Topology, Novosibirsk State University, Head of the Laboratory of Dynamical Systems, Sobolev Institute of Mathematics, Russia

## COMMITTEES

### HONORARY CHAIR

- Prof. **Tynysbek Kalmenov**, Institute of Mathematics and Mathematical Modelling, Kazakhstan

### CHAIRS

- Prof. **Allaberen Ashyralyev**, Emeritus Professor, Turkey and Turkmenistan
- Prof. **Mukhtarbay Otelbaev**, Institute of Mathematics and Mathematical Modelling, Kazakhstan

### CO-CHAIRS

- Prof. **Makhmud Sadybekov**, Institute of Mathematics and Mathematical Modelling, Kazakhstan
- Assoc. Prof. **Abdullah S. Erdogan**, Turkey

### INTERNATIONAL ADVISORY BOARD

- Prof. Mahmoud Abdel-Aty, Zewail City of Science and Technology, Egypt
- Prof. Ravi P. Agarwal, Texas A&M University, USA
- Prof. Shavkat Alimov, The Institute MIMOS, Malaysia
- Aslanbek Amrin, Ministry of Education and Science of the Republic of Kazakhstan, Kazakhstan
- Prof. Alexander Aptekarev, Keldysh Institute of Applied Mathematics, Russia
- Prof. Ravshan R. Ashurov, Institute of Mathematics of Academy of Science, Uzbekistan
- Prof. Sergey V. Astashkin, Samara State University, Russia
- Prof. Takir Balykbaev, Executive Office of the President of the Republic of Kazakhstan, Kazakhstan
- Prof. Feyzi Basar, Turkey
- Prof. Nazarbai Bliev, Institute of Mathematics and Mathematical Modelling, Kazakhstan
- Prof. Martin Bohner, Missouri University of Science and Technology, USA
- Prof. Valry C. Covachev, Institute of Mathematics and Informatics, Bulgarian Academy of Sciences, Bulgaria
- Prof. Vasilii Denisov, Lomonosov Moscow State University, Russia
- Prof. Askar Dzhumadil'daev, Institute of Mathematics and Mathematical Modelling, Kazakhstan
- Prof. Lyudvig Faddeev, St. Petersburg Department of Steklov Mathematical Institute, Russia
- Prof. Sergei Kharibegashvili, A. Razmadze Mathematical Institute, Georgia
- Prof. Stanislav Kharin, Institute of Mathematics and Mathematical Modelling, Kazakhstan
- Prof. Mokhtar Kirane, University of La Rochelle, France
- Prof. Michael Klibanov, The University of North Carolina at Charlotte, USA
- Prof. Ljubisa D.R. Kocinac, University of Nis, Serbia

- Prof. Ismail Kucuk, American University of Sharjah, UAE
- Prof. Vladimir Levin, Lomonosov Moscow State University, Russia
- Prof. Hongyu Liu, Hong Kong Baptist University, Hong Kong
- Prof. Ihor Lubashevsky, University of Aizu, Japan
- Prof. Alexey Lukashov, Saratov State University, Russia
- Prof. Eberhard Malkowsky, University of Giessen, Germany
- Prof. Mohammad Mursaleen, Aligarh Muslim University, India
- Prof. Adam Nakhushev, Institute of Applied Mathematics and Automation, Russia
- Prof. Zuhair Nashed, University of Central Florida, USA
- Prof. Sandra Pinelas, Academia Militar, Portugal
- Prof. Sergey Piskarev, Lomonosov Moscow State University, Russia
- Prof. Nedyu Popivanov, University of Sofia "St. Kliment Ohridski", Bulgaria
- Prof. Reimund Rautmann, University of Paderborn, Germany
- Prof. Mikhail Ruzhansky, Imperial College London, UK
- Prof. Reza Saadati, Islamic Azad University, Iran
- Prof. Valery Sergejevich Serov, University of Oulu, Finland
- Prof. Andrei Shkalikov, Lomonosov Moscow State University, Russia
- Prof. Yury Shestopalov, Karlstad University, Sweden
- Prof. Alexander Skubachevskii, Peoples Friendship University of Russia, Russia
- Prof. Pavel Sobolevskii, Hebrew University, Israel & Universidade Federal do Ceara, Brazil
- Prof. Hari M. Srivastava, University of Victoria, Canada
- Prof. Iskander Taimanov, Sobolev Institute of Mathematics, Russia
- Prof. Murat Zhurinov, National Academy of Sciences of the Republic of Kazakhstan, Kazakhstan


## INTERNATIONAL ORGANIZING COMMITTEE


- Prof. Ljudmila Alexeyeva, Institute of Mathematics and Mathematical Modelling, Kazakhstan
- Prof. Fikret Aliev, Baku State University, Azerbaijan
- Prof. Tolegen Amanbaev, M. Auezov South Kazakhstan State University, Kazakhstan
- Prof. Anar Asanova, Institute of Mathematics and Mathematical Modelling, Kazakhstan
- Prof. Bektur Baizhanov, Institute of Mathematics and Mathematical Modelling, Kazakhstan
- Prof. Daurenbek Bazarkhanov, Institute of Mathematics and Mathematical Modelling, Kazakhstan
- Prof. Maktagali Bektemesov, Al-Farabi Kazakh National University, Kazakhstan
- Prof. Amantay Berzhanov, Zhubanov State University, Kazakhstan
- Prof. Yessen Bidaibekov, Abai Kazakh National Pedagogical University, Kazakhstan
- Prof. Galina Bizhanova, Institute of Mathematics and Mathematical Modelling, Kazakhstan
- Prof. Nurzhan Bokaev, Gumilev Eurasian National University, Kazakhstan
- Prof. Huseyin Cakalli, Maltepe University, Turkey

- Prof. Claudio Cuevas, Federal University of Pernambuco, Brazil
- Prof. Gennadii Demidenko, Sobolev Institute of Mathematics, Russia
- Prof. Huseyin Demir, Ondokuz Mayis University, Turkey
- Prof. Muvasherhan Dzhenaliev, Institute of Mathematics and Mathematical Modelling, Kazakhstan
- Prof. Dulat Dzhumabaev, Institute of Mathematics and Mathematical Modelling, Kazakhstan
- Assoc. Prof. Okan Gercek, Turkey
- Assoc. Prof. Selim Guvercin, Suleyman Demirel University, Kazakhstan
- Prof. Baltabek Kanguzhin, Al-Farabi Kazakh National University, Kazakhstan
- Prof. Viktor Korzyuk, Institute of Mathematics of the NAS of Belarus, Belarus
- Prof. Bakhytbek Koshanov, Institute of Mathematics and Mathematical Modelling, Kazakhstan
- Prof. Aleksandr Kozhanov, Sobolev Institute of Mathematics, Russia
- Assoc. Prof. Leonid Kritskov, Lomonosov Moscow State University, Russia
- Prof. Misir Mardanov, Institute of Mathematics and Mechanics of the National Academy of Sciences of Azerbaijan, Azerbaijan
- Prof. Mukhammed Meredov, Turkmenistan
- Prof. Saltanbek Mukhambetzhanov, Al-Farabi Kazakh National University, Kazakhstan
- Prof. Oktay Mukhtarov, Gaziosmanpasa University, Turkey
- Prof. Musakhan Muratbekov, Taraz State Pedagogical Institute, Kazakhstan
- PhD. Daulet Nurakhmetov, S. Seifullin Kazakh AgroTechnical University, Kazakhstan
- Prof. Erlan Nursultanov, Kazakhstan Branch of Lomonosov Moscow State University, Kazakhstan
- Prof. Ryskul Oinaruly, Gumilev Eurasian National University, Kazakhstan
- Prof. Kordan Ospanov, Gumilev Eurasian National University, Kazakhstan
- Prof. Pavel Pankov, International University of Kyrgyzstan, Kyrgyzstan
- Prof. Arsen Pskhu, Institute of Applied Mathematics and Automation, Russia
- Prof. Nusrat Radjabov, Tajik State National University, Tajikistan
- Prof. Stasys Rutkauskas, Institute of Mathematics and Computer Science, Lithuania
- Prof. Makhmud S. Salakhitdinov, National University of Uzbekistan, Uzbekistan
- Prof. Abdizhahan Sarsenbi, M. Auezov South Kazakhstan State University, Kazakhstan
- Assoc. Prof. Merey Sarsengeldin, Institute of Mathematics and Mathematical Modelling, Kazakhstan
- Prof. Vladimir Savchin, Peoples' Friendship University of Russia, Russia

- Prof. Mirbolat Sikhov, Al-Farabi Kazakh National University, Kazakhstan
- Prof. Esmuhanbet Smailov, Institute of Applied Mathematics, Kazakhstan
- Prof. Alexandr Soldatov, Belgorod State University, Russia
- PhD. Durvudkhan Suragan, Institute of Mathematics and Mathematical Modelling, Kazakhstan
- Prof. Murat Ramazanov, Buketov Karaganda State University, Kazakhstan
- Prof. Pavel Terekhin, Saratov State University, Russia
- Prof. Marat Tleubergenov, Institute of Mathematics and Mathematical Modelling, Kazakhstan
- Prof. Nazerke Tleukhanova, Gumilev Eurasian National University, Kazakhstan
- Prof. Batirkhan Turmetov, A. Yasavi International Kazakh-Turkish University, Kazakhstan
- Prof. Umirbaev Ualbay, Wayne State University, USA
- Prof. Victor G. Zvyagin, Voronezh State University, Russia

**TECHNICAL PROGRAM COMMITTEE**

- Asst. Prof. Necmettin Aggez, Turkey
- Assoc. Prof. Deniz Agirseven, Trakya University, Turkey
- Meyram Akhymbek, Institute of Mathematics and Mathematical Modelling, Kazakhstan
- Gaukhar Arepova, Institute of Mathematics and Mathematical Modelling, Kazakhstan
- Assoc. Prof. Maksat Ashyraliyev, Baheehir University, Turkey
- Asst. Prof. Abil Aubakirov, Institute of Mathematics and Mathematical Modelling, Kazakhstan
- Samat Kassabek, Institute of Mathematics and Mathematical Modelling, Kazakhstan
- Aidyn Kassymov, Institute of Mathematics and Mathematical Modelling, Kazakhstan
- Assoc. Prof. Mehmet Emir Koksal, Ondokuz Mayis University, Turkey
- Asst. Prof. Yildirim Ozdemir, Duzce University, Turkey
- Lecturer Ali Ugur Sazaklioglu, University of Turkish Aeronautical Association, Turkey
- Asst. Prof. Fatma S. Ozesenli Tetikoglu, Turkey
- Berik Torebek, Institute of Mathematics and Mathematical Modelling, Kazakhstan
- Nurgissa Yessirkegenov, Institute of Mathematics and Mathematical Modelling, Kazakhstan
- Asst. Prof. Ozgur Yildirim, Yildiz Technical University, Turkey

## FOREWORD

The Organizing Committee of ICAAM and Institute of Mathematics and Mathematical Modelling are pleased to invite you to the Third International Conference on Analysis and Applied Mathematics, ICAAM 2016. The meeting will be held on September 7-10, 2016 in Almaty, Kazakhstan. This conference is dedicated to 70th birthday of Prof. Tynysbek Kalmenov.

The conference is organized biannually. Previous conferences were held in Gumushane, Turkey in 2012 and in Shymkent, Kazakhstan in 2014. The proceedings of ICAAM 2012 and ICAAM 2014 were published in AIP (American Institute of Physics) Conference Proceedings. Institute of Mathematics and Mathematical Modelling is pleased to host the third conference which is focused on various topics of analysis and its applications, applied mathematics and modeling.

The conference will consist of plenary lectures, mini symposiums and contributed oral presentations. The proceedings of ICAAM 2016 will be published in AIP (American Institute of Physics) Conference Proceedings. Selected full papers of this conference will be published in peer-reviewed international journals:

- FILOMAT (Science Citation Index),
- BOUNDARY VALUE PROBLEMS (Science Citation Index),
- CONTEMPORARY ANALYSIS AND APPLIED MATHEMATICS.

The aim of the International Conference on Analysis and Applied Mathematics (ICAAM) is to bring mathematicians working in the area of analysis and applied mathematics together to share new trends of applications of mathematics. In mathematics, the developments in the field of applied mathematics open new research areas in analysis and vice versa. That is why, we plan to found the conference series to provide a forum for researches and scientists to communicate their recent developments and to present their original results in various fields of analysis and applied mathematics. The Conference Organizing Committee would like to thank our sponsors. The main organizer of the conference is Institute of Mathematics and Mathematical Modelling, Almaty, Kazakhstan. The conference is also supported by Al-Farabi Kazakh National University, Almaty and L. N. Gumilyov Eurasian National University, Astana, Kazakhstan. We would like to thank Institute of Mathematics and Mathematical Modeling, Al-Farabi Kazakh National University and L. N. Gumilyov Eurasian National University for their support. We alsa would like to thank to all invited speakers, International Organizing Committee, International Organizing Committe, and Technical Program Committee Members. With our best wishes and warm regards,

**Chairs:**

**Prof. Allaberen Ashyralyev**

**Prof. Mukhtarbay Otelbaev**

## CONTENTS

70th anniversary OF TYNYSBEK SHARIPOVICH KAL'MENOV 17

# 1 ANALYSIS 23

*Akbota ABYLAYEVA*
Weighted estimates for the integral operator with a logarithmic singularity 24

*Akbota ABYLAYEVA*
Criterion of the boundedness of a fractional integration type operator with variable upper limit in weighted Lebesgue spaces . . . . . 25

*Seyed Masoud AGHAYAN, Ahmad ZIREH*
Common best proximity points in complex valued metric spaces . . . 26

*Meiram AKHYMBEK, Makhmud SADYBEKOV*
On a difference scheme for nonlocal heat transfer boundary-value problem 27

*Gabdolla AKISHEV*
Estimations of the best $M-$ term approximations of functions in the Lorentz space with constructive methods. . . . . 28

*Abdigali SH. AKYSH*
The maximum principle for the Navier-Stokes equations . . . . . 29

*Serik A. ALDASHEV, Erlan T. KITAYBEKOV*
The correctness Dirichlet problem in a cylindrical domain for three-dimensional elliptic equations with type and order extinction . . . . . 30

*Javid ALI*
Relation theoretic metrical fixed point theorems with an application . . 31

*Alexander I. APTEKAREV*
The Steklov problem and estimates for orthogonal polynomials . . . . 32

*Allaberen ASHYRALYEV, Fatih Sabahattin TETİKOGLU*
The positivity of the difference operator with periodic conditions and its applications . . . . . 33

*Anar ASSANOVA*
On the unique solvability of a nonlocal problem with integral conditions for the system of partial differential equations of second order . . . . 34

*Nilupar ATAKHAN*
Asymptotical estimation of boundary value problems for singularly perturbed integro-differential equations . . . . . 35

*Anatoly ATTAEV*
Boundary value problems for loaded hyperbolic equations . . . . . 36

*Sh.BAITUYAKOV, Zh. BAITUYAKOVA, M. ILYASOVA*
On the boundedness of the periodic Hilbert transform on generalized periodic Morrey spaces . . . . . 37

*Sholpan A. BALGIMBAYEVA*
Recovery of partial differential operators on classes of periodic functions with mixed smoothness . . . . . 38

*Feyzi BAŞAR*
Domain of the composition of some triangles in the space of $p-$summable sequences, $(0 < p \leq 1)$ . . . . . 39

*Feyzi BAŞAR, Hüsamettin ÇAPAN*
On some spaces of $q-$bounded double sequences . . . . . 40

*Dauren BAZARKHANOV*
Fourier widths of some function classes associated with $m-$multiple Haar system . . . . . 40

*Turdebek N. BEKJAN*
Noncommutative Holder-type inequalities . . . . . 41

*K.A. BEKMAGANBETOV, YE. TOLEUGAZY*
Interpolation properties of anisotropic Nikol'skii-Besov spaces and embedding theorems . . . . . 42

*Sh. BILALL, S. Kh. SHALGINBAYEVA*
Weighted inequality of Hardy type. . . . . 43

*Nurzhan BOKAYEV, Victor BURENKOV, Dauren MATIN*
On the compactness of set in generalized Morrey spaces . . . . . 44

*Huseyin CAKALLI, Mikail ET, Hacer SENGUL*
A variation on lacunary statistical quasi-Cauchy sequences . . . . . 45

*Ekrem SAVAS, Huseyin CAKALLI*
Ideal statistically quasi Cauchy sequences . . . . . 46

*Nurlan DAIRBEKOV, Oleg PENKIN, Lyazzat SARYBEKOVA*
Solvability of the Dirichlet problem for the $p$-Laplacian on stratified sets . 47

*Muratkhan DAUYLBAYEV, Aziza MIRZAKULOVA*
Asymptotic estimates of the solution of the Cauchy problem for singularly perturbed integro-differential equations . . . . . 49

*Gennadii DEMIDENKO*
Weighted Sobolev spaces, quasielliptic operators and equations not solvable with respect to the highest-order derivative . . . . . 50

*Vasilii N. DENISOV*
The stabilization rate of a solution to the Cauchy problem for parabolic equation with lower order coefficients . . . . . 51

*Gulnar DILDABEK, Isabek ORAZOV*
On the solvability of a nonlocal boundary value problem for the Laplace operator with opposite flows at the part of the boundary . . . . . 52

*Sirojiddin DJAMALOV*
About the correctness to some nonlocal boundary value problem for the equation of the mixed type of the first kind and the second order in space 53

*Zhanat DZHOBULAEVA*
Solution of the linearized nonregular free boundary problem for the parabolic equations . . . . . 54

*Dulat S.DZHUMABAEV, Bayan B. MINGLIBAYEVA*
Properties of an isolated solution to a nonlinear boundary value problem with parameters . . . . . 55

*Abdullah Said ERDOGAN, Ali Ugur SAZAKLIOGLU, Allaberen ASHYRALYEV*
Unique solvability of an inverse problem for a semilinear equation with final overdetermination . . . . . 56

*Nurzhan E. ERZHANOV*
Green's function of a heat problem with a periodic boundary condition . 57

*Leonid FALALEEV*
On the properties of operators with the semi–power type lacunas . . . . 58

*Vladimir E. FEDOROV, Roman R. NAZHIMOV, Dmitriy M. GORDIEVSKIKH*
Initial value problem for a class of fractional order inhomogeneous equations in Banach spaces . . . . . 59

*Ulzada ISKAKOVA*
On an illposed model of oscillations of a flat plate with a variety of mounts on opposite sides . . . . . 60

*Daniyal ISRAFILOV, Ahmet TESTICI*
Approximation properties of some summation methods in the Smirnov classes with variable exponent . . . . . 61

*Symbat KABDRAKHOVA*
Algorithm of finding of a solution bounded on stripe of linear hyperbolic equation with the mixed derivative . . . . . 62

*Bakhtiyor J. KADIRKULOV, Jambul A. SERIKBAEV*
On boundary value problem for the fourth order mixed type equation with fractional derivative . . . . . 63

*Bakhtiyor KADIRKULOV, Jambul SERIKBAEV*
On boundary value problem for the fourth order mixed type equation with fractional derivative . . . . . 64

*Burkhan KALIMBETOV, Dinmukhambet SAPAKOV*
Asymptotics solutions of a singularly perturbed integro-differential problem with rapidly oscillating coefficients . . . . . 65

*Tynysbek KAL'MENOV, Makhmud SADYBEKOV*
On a problem of the Frankl type for an equation of the mixed parabolic-hyperbolic type . . . . . 66

*Aigerim KALYBAY, Saltanat SHALGINBAYEVA*
Weighted multiplicative estimate for norm of discrete Hardy operator . . 67

*Aigerim KALYBAY, Saltanat SHALGINBAYEVA*
Weighted additive estimate for norm of discrete Hardy operator . . . 68

*Huseyin KAPLAN, Huseyin CAKALLI*
Variations on strongly lacunary quasi Cauchy sequences . . . . . 69

*Valery V. KARACHIK, Berikbol T. TOREBEK*
On an Uniqueness and Correct Solvability of the Biharmonic Boundary Value Problem . . . . . 70

*Mahmut KARAKUŞ*
On Generalized Quasi-Convex Bounded Sequences . . . . . 71

*Mahmut KARAKUŞ*
On $AK(S)$ and $AB(S)$ Properties of a $K$-Space . . . . . 71

*Mahmut KARAKUŞ*
On $q^{\lambda}$ and $q_0^{\lambda}$ Invariant Sequence Spaces . . . . . 72

*Shakhobiddin T. KARIMOV*
Solution of a Cauchy problem for iterated generalized two-axially symmetric equation of hyperbolic type . . . . . 73

*Gulden KARSHYGINA, Alua IGENBERLINA, Zauresh SULEYMENOVA*
On the sufficient condition for embedding of generalized Lizorkin-Triebel spaces . . . . . 74

*Ermek KHAIRULLIN*
About one special boundary value problem for multidimensional parabolic integro-differential equation . . . . . 75

*M.KOSHANOVA, K.I.USMANOV, B.TURMETOV*
About solvability of some boundary value problems for Poisson equation in the ball . . . . . 76

*Vladimir V. KOVRIZHKIN*
On weak solutions of the Dirichlet problem for the Tricomi equation . . 76

*Aleksandr KOZHANOV*
Global solvability in some models related to equations of Sobolev type . 77

*Leili KUSSAINOVA, Bakhytty KOSHKAROVA*
Oscillation and nonoscillation of half-linear second order differential equation with alternating potential . . . . . 78

*Damir KURMANBAEV*
Soliton deformations of some minimal surfaces . . . . . 79

*Nurgul KYDYRMINA*
Necessary and sufficient condition that function belong to the Nikol'skii–Morrey space $H^{(r)}M^{\lambda}_{p\,x_1}$ . . . . . 80

*Alisher MATYAKUBOV*
Cauchy problem for a nonlinear degenerate parabolic system not in divergence form . . . . . 81

*Mussakan MURATBEKOV, Sabit IGISINOV, Abai SHYRAKBAEV*
On separability of a differential operator of non-classical type in an unbounded domain . . . . . 82

*Mussakan MURATBEKOV, Madi MURATBEKOV*
On spectral properties of a Shrödinger operator with a negative parameter 83

*Ergashboy MUKHAMADIEV, Alizhon NAIMOV*
On solvability of a class of nonlinear two-point boundary value problems . 84

*Kairat MYNBAEV*
Convergence of kernel density estimators on compact sets . . . . . 85

*Mostepha NACERI, Amir ELHAFFAF*
Existence and nonexistence of positive solutions for singular $n$th-order three-point nonhomogeneous boundary value problem . . . . . 86

*Igor NOVIKOV*
Riecz Lemma and Bernstein Polynomials deformations of some minimal surfaces . . . . . 87

*Erlan NURSULTANOV, Nazerke TLEUKHANOVA*
Summability of the Fourier coefficients of functions from anisotropic Lorentz space . . . . . 88

*Ryskul OINAROV, Ainur TEMIRKHANOVA*
Integral operators with two variable integration limits on the cone of monotone functions . . . . . 89

*Ryskul OINAROV*
Additive and multiplicative weighted estimates of intermediate integral operator . . . . . 90

*Gulaiym ORALSYN*
Inverse coefficient problems for one-dimensional heat transfer with a preservation of medium temperature condition . . . . . 91

*Issabek ORAZOV, Sultanbek AYAZ*
Construction of mathematical models of extraction processes with nonlocal conditions by a spatial variable . . . . . 92

*Kordan OSPANOV*
Nonlocal estimates for solutions of a singular higher order differential equation . . . . . 93

*Kordan N. OSPANOV, Assylbek ZULKHAZHAV*
Coercive solvability of degenerate system of second order difference equations . . . . . 94

*Sergey PISKAREV*
Approximation in time of fractional equations . . . . . 95

*Marina PLEKHANOVA*
Sobolev type equations of time-fractional order with periodical boundary conditions . . . . . 96

*Nedyu POPIVANOV*
Pohozhaev identities and applications for quasilinear equations of mixed elliptic-hyperbolic type . . . . . 97

*Arsen PSKHU*
To the theory of linear ordinary differential equations of fractional order with constant coefficients . . . . . 98

*Asghar RANJBARI*
Some properties of linear operators in locally convex cones . . . . . 98

*Ludmila PULKINA*
Some problems with nonlocal conditions for hyperbolic equations . . . 99

*Nusrat RAJABOV*
To theory one class of integral equation by tube domain . . . . . 100

*Alexander ROGOVOY*
About solutions properties of Tricomi problem for Gellerstedt equation . 101

*Ekrem SAVAS*
On $I_\lambda$-statistical convergence of weight $g$ . . . . . 102

*Ekrem SAVAS*
Asymptotically $\mathcal{I}_\lambda$- Statistical Equivalent Sequences of weight $g$ . . . 103

*Djamila SEBA, Rebai HAMZA*
Weak solutions for nonlinear fractional differential equations with fractional separated boundary conditions in Banach spaces . . . . . 104

*S. SHAIMARDAN*
Hardy-type inequality for a fractional integral operator in q-analysis . . 105

*Yu.R. SHPADI, A.A. KAVOKIN, A.T. KULAKHMETOVA*
On the applicability of Picard's successive approximations to the numerical solving the problem of heat transfer in domain, degenerate at the initial time . . . . . 106

*Yesmukhanbet SMAILOV*
The Hardy-Littlewood type theorems in Besov spaces with the Haar basis 107

*Alexander P. SOLDATOV*
Integrals with homogeneous - difference kernels on the plane and their applications . . . . . 108

*Zhaksylyk TASMAMBETOV*
Confluent hyper geometric function sand two variables Laguerre polynomials as a solutions of Wilczynski type system . . . . . 108

*Svetlana TEMESHEVA*
On one algorithm to find a solution to a linear two-point boundary value problem . . . . . 110

*Balwant Singh THAKUR*
On the convergence of some iterative algorithms to approximate fixed points of nonexpansive mappings . . . . . 111

*Ainur TEMIRKHANOVA*
Oscillation of second order half-linear difference equation and discrete Hardy inequality . . . . . 111

*Agila TLEULESOVA*
Periodic boundary value problem for a system of ordinary differential equations with impulse effects . . . . . 112

*Zhanuzak TOKIBETOV, Gulzhan ABDUAKHITOVA, Ulbike KUSHERBAYEVA*
About sedately growing solutions of one generalized Cauchy-Riemann system . . . . . 113

*Batirkhan Kh. TURMETOV, Berikbol T. TOREBEK*
On a class of fractional elliptic problems with an involution perturbation 114

*Orhan TUĞ, Feyzi BAŞAR*
Four-dimensional generalized difference matrix and some double sequence spaces . . . . . 115

*Orhan TUĞ, Feyzi BAŞAR*
Four-dimensional generalized difference matrix and almost convergent double sequence spaces . . . . . 116

*Faruk UYGUL*
Multiplier Amenability of Banach Algebras . . . . . 117

*Vijay Kumar VYAS*
$q$-Sumudu transforms of basic generalized hypergeometric functions . . 117

*Vladimir VASILYEV*
On periodic wave factorization . . . . . 118

*Aizhan YDYRYS, Nazerke TLEUKHANOVA*
On multipliers of Fourier series in the Lorentz space . . . . . 119

*Alibek YESKERMESSULY*
About the defect indices of fourth-order differential operator with rapidly oscillating coefficients . . . . . 120

*Nurgissa YESSIRKEGENOV*
On a problem for wave equation with data on the whole boundary . . . 121

2 APPLIED MATHEMATICS 123

*Akmaral ABDIGALIYEVA, Zhailaubai ZHUBATOV, Dauren ZHAKEBAYEV*
High order finite-difference method for simulation isotropic turbulence flow 124

*Emmanuel ADEYEFA, R.O. FOLARANMI*
A new class of orthogonal polynomials (Adeolu polynomials) for derivation of initial value solvers . . . . . 125

*D. ALIBIYEV, G. KHAKIMZYANOV, A. KAZHIKENOVA, E. IBRAEVA*
Spreading of flame front I by the predictor-corrector scheme . . . . 126

*Allaberen ASHYRALYEV, Begench GURBANOV, Ramil SALIMOV*
The identification problem for hyperbolic Schrodinger equation . . . 127

*Allaberen ASHYRALYEV, Ramil SALIMOV, Begench GURBANOV*
Fractional derivatives and fractional integrals linked to the fractional powers of positive operators . . . . . 128

*Allaberen ASHYRALYEV, Fatma Songul OZESENLI TETIKOGLU, Tulay KAHRAMAN*
Source identification problem for an elliptic-hyperbolic equation . . . 129

*Allaberen ASHYRALYEV, Baktygul KARABAEVA, Abdizhahan SARSENBI*
Stable difference scheme for the solution of an elliptic equation with the involution . . . . . 130

*Allaberen ASHYRALYEV, Bahriye KARACA*
A note on the Dirichlet problem for model complex partial differential equations . . . . . 131

*Allaberen ASHYRALYEV, Uelker OKUR*
Crank-Nicolson difference scheme for the stochastic parabolic equation with the dependent operator coefficient . . . . . 132

*Kadriye AYDEMİR, O. Sh. MUKHTAROV, Hayati OLĞAR*
Differential operator equations with interface conditions in modified direct sum spaces . . . . . 133

*Anvar AZIMOV, Syrym KASENOV, Daniyar NURSEITOV, Simon SEROVAJSKY*
Inverse problem for the Verhulst equation of limited population growth with discrete experiment data . . . . . 134

*Kheireddine BELAKROUM, Allaberen ASHYRALYEV, Assia GUEZANELAKOUD*
A note on the nonlocal boundary value problem for a third order partial differential equation . . . . . 135

*Galina BIZHANOVA*
Solution of the free boundary problem for the parabolic equations with unknown temperature and velocity . . . . . 136

*M.B. BORIKHANOV, B.Kh. TURMETOV*
On construction of solutions of linear fractional differential equations with constant coefficients . . . . . 136

*Hakima BOUHADJERA*
More general common fixed point theorems under a new concept . . . 137

*Ebru ÇOPUROĞLU, Bahtiyar A. MAMEDOV*
Analytical treatment of two-center overlap integral with respect to same screening constants over Slater-type orbitals . . . . . 137

*Azhibek DASIBEKOV, Azimkhan ABZHAPBAROV, Peruza DUISEBAYEVA, Aigul POLATBEK*
Problems of the theory of the consolidation solved in the special functions 138

*Michael DMITRIEV, Dmitry MAKAROV*
The near optimality of the stabilizing control in a weakly nonlinear system with state-dependent coefficients . . . . . 139

*Yildirim OZDEMIR, Sevilay ERDOGAN*
Numerical solution of the hyperbolic-Schrödinger . . . . . 140

*K. ESHKUVATOV, F. S. ZULKARNAIN, Z. MUMINOV, N. M. A. NIK LONG*
Convergence of Modified Homotopy Perturbation Method for Fredholm-Volterra Integro-Differential Equation of order $m$ . . . . . 141

*Assia GUEZANE-LAKOUD, Rabah KHALDI*
Successive approximations to solve higher order fractional differential equations . . . . . 142

*Farkhondeh HOSSEINI SHEKARABI, Hamed HATAMI*
Operational matrices for finding numerical solution of stochastic differential equations arisen in financial mathematics . . . . . 143

*Djamila ISKENDEROVA, Aibek TOKTORBAEV*
Problem with inhomogeneous boundary values for the equations of magnetic elektrogazodinamik . . . . . 144

*Aizhan ISSAGALI, Auzhan SAKABEKOV*
Principle of Mass Conservation for the Boltzmann's Moment System of Equations in Fourth Approximation . . . . . 145

*Aizhan ISSAGALI, Auzhan SAKABEKOV*
On the well-posedness of the Boltzmann's moment system of equations in fourth approximation . . . . . 146

*V.KARACHIK, S.MASANOV, B.Kh.TURMETOV*
On solvability of some boundary value problems for a biharmonic equation with periodic conditions . . . . . 147

*Syrym KASENOV, Altyn NURSEITOVA, Daniyar NURSEITOV*
A conditional stability estimate of continuation problem for the Helmholtz equation . . . . . 148

*Hossein Jabbari KHAMNEI, Roghaye MAKOUYI*
Recurrence relation for the moments of order statistics from a beta-Pareto distribution . . . . . 149

*Amir KHAN, Gul ZAMAN*
The Motion of Fractional Order Jeffrey Fluid Model . . . . . 150

*Anna KORNEEVA, Vladimir SHAIDUROV*
Computational algorithms for analysis of data from thin-film thermoresistors on a radio-electronic printed circuit board . . . . . 151

*Bakytbek D. KOSHANOV, Gulzhazira D. SMATOVA*
On solvability of boundary value problem of magnetic gas dynamics with cylindric and spherical symmetries . . . . . 152

*Sanjeev KUMAR, Sanjay CHAUDHARY*
Diffusion equation model for the tumors cell density and immune response 153

*Selahattin MADEN, Tamilla I. NASIROVA, Ulviyya Y. KARIMOVA*
Investigation of Laplace transforms for Erlangen distribution of the first passage time into zero level of the semi-Markov random process with positive tendency and negative jumps . . . . . 153

*Iman MAKHDOOM, Parviz NASIRI, Abbas PAK, Masoud YARMOHAMADI*
Statistical inference for the parameter of Burr-type 2 model using vague information as fuzzy data . . . . . 155

*Aigul MANAPOVA*
Differentiation of the functional in an optimization problem for diffusion and convective transfer coefficients of elliptic imperfect contact interface problems . . . . . 156

*H. MOGHADDASI, K. KHALIFEH, A.H.DAROONEH*
Ranking DNA words Using Nonextensive Statistical Mechanics . . . . 157

*O. Sh. MUKHTAROV, Hayati OLĞAR, F. S. MUHTAROV*
Positiveness of second order differential operators with interior singularity 158

*Zahir MURADOGLU*
Transmission conditions for biharmonic equations with discontinuous coefficients and numerical solution of boundary value problem . . . . . 159

*Hayati OLĞAR, O. Sh. MUKHTAROV, Kadriye AYDEMİR*
Some properties of eigenvalues and generalized eigenvectors of one boundary-value problem . . . . . 160

*Nurgul ORUMBAYEVA, Beybitgul SHAYAKMETOVA*
On a method of finding a solution of semi-periodic boundary value problem for hyperbolic equations . . . . . 161

*Yildirim OZDEMIR, Mustafa ALP*
A note on numerical solution of the parabolic-Schrödinger equation . . 162

*Elif OZTURK BEIGMOHAMMADI*
Well-posedness of high order of accuracy difference scheme . . . . . 163

*Alexandr PANOV*
Rank 0 invariant solutions of dynamics of two-phase medium . . . . . 164

*Yildirim OZDEMIR, Esra PEKONUR*
Numerical solution of the elliptic-Schrödinger equation . . . . . 165

*Sergey PYATKOV*
On parabolic inverse problems with overdetermination data on spatial manifolds . . . . . 166

*S.REZAEI, A.H. DAROONEH*
Statistical property of earthquakes network: active and passive points . 167

*Azimbay SADULLAEV, Bakhrom ABDULLAEV*
Monge-Ampere equation and pluripotential theory . . . . . 168

*Auzhan SAKABEKOV, Yerkanat AUZHANI*
Maxwell-Auzhan boundary conditions for Boltzmann's moment system equations in third approximation . . . . . 170

*Kanat SHAKENOV, Marzhan SULTANOVA*
Numerical solution of the relaxational filtration model in relaxationaly-compressed porous environment realized by the linear Darcy law by Monte Carlo and probability difference methods . . . . . 171

*S. SHARIPOV, KUB.S. SHARIPOV, KAD.S. SHARIPOV*
New Economy Form (NEF) as a Plan-Market Economy . . . . . 172

*Alexander L. SKUBACHEVSKII*
Initial–Boundary–Problems for Vlasov–Poisson Equations with Homogeneous Magnetic Field . . . . . 173

*Nurlan TEMIRBEKOV, Amankeldy TURAROV*
Numerical solution of the multiphase one dimensional model of gas lift process . . . . . 173

*Almas TEMIRBEKOV*
Numerical implementation of the method of fictitious domains for elliptic equations . . . . . 175

*A.N. TYUREKHOJAYEV, G.U. MAMATOVA*
The gyroscope movement with variable moments of inertia . . . . . 176

*Koray TÜRK, Allaberen ASHYRALYEV, Deniz AĞIRSEVEN*
On the stability of the telegraph equation with time delay . . . . . 177

*Mesut URUN, Allaberen ASHYRALYEV*
Determination of a control parameter of the $r$-modified Crank-Nicholson difference scheme for the Schrödinger equation . . . . . 178

*Ozgur YILDIRIM, Meltem UZUN*
On the analytical solution methods for sine-Gordon equation . . . . . 179

*Ozgur YILDIRIM, Meltem UZUN*
On the numerical solution of sine-Gordon equation . . . . . 180

*Gulmira K. ZAKIR'YANOVA*
Dynamic analogues of Green and Gauss's formulas for unsteady dynamics of anisotropic elastic medium at antiplane deformation . . . . . 181

*Uzak ZHAPBASBAYEV, Gaukhar RAMAZANOVA, Madina ZHAKENOVA*
Optimization of turbulent flow in a radial reactor with fixed bed . . . 182

*Sailaubay ZHUMATOV*
Instability of control system in the neighborhood of program manifold . 183

*Amir Hadi ZIAIE*
Role of non-commutative geometry in gravitational collapse scenarios . . 184

3 MATHEMATICS EDUCATION 185

*Bayramsoltan ANNADURDYYEVA, Nursoltan NURMYRADOVA*
Geometry study using the interactive whiteboard . . . . . 186

*Maral A. ASHYRALYYEVA, Abdyresul N. CHARYYEV*
Usage of recognizing algorithm handwriting in mathematical education . 187

*Yessen BIDAIBEKOV, Victor KORNILOV, Guldina KAMALOVA, Nagima AKIMZHAN*
Fundamentalization of mathematical knowledge while teaching students inverse and incorrect problem . . . . . 188

*Yessen BIDAIBEKOV, Guldina KAMALOVA, Bektas BOSTANOV, Kairat UMBEBAEV*
Geometric heritage of al-Farabi in education. . . . . 189

*Yessen BIDAIBEKOV, Vadim GRINSHKUN, Guldina KAMALOVA*
Features and advantages of training teachers to use means of informatization to profile teaching mathematics . . . . . 190

*Abdyresul N. CHARYYEV, Maral A. ASHYRALYYEVA*
Value of new technologies in improving the branch of learning . . . . 191

*Burkhan KALIMBETOV, Indira OMAROVA*
Formation of project and research skills of students in calculation of limits 192

*Bahytgul KASKATAEVA, Yssen BIDAIBEKOV, Nurbol ABISHEV*
Methods of professionally-oriented training of complex function theory . 193

*Margarita KLUNNIKOVA, Nikolai PAK, Tatyana PUSHKARYEVA, Tatian STUPINA*
Student-centered model for teaching numerical methods course . . . . 194

*Dosymkhan RAKHYMBEK, Marzhan ABDUALIYEVA, Nurlibai MADIYAROV, Shadyar ALTYNBEKOV*
Didactic bases of system formation of learners' methodological knowledge 195

*Beybitgul SHAYAKHMETOVA, Nurgul ORUMBAYEVA, Sholpan OMAROVA*
Analysis of theoretical and methodological bases of teaching object-oriented programming languages in higher school . . . . . 196

*Simon SEROVAJSKY*
History of mathematics and its teaching in higher school . . . . . 197

*Aigerim ZHANBOLOVA, Bogdat ZHANBUSINOVA, Gaukhar ISKAKOVA, Kelbet SHAUKENOVA, Beybitgul SHAYAKHMETOVA*
Methodological questions of filling the virtual university program modules 197

4 OTHER TOPICS 199

*Nursadyk AKANBAY*
Furstenberg type theorem for a multiplicative product of Markov random matrices . . . . . 200

*Nursadyk AKANBAY, Zoiya SULEYMENOVA, Samal TAPEEVA*
On asymptotic normality of solutions of the Cauchy problem of a parabolic equation with random right side . . . . . 201

*Valentina ALEKHINA, Ilya STARODUMOV, Oleg STARODUMOV*
Modeling of thorium transport by the water flow in the river Kechi-Kemin 202

*Bektur BAIZHANOV*
TOne-types in ordered theories . . . . . 203

*Cigdem GUNDUZ ARAS, Sadi BAYRAMOV*
On the Tietze extension theorem in soft topological spaces . . . . . 204

*Cigdem GUNDUZ ARAS*
A study on intuitionistic fuzzy supra soft topological spaces . . . . . 205

*Yu. FATEEV, D. DMITRIEV, V. TYAPKIN, N. KREMEZ, V. SHAIDUROV*
The probability distribution function for the sum of squares of independent random variables . . . . . 206

*Assylbek ISSAKHOV*
Hyperimmunity and $A$-computable universal numberings . . . . . 207

*Seyit KERIMKHULLE*
A system dynamics model to forecasting of the United States gross savings 208

*Ljubiša D.R. KOČINAC*
Selections, games and Karamata theory . . . . . 209

*Damir KULBAEV, A.Z. ABDENOV*
Application of the theory of games for selection of effective defense. . . 210

*Beibut KULPESHOV*
Binarity for almost $\omega$-categorical quite o-minimal theories . . . . . 211

*Ainur KUSSAINOVA, Tamara ZHUKABAYEVA, Abai JUSSUPOV, Kazizat ISKAKOV, Zhanar ORALBEKOVA*
Prospective study on the potential of big data in railways . . . . . 212

*Bahtiyar Akber MAMEDOV, Elif SOMUNCU, Iskender M.ASKEROV*
An Efficient Method for the Determination of Fourth Virial Coefficient with Lennard-Jones (12-6) Potential and Its Application . . . . . 212

*K. MOTARJEM, M. MOHAMMADZADEH, A. ABYAR*
Analyzing High-Dimensional Survival Data Using Random Forests . . . 213

*Ilya STARODUMOV, Nikolai KROPOTIN*
Features in simulation of crystal growth using the hyperbolic PFC equation and the dependence of the numerical solution on the parameters of the computational grid . . . . . 214

*Iskander A. TAIMANOV*
The Moutard transformation of two-dimensional Dirac operators and the Mobius geometry . . . . . 216

*Marat TLEUBERGENOV, Gulmira VASSILINA*
On stochastic stability of the integral manifold under permanently acting random perturbations. . . . . 217

*Ualbai UMIRBAEV*
Automorphisms of differential polynomial algebras . . . . . 218

*Vassilly VOINOV, Natalya PYA, Rashid MAKAROV, Yevgeniy VOINOV, Roza RAKHIMOVA*
Combining goodness of fit tests for multivariate normality . . . . . 219

*Aibat YESHKEYEV, Olga ULBRIKHT*
The stability of forcing companion for center of Jonsson set's fragment . 220

*Aibat YESHKEYEV, Nazgul SHAMATAYEVA*
Categorical Jonsson fragments in existentially prime convex Jonsson theory . . . . . 221

*Aibat YESHKEYEV, Maira KASSYMETOVA*
Strongly minimal fragments in existentially prime convex Jonsson theory . 222

5 MS1:MATHEMATICAL MODELLING 223

*Kamil AIDA-ZADE, Anar RAHIMOV*
Numerical solution to inverse source problems for hyperbolic equation . . 224

*Aidyn AITZHAN, Medet INKARBEKOV, Asetzhan KOLDAS, Aidarkhan KALTAYEV*
Implementation of scalar filtered density function for large eddy simulation of turbulent reacting flow using a high-order discontinuous Galerkin method . . . . . 225

*Lyudmila ALEXEYEVA, Makpal AHMETZHANOVA, Gulnar KAYSHIBAYEVA*
Dynamics of the massif in the vicinity of the tunnel any profile of section under action of transport loadings . . . . . 226

*F.A. ALIEV, N.S. HAJIEVA, N.A. SAFAROVA, M.F. RAJABOV*
An asymptotical method for determining hydraulic resistance coefficient of gas-lift process . . . . . 227

*Tulegen AMANBAEV, Gamidulla TILLEUOV, Bibigul TULEGENOVA*
Modeling and calculation of flotation process in one-dimensional formulation . . . . . 228

*Mersaid ARIPOV*
The Fujita type a critical exponent for a double nonlinear parabolic equation and system . . . . . 229

*Nurgali ASHIRBAYEV, Zhansaya ASHIRBAYEVA, Azimkhan ABZHAPBAROV, Manat SHOMANBAYEVA*
The features of a non-stationary state of stress in the elastic multisupport construction . . . . . 230

*Nurgali ASHIRBAYEV, Zhansaya ASHIRBAYEVA, Turlybek SULTANBEK, Raina BEKMOLDAYEVA*
Modeling and solving the two-dimensional non-stationary problem in an elastic body with a rectangular hole . . . . . 231

*Galitdin B. BAKANOV*
On the stability estimation of differential-difference analogue of the integral geometry problem with a weight function . . . . . 232

*Arzu ERDEM COSKUN, Cigdem GUNDUZ ARAS, Ayse SONMEZ, Huseyin CAKALLI*
Soft matrices on soft multisets in an optimal decision process . . . . 233

*Evgeny DEREVTSOV, Svetlana MALTSEVA,Ivan SVETOV, Murat SULTANOV*
On a problem of reconstruction of a discontinuous function by its Radon transform . . . . . 234

*Medet INKARBEKOV, Aydyn AYTZHAN, Aidarkhan KALTAYEV*
A new combined DG-MC solver for large eddy simulation of reacting turbulent flows . . . . . 235

*Bakhtiyar ISMAILOV, Khairulla ISMAILOV, Saule MELDEBEKOVA*
On a method of mathematical modeling of the pyrolysis process to produce nanomaterials from natural gas . . . . . 236

*Alibek ISSAKHOV*
Mathematical modeling of the thermal effect on the aquatic environment from thermal power plant by using two water discharged pipes . . . . . 237

*Ibragim KALIEV, Saltanbek MUKHAMBETZHANOV, Gulnara SABITOVA, Anghyz SAKHIT*
Mathematical modeling of non-equilibrium sorption . . . . . 238

*Alex KAVOKIN, Targyn NAURYZ, Nazerke BIZHIGITOVA*
Exact solution of two phase spherical Stefan problem with two free boundaries . . . . . 239

*Aigul Sh. KAZHIKENOVA, D.B.ALIBIYEV, E.S. IBRAYEVA*
Mathematical modeling of the temperature dependence of molten viscosity of metal according to the concept of chaotized particles . . . . . 240

*Stanislav N.KHARIN, Merey M.SARSENGELDIN, Samat KASSABEK*
The mathematical models of electromagnetic field dynamics and heat transfer in closed electrical contacts including Thomson effect . . . . . 241

*Khonatbek KHOMPYSH, Sharypkhan SAKHAEV*
Solvability of a stationary problem of magnetohydrodynamics . . . . . 242

*Sonu LAMBDA*
Mathematical modelling and control theory analysis of artificial satellite . 243

*Vladimir LEVIN, Pavel GEORGIEVSKIY*
New effects for high speed collision of blunt bodies with atmospheric inhomogeneities . . . . . 244

*Vladimir LEVIN, Ivan MANUYLOVICH, Vladimir MARKOV*
Initiation of 3D Detonation in Supersonic Flows in Channels of Special Geometry . . . . . 246

*Altyn MAKASHEVA, Altynshash NAIMANOVA*
Supersonic multi-species flow with particle dispersion on a mixing layer . 247

*Ernazar NYSSANOV, Marina SAPRYGINA*
Modeling and solving an optimization problem of supply of chemical raw materials . . . . . 248

*Anna PYRKOVA, Anatoly IVASHCHENKO, Raigul NIYAZOVA*
Fragmenting of algorithm for solving the problem of clustering miRNA families . . . . . 249

*Bolatbek RYSBAIULY, Abilmazhin ADAMOV*
Iterative method of finding hydraulic conductivity characteristics of soil moisture . . . . . 250

*Makhmud SADYBEKOV, Abdizhahan SARSENBI, Aizhan TENGAYEVA*
Spectral properties of a Laplace operator with Samarskii–Ionkin type boundary conditions in a disk . . . . . 251

*Makhmud SADYBEKOV, Nurgissa YESSIRKEGENOV*
Spectral properties of a Laplace operator with Samarskii–Ionkin type boundary conditions in a disk . . . . . 252

*Merey M.SARSENGELDIN, Stanislav N.KHARIN, Hassan NOURI*
Analytical solution of two-phase spherical Stefan problem by heat polynomials and integral error functions . . . . . 253

*V. SHAIDUROV, Viktoriya KORNIENKO, A. VYATKIN*
The mathematical modeling of grouping the dipole water clusters . . . 254

*Nurtoleu SHAKHAN, Asel BEKETAEVA, Altynshash NAIMANOVA*
Simulation of shock wave boundary layer interaction in flat channel with jet injection . . . . . 255

*R.Y. SHIKHLINSKAYA, S.M. SALIMOV*
Application of of fuzzy logic for matching modes use of wind power plant with an electrical load schedule . . . . . 256

*Ivan SVETOV, Svetlana MALTSEVA, Anna POLYAKOVA*
Numerical solution of $2D$-vector tomography problem using the method of approximate inverse . . . . . 257

*Kanat TULENOV*
Duality property of the noncommutative $\ell_\infty$ and $\ell_1$ valued symmetric Hardy spaces. . . . . 258

*Kanat TULENOV, Dostilek DAUITBEK*
Lateral boundary conditions for the Klein-Gordon-Fock equation . . . 259

*Ainur ZHUMALI*
A modelling the dynamics of in-situ leaching process at the microscale . 260

6 MS2:SPECTRAL THEORY OF DIFFERENTIAL EQUATIONS 261

*Aizat AIMAKHANOVA, Saltanat SHALGINBAYEVA, Lyazzat ZHUMANOVA*
The first regularized trace of integro-differential Sturm-Liouville operator on the segment with punctured points at integral perturbation of transmission conditions . . . . . 262

*Shavkat ALIMOV*
On the solvability of differential equations in logarithmic scale of Hilbert spaces . . . . . 263

*Meiramkul AMANGALIYEVA, Muvasharkhan JENALIYEV, Kanzharbek IMANBERDIYEV, Murat RAMAZANOV*
On spectral problems for loaded two-dimension Laplace operator . . . 264

*Meiramkul AMANGALIYEVA, Muvasharkhan JENALIYEV, Madi ERGALIYEV, Murat RAMAZANOV*
Spectrum of Volterra integral operator of the second kind . . . . . 265

*Gaukhar AREPOVA*
On a regular problem for an elliptic-parabolic equation with a potential boundary condition . . . . . 266

*Allaberen ASHYRALYEV, Abdulgafur TASKIN*
Structure of fractional spaces generated by the two dimensional neutron transport operator. . . . . 267

*Allaberen ASHYRALYEV, Abdisalam SARSENBI*
Green's function of the second order differential operator with involution 268

*Bazarkan N. BIYAROV Gulnara K. ABDRASHEVA*
Bounded perturbations of the correct restrictions and extensions . . . 269

*Ziganur FAZULLIN, Irina NUGAEVA*
On spectrum of perturbed two-dimensional harmonic oscillator in a strip . 270

*Tuba GULSEN, Etibar S. PANAKHOV*
Dirac systems that contain discontinuity conditions . . . . . 271

*Nurlan IMANBAEV, Makhmud SADYBEKOV*
Stability of basis property of a type of problems with nonlocal perturbation of boundary conditions. . . . . 272

*Muvasharkhan JENALIYEV, Murat RAMAZANOV*
On a homogeneous parabolic problem in an infinite corner domain . . . 273

*Tynysbek Sh. KALMENOV, Durvudkhan SURAGAN and Bolys SABITBEK*
Spectral geometry inequalities for Schatten $p$-norms of compact operators 274
*Tynysbek KAL'MENOV, Gaukhar AREPOVA*
On a heat transfer model for the locally inhomogeneous initial data . . 275
*Baltabek KANGUZHIN, Niyaz TOKMAGAMBETOV*
A uniqueness theorem of a boundary inverse problem of a differential operator on an interval with integro–differential boundary conditions . . . 276
*Baltabek KANGUZHIN, Gulzat NALZHUPBAYEVA*
On identities for eigenvalues of a well-posed pertubation of the Laplace operator in a punctured domain . . . . . 277
*Baltabek KANGUZHIN, Dostilek DAUITBEK*
On first eigenvalue of Laplace operator . . . . . 278
*Baltabek KANGUZHIN, Lyailya ZHAPSARBAYEVA*
Estimates of the eigenvalues of operator arising in swelling pressure model 279
*Akylbek KERIMBEKOV*
The resolvent equation of nonlinear Fredholm integral equations . . . 280
*Leonid KRITSKOV, Abdizhahan SARSENBI*
On Spectral Properties of Perturbations of the Operator $-u''(-x)$ with Initial Data . . . . . 281
*Daulet NURAKHMETOV*
Well-posed problems for the Laplace operator in the multiply connected domains . . . . . 282
*Michael RUZHANSKY*
Very weak solutions to wave equations . . . . . 283
*Bolys SABITBEK, Durvudkhan SURAGAN*
An embedding theorem of Sobolev type spaces on stratified Lie groups . 283
*Makhmud A. SADYBEKOV, Aidyn A. KASSYMOV*
An isoperimetric inequality for heat potential and heat equation . . . 284
*Makhmud A. SADYBEKOV, Berikbol T. TOREBEK*
On some spectral inequalities for a nonlocal elliptic problem . . . . . 285
*Makhmud SADYBEKOV, Abdizhahan SARSENBI, Aizhan TENGAYEVA*
Spectral properties of a Laplace operator with Samarskii–Ionkin type boundary conditions in a disk . . . . . 286
*Makhmud SADYBEKOV, Nurgissa YESSIRKEGENOV*
Spectral properties of a Laplace operator with Samarskii–Ionkin type boundary conditions in a disk . . . . . 287
*Abdizhahan SARSENBI*
The theorem on the basis property of eigenfunctions of second order differential operators with involution . . . . . 288
*Amir SHALDANBAYEV, Manat SHOMANBAYEVA, Asylzat KOPZHASSAROVA*
Solution of a singularly perturbed Cauchy problem for linear systems of ordinary differential equations by the method of spectral decomposition . 289
*Andrei SHKALIKOV*
Differential operators with singular coefficients . . . . . 290
*Aizhan TENGAYEVA, Gulnar DILDABEK*
Existence of eigenvalues of problem with shift for an equation of parabolic-hyperbolic type . . . . . 291
*Kanat TULENOV*
Duality property of the noncommutative $\ell_\infty$ and $\ell_1$
valued symmetric Hardy spaces. . . . . 292
*Kanat TULENOV, Dostilek DAUITBEK*
Lateral boundary conditions for the Klein-Gordon-Fock equation . . . 293
*Nurgissa YESSIRKEGENOV*
Euler-Hilbert-Sobolev spaces on homogeneous groups . . . . . 294

7 MS3:DINAMICAL SYSTEMS 295

*Nurlan ABIEV*
On a dynamical system relating to the Ricci flow on Aloff-Wallach spaces 296

*Serikbai A. AISAGALIEV, A.M. AYAZBAYEVA*
Investigation on absolute stability of regulated systems in a simple critical case . . . . . 297

*Serikbai A. AISAGALIEV, A.M. AYAZBAYEVA*
Controllability of process described by system of linear integro-differential equations with resrtictions . . . . . 298

*Daulet Sh. AKHMEDOV, Almat S. RASKALIYEV*
Localization of an air target by means of GNSS-based multistatic radar . 299

*Hikmet AKHUNDOV, Mutallim MUTALLIMOV*
Gradient in the optimal control problem described by the differential equations . . . . . 300

*F.A. ALIEV, N.A. ALIEV, A.P. GULIEV, K.G. GASIMOVA*
Time frequency method of solving one boundary value problem for a hyperbolic system and its application to the oil extraction . . . . . 301

*Allaberen ASHYRALYEV, Kadriye Tuba TURKCAN, Mehmet Emir KOKSAL*
On Numerical Solutions of Telegraph Equations with the Dirichlet Boundary Condition . . . . . 302

*Guzel BAYBULATOVA*
Optimal control problem for the degenerate phase field system of equations 303

*Yulia DANIK*
About the robustness of the middle stabilizing controller for quasi-linear state dependent coefficients discrete-time systems . . . . . 304

*F.G. FEYZIEV, G.H. MAMMADOVA, F.N. NABI-ZADE*
On a controllability for a class of $2D$-linear modular dynamic systems with periodic parameters . . . . . 306

*Allaguly GURBANLYYEV*
On the systems of rational difference equations . . . . . 307

*Zainel MURZABEKOV*
Design of PI regulators for dynamic systems with constrained control and fixed endpoints of trajectories . . . . . 308

*Zainel MURZABEKOV, Shamshi AIPANOV, Saltanat USUBALIEVA*
Optimal control problem for the three-sector economic model of a cluster . 309

*Zhussip SULEIMENOV*
The method of accelerated convergence for constructing conditional-periodical solutions . . . . . 310

*Mikhail M. TUROV*
Martinet-Ramis modulus for one Quadratic System . . . . . 311

*Kuralay YESMAKHANOVA, Gaukhar SHAIKHOVA, Guldana BEKOVA*
Soliton solutions of the Hirota equation . . . . . 312

AUTHOR'S INDEX 313

# TO 70th ANNIVERSARY

**May 2016 was the 70th anniversary of Tynysbek Sharipovich Kal'menov, the outstanding Kazakhstani mathematician, the academician of the National Academy of Sciences of the Republic of Kazakhstan.**

**He was born in the South-Kazakhstan region of the Kazakh SSR, in the village Koksaek of Lenger district, on May 5, 1946.**

Tynysbek Sharipovich is a graduate of the Novosibirsk State University (1969) and a representative of the school of A.V. Bitsadze, the outstanding scientist, the member-correspondent of the Academy of sciences of the USSR. Scientific researches held by T.Sh. Kal'menov in postgraduate years (1969 - 1972) were devoted to studying the well-posedness of initial and initial-boundary value problems for degenerated hyperbolic equations of the form

$$k(y)u_{xx} - u_{yy} + au_x + bu_y + cu = f; \quad k(0) = 0; \ k(y) > 0, y \neq 0.$$

For this general equation T.Sh. Kal'menov found sufficient conditions of the well-posedness of the first and second Darboux problem including all the previously known conditions for the well-posedness. For various model cases of this equation there were first found: a criterion of the uniqueness of a solution; sufficient conditions of the well-posedness and a class of smoothness of a solution of the Darboux problem; a criterion for the continuity of a solution of the Goursat problem, and in this case it was first indicated on a different impact of characteristics for the well-posedness of a problem; a new characteristic Cauchy problem (incomplete Goursat problem) was formulated and its well-posedness in a class of analytical functions was proved.

From 1972 till 1985 T.Sh. Kal'menov worked in the Institute of Mathematics of the Academy of Sciences of the KazSSR, where he made a career from a junior researcher to the head of laboratory. In these years the field of his scientific interests belonged to theory of differential operators and he obtained significant results.

In case of boundary value problems for equations of hyperbolic and mixed types in researching problems in the weak sense there arise sufficient difficulties in clarifying the relation between generalized and classical solutions and, as a result, the lack of proof of the uniqueness of solutions, hence there is the one-sided investigation of problems. In this regard, the following question is of great importance: Under what conditions weak solutions are classical. In a more precise setting, when the generalized solution $u$ of a boundary problem for the equation $Lu = f$ can be approximated by the sequence $u_n \in W_2^2(\Omega)$ of the classical solutions in the sense that $u_n \to u$ and $Lu_n \to f$ in some metric, as a rule, by the norm $L_2(\Omega)$. The generalized solutions having the last above-mentioned property is called *strong*.

In this direction T.Sh. Kal'menov proved the well-posedness of the semiperiodic Dirichlet problem for the degenerated equation of the mixed type

$$Lu = k(y)u_{xx} + u_{yy} + a(y)u_x + b(y)u_y + c(y)u = f(x, y);$$

and for the degenerated equation

$$Lu = signy|y|^m u_{xx} + u_{yy} = f(x, y).$$

He showed the existence of strong solutions of the Darboux and Tricomi problems.

For the Lavrent'ev–Bitsadze equation

$$Lu = -signy \ u_{xx} - u_{yy} = f(x, y),$$

T.Sh. Kal'menov proved the criterion for strong solvability of the Tricomi problem in spaces $L_p$, in terms of approach angles of an elliptical part of the boundary to a line changing the type. In this case, in contrast to previously known works, non-uniqueness of weak solutions was shown for the first time and criteria for the strong solvability of these problems were given.

Developing the operator approach, T.Sh. Kal'menov constructed the general theory of regular boundary value problems for equations of hyperbolic and mixed types. The classical construction of a boundary value problem is as follows: an equation and a boundary condition are given. It is necessary to investigate the solvability of this problem and properties of the solution if it exists ( in the sense of belonging to some space). Beginning with the papers of J. von Neumann and M.I. Vishik, there exists another more general approach: an equation and a space are given, right-hand parts of the equation and boundary conditions, and a solution must belong to this space. It is necessary to describe all the boundary conditions, for which the problem is correctly solvable in this space.

T.Sh. Kal'menov described all the well-posed boundary value problems: for a wave equation in a characteristic triangle, for a multidimentional wave equation in a characteristic cone, for a Lavrent'ev-Bitsadze equation, for a Tricomi equation and for a multidimensional equation of mixed type. For the first time a multidimensional correct analogue of the Darboux problem was constructed, and its solution in the explicit form was given.

A large series of papers of T.Sh. Kal'menov is devoted to the spectral theory of equations of hyperbolic and mixed types. In contrast to the theory of solvability, spectral problems for the equations of hyperbolic and mixed types are poorly understood. Note that the well-known methods (in particular, the abstract spectral theory of linear operators), being a powerful tool in the study of elliptic operators, are little adapted in application to the boundary value problems for equations of hyperbolic and mixed types in the domain, part of which coincides with a characteristic cone.

For this reason, many actual problems of equations of hyperbolic and mixed types require special studies and attracting new tools and techniques. Some of such problems, in particular, are the questions of spectrum and strong solvability of local and non-local non-elliptic problems. The systematic study of spectral problems for mixed type equations started recently with the work of T.Sh. Kal'menov, E.I. Moiseev, S. Ponomarev.

T.Sh. Kal'menov formulated and proved the extremum principle for equations of mixed type, having now in literature the name "the maximum principle of Kal'menov". On the basis of this new principle of extremum he solved the problem of the existence of an eigenvalue of the Tricomi problem (this problem is not Volterra), and this fact laid the basis of a new perspective scientific direction - the spectral theory of equations of mixed type. The existence of eigenvalues of the Tricomi problem was proved for both a Lavrent'ev–Bitsadze problem and a general Gellerstedt's problem.

All these results became the basis for the doctoral thesis on "On regular boundary value problems and spectrum for equations of hyperbolic and mixed types" , defended at the Moscow State University named after M.V. Lomonosov in 1983.

From 1985 till 1991 T.Sh. Kal'menov worked as the dean of mathematical faculty of the Kazakh State University, from 1991 till 1997 he was the rector of the Kazakh Chemical-Technological University, the South-Kazakhstan Technical University. In the period from 1998 till 2003 he was the head of department of the South-Kazakhstan State University.

In these years T.Sh. Kal'menov appealed to problems in more general setting. After the appearance of the famous paper of A.V. Bitsadze and A.A.

Samarskii, nonlocal problems of mathematical physics are increasingly attracting the attention of mathematicians. Such problems are significantly nonself-adjoint and classical well-developed investigation methods are not applicable for their investigation. For the general elliptic equation

$$Lu = -\sum_{i,j=1}^{n} a_{ij}(x)\frac{\partial^2 u}{\partial x_i \partial x_j} + \sum_{i=1}^{n} a_i(x)\frac{\partial u}{\partial x_i} + a(x)u$$

in an arbitrary domain with smooth boundary, the completeness of root vectors of the main Bitsadze-Samarskii problems in general setting was proved by T.Sh. Kal'menov for the first time.

In 2004 T.Sh. Kal'menov returned to Almaty, where he became the General director of the Center of Physical-Mathematical Researches of the Ministry of Education of the Republic of Kazakhstan, reformed into the Institute of Mathematics, Informatics and Mechanics, then into the Institute of Mathematics and Mathematical Modeling in 2012. During these years T.Sh. Kal'menov obtained new significant results concerning classical problems of partial equations.

In the theory of mathematical physics, for example, in the theory of elasticity, an explicit representation of a solution of problems for an inhomogeneous biharmonic equation takes an important place. In papers of T.Sh. Kal'menov a polyharmonic Greens function of the Dirichlet problem

$$\triangle_x^m u(x) = f(x), f(x) \in L_2(\Omega_\delta),$$

$$\frac{\partial^i}{\partial n_x^i} u|_{|x|=\delta} = 0, \quad i = 0, 1, 2, ..., m-1$$

for the first time was constructed in the explicit form (in the terms of elementary functions) in the ball $\Omega_\delta \subset \mathbb{R}^n$ for the case of an arbitrary dimension of the space.

The spectral theory of non-selfadjoint differential operators is not complete yet even for a case of simplest ordinary differential equations. For partial equations and all the more for equations of the mixed type in non-selfadjoint spectral theory only some separate results have been obtained. Up to date the example of regular boundary value problem for differential equations is not found, a spectrum of this problem (except an empty set) is a finite set. T.Sh. Kal'menov was the first to establish that the spectrum of differential operators of the general form

$$Lu = \sum_{|\alpha| \le p} a_\alpha(x) D^\alpha u = \lambda u(x),$$

generated by regular boundary conditions, is either empty, or an infinite set.

The classical volume Newton potential

$$u(x) = \varepsilon_n * f \equiv \int_\Omega \varepsilon_n(x-y) f(y) dy = L^{-1} f$$

defines the value of mass or charge distributed in $\Omega$ with density $\frac{f}{(n-2)\sigma_n}$. This potential also is often used in the theory of functions and in the embedding theorems. Since, the fundamental solution $\varepsilon_n(x-y)$ is symmetric and real-valued, and has a weak singularity, then the integral operator $L^{-1}$ is the completely continuous self-adjoint operator in $L_2(\Omega)$ and the function $u(x) = \varepsilon_n * f(x)$ satisfies the Poisson equation $-\triangle u = f$.

In papers of T.Sh. Kal'menov the boundary condition for the volume Newton potential:

$$\frac{1}{2}u(x) = -\int_{\partial\Omega} \varepsilon_n(x-y)\frac{\partial u(y)}{\partial n_y}dS_y + \int_{\partial\Omega} \frac{\partial \varepsilon_n(x-y)}{\partial n_y}u(y)dS_y,$$

$$\text{for all } x \in \partial\Omega.$$

for the first time was found.

It turned out that these boundary conditions describe the long-known in theoretical physics effect of "transparent boundary conditions" , skipping the outgoing waves and reflecting the incoming waves. The presence of such boundary conditions of the volume potential permits reducing of the problem with conditions of Sommerfeld type radiation in the infinite domain to a problem in a bounded domain and effectively apply numerical methods.

These results were further developed on a case of heat and wave potentials as well as in non-cylindric domains.

The classical Hadamard example demonstrates the ill-posedness of the Cauchy problem for a Laplace equation. The outstanding Soviet mathematicians - the academicians A.N. Tikhonov and M.M. Lavrent'ev, their disciples and followers found the conditions of well-posedness of the Cauchy problem for the Laplace equation, and of the other ill-posed problems. Also regularization methods for ill-posed problems were constructed. In papers of T.Sh. Kal'menov by the method for the expansion on eigenfunctions of a mixed Cauchy problem for the Laplace equation with a deviating argument we established necessary and sufficient well-posedness condition in the domain $\Omega = \{(x,t) : 0 < x < \pi, -1 < t < 1\}$ of the mixed Cauchy problem for the Poisson equation:

$$Lu = u_{tt}(x,t) + u_{xx}(x,t) = f(x,t), (x,t) \in \Omega,$$

$$u|_{t=-1} = \varphi_1(x), u_t|_{t=-1} = \varphi_2(x),$$

$$u|_{x=0} = 0, u|_{x=\pi} = 0.$$

Furthermore, it is shown that the solution of the Cauchy problem is unique and without setting of boundary conditions.

A characteristic feature of the mathematical creation of T.Sh. Kal'menov is the plenty of original methods and non-standard approaches to solving mathematical problems. T.Sh. Kal'menov belongs one of the few scientists managed to leave a mark almost in all fields of mathematics that he has studied. He is the worthy successor to his teacher Andrei Vasil'evich Bitsadze, the member-correspondent of AS SSSR and the bright continuer of traditions of the Soviet mathematical school.

More than 50 master's theses and 11 doctoral theses have been defended under supervision of T.Sh. Kal'menov. He has more than 150 published scientific works, a great amount of which have been published in international mathematical journals. Among them 11 works have been published in the journal "Proceedings of the USSR Academy of Sciences / Doklady Mathematics". He has published 14 works in highly rated international journals for last 5 years.

T.Sh. Kal'menov has been honored by a number of high awards and titles for his longstanding scientific, pedagogical and social organizational activity. In 1978 he was honored with the premium of Lenin Komsomol of Kazakhstan, in 1996 he became the Honored Worker of Science and Technology of the Republic

of Kazakhstan. In 1989 he was elected as a member-correspondent of the Academy of Sciences of Kazakh SSR, and in 2003 he became an academician of the National Academy of Sciences of the Republic of Kazakhstan.

T.Sh. Kal'menov was honored with orders and medals of the Republic of Kazakhstan, and in 2013 he was awarded the State premium in the field of science and technology for the series of papers "To the theory of initial-boundary value problems for differential equations".

The academician T.Sh. Kal'menov as professional scientist and the head of The Institute of Mathematics and Mathematical Modeling is demanding of himself and of all scientists, that helps to keep the bar of mathematical science in Kazakhstan at a high international level. At the present moment he is one of the brighest mathematician of Kazakhstan and its principal leader.

With unflagging energy T.Sh. Kal'menov is engaged in scientific work, generously presents scientific ideas to his disciples and colleagues, realizes all new creative plans. He is full of new and original mathematical ideas.

It is wonderful that T.Sh. Kal'menov is fruitful and in the family life. He is the successful father of ten children, and now he is bringing up more than twenty five grandchildren!

# ANALYSIS

# Weighted estimates for the integral operator with a logarithmic singularity

Akbota ABYLAYEVA
L.N. Gumilyov Eurasian National University,
010008 Astana, Satpayev str., 2, Kazakhstan
E-mail: abylayeva_b@mail.ru

**Abstract:** Let $0 < q < \infty$, $1 < p < \infty$, $\frac{1}{p} + \frac{1}{p'} = 1$, $R_+ = (0, \infty)$, $U : R_+ \to R$ and $V : R_+ \to R$ are wheighted functions, i.e. non-negative measurable functions on $R_+$.

Since the 70-s of the last century the weighted estimate of the form

$$\|UKf\|_q \le C\|Vf\|_p \tag{1}$$

intensively studied in the world mathematical literature for different classes of the operators $K$, where $\|\cdot\|_p$ is the usual norm of the space $L_p \equiv L_p(R)$. Some directions of the research of the estimate (1) until 2000 year for the integral operators are summarized in the books [1], [2], [3]. In the paper [4] the estimate (1) was studied for some classes of the operator $Kf$ and also a full description of weights $U$ and $V$ was given there.

In this paper we study the estimate (1) for the singular operator in the form $Kf(x) = \int\limits_0^x \ln \frac{x}{x-s} f(s)ds$, when $V(x) = x^{1-\gamma}$, $\gamma > 0$.

**Theorem.** *Let $1 < p \le q < \infty$, $\gamma > \frac{1}{p}$. The inequality (1) holds if and only if*

$$A = \sup_{x>0} x^{\gamma+\frac{1}{p'}} \left( \int\limits_x^\infty U^q(t)t^{-q}dt \right)^{\frac{1}{q}} < \infty,$$

*wherein $C \approx A$, where $C$ is the best constant (1).*

**Keywords:** integral operator with logarithmic singularity, weighted estimate.

**2010 Mathematics Subject Classification:** 47A63, 47G10

**References:**

[1] Opic B., Kufner A., "Hardy-type inequalities", *Pitman Research Notes in Math. Series. Longman Scientific and Technical*, Horlow, 1990.

[2] Kufner A., Persson L-E., "Integral inegualities with weights", Prague V.211, 2000.

[3] Kufner A., Maligranda L., Persson L.-E. "The Hardy inequality -about its history and some related results, *University of West Bohemia, Plzen*, p. 152, 2007.

[4] Oinarov R., "Boundedness and compactness of integral operators Voltaire types", *Sib. Math. J.*, Vol.48, No.5, pp. 1100-1115, 2007 (in Russian).

## Criterion of the boundedness of a fractional integration type operator with variable upper limit in weighted Lebesgue spaces

Akbota ABYLAYEVA
L.N. Gumilyov Eurasian National University,
010008 Astana, Satpayev str., 2, Kazakhstan
E-mail: abylayeva_b@mail.ru

**Abstract:** Let $0 < p, q < \infty$, $I = (a, b)$, $-\infty \le a < b \le \infty$, $0 < \alpha < 1$, $\frac{1}{p} + \frac{1}{p'} = 1$, $R_+ = (0, \infty)$, $W : I \to R$ is a nondecreasing, locally absolutely continuous function and $W(a) = \lim\limits_{t \to a^+} W(t) > -\infty$. $\frac{dW(t)}{dt} = w(t)$, $v : I \to R$ is a non-negative locally integrable function on $I$ and $\varphi : I \to I$ is a strictly increasing locally absolutely continuous function with the properties $\lim\limits_{x \to a^+} \varphi(x) = a$, $\lim\limits_{x \to b^-} \varphi(x) = b$, $\varphi(x) \le x$, $\forall x \in I$. Consider the operator in a form $K_{\alpha,\varphi} f(x) = \int\limits_a^{\varphi(x)} \frac{f(s) w(s) ds}{(W(x) - W(s))^{1-\alpha}}$, $x \in I$ from $L_{p,w} = L_{p,w}(I)$ to $L_{q,v} = L_{q,v}(I)$, where $L_{p,w}$ is a space of measurable functions $f : I \to R$ for which the functional $\|f\|_{p,w} = \left( \int\limits_a^b |f(x)|^p w(x) dx \right)^{\frac{1}{p}}$, $0 < p < \infty$ is finite. In the case $\varphi(x) \equiv x$ the operator $K_{\alpha,\varphi}$ is studied in papers [1], [2].

**Theorem.** The operator $K_{\alpha,\varphi}$ is bounded from $L_{p,w}$ to $L_{q,v}$ if and only if

1) $A = \sup\limits_{a<z<b} \left( \int\limits_z^b W^{q(\alpha-1)}(x) v(x) dx \right)^{\frac{1}{q}} W^{\frac{1}{p'}}(\varphi(x)) < \infty$, for $1 < p \le q < \infty$ and $\frac{1}{p} < \alpha < 1$, where $\|K_{\alpha,\varphi}\| \approx A$;

2) $B = \left( \int\limits_a^b \left( \int\limits_t^b W^{q(\alpha-1)}(x) v(x) dx \right)^{\frac{q}{p-q}} W^{\frac{q(p-1)}{p-q}}(\varphi(t)) \frac{v(t) dt}{W^{q(1-\alpha)}(t)} \right)^{\frac{p-q}{pq}} < \infty$, when $0 < q < p < \infty$, $p > \frac{1}{\alpha}$, $0 < \alpha < 1$ and $\|K_{\alpha,\varphi}\| \approx B$, where $\|K_{\alpha,\varphi}\|$ is a norm of the operator $K_{\alpha,\varphi} : L_{p,w} \to L_{q,v}$.

**Keywords:** fractional integration operator, weighted estimate.

**2010 Mathematics Subject Classification:** 47A63, 47G10

**References:**

[1] Abylayeva A.M., and Oinarov R., "Criterion of the boundedness of a class of fractional integration operators", *Mathematical Journal.*, Almaty, Vol.4, No.2(12), pp. 5-14, 2004 (in Russian).

[2] Abylayeva A.M., "Criterion of the boundedness of $0 < \alpha < 1$ - order fractional integration type operator", Astana. No.6(46), pp. 130-136, 2005 (in Russian).

# Common best proximity points in complex valued metric spaces

Seyed Masoud AGHAYAN [1], Ahmad ZIREH [2]

[1] *Department of Mathematics, Payame Noor University, Tehran, Iran*
*E-mail: masoud_aghayan200@yahoo.com*
[2] *Department of Mathematics, Shahrood University of Technology, Shahrood, Iran*
*E-mail: azireh@shahroodut.ac.ir*

**Abstract:** In this paper, we introduce the concept of common best proximity points for non-self mappings between two subsets of a complex valued metric space which is an extension of the classical metric spaces obtained by allowing the metric function to assume values from the field of complex numbers and was recently introduced by Azam et al. [2]. Let A and B be non-empty subsets of a complex valued metric space and $S, T : A \to B$ be non-self mappings. Due to the fact that S and T are non-self mappings, the equations $Sx = x$ and $Tx = x$ are likely to have no common solution, known as a common fixed point of the mappings $S$ and $T$. Consequently, when there is no common solution, it is considered to determine an element x that is in close proximity to $Sx$ and $Tx$ in the sense of that $d(x, Sx)$ and $d(x, Tx)$ are minimum. In fact, common best proximity point theorems focus on the existence of such optimal approximate solutions, called common best proximity points, that the equations $Sx = x$ and $Tx = x$ in the case have no common solutions. As a matter of fact, given any element x in A, the distance between $x$ and $Sx$ and between $x$ and $Tx$ are at least $d(A, B)$, a common best proximity point theorem affirms global minimum of both complex valued functions $x \to d(x, Sx)$ and $x \to d(x, Tx)$ by imposing a common approximate solution of the equations $Sx = x$ and $Tx = x$ to satisfy the condition that $d(x, Sx) = d(x, Tx) = d(A, B)$. For this purpose, we extend some well-known results that was proved in classic metric space to the complex valued metric space by some new definitions. Our results supported by some examples.

**Keywords:** common best proximity point, complex valued metric space, common fixed points

**2010 Mathematics Subject Classification:** 54H25, 47H10

**References:**

[1] Amini-Harandi A., "Common best proximity points theorems in metric spaces $\mathbb{Z}_4$", *Optim. Lett.*, doi: 10.1007/s11590-012-0600-7, 2014.

[2] Azam A., Fisher B., Khan M., "Common fixed point theorems in complex valued metric spaces $\mathbb{Z}_4$", *Num. Func. Anal. and Optim.*, Vol.32, No.3, pp. 243-253, 2011.

[3] Ahmad J., Azam A., Saejung S., "Common fixed point results for contractive mappings in complex valued metric spaces $\mathbb{Z}_4$", *Fixed Point Theory and Appl.*, Vol. 67, 2014.

## On a difference scheme for nonlocal heat transfer boundary-value problem

Meiram AKHYMBEK [1], Makhmud SADYBEKOV [2]

[1] *Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan, E-mail: akhymbek@math.kz*

[2] *Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan E-mail: sadybekov@math.kz*

**Abstract:** In this paper, the family of boundary value problems for heat equation and finite difference schemes approximating these problems are considered. The peculiarity of the initial-boundary value problems is a special choice of the boundary conditions, which are not strongly regular. The corresponding difference schemes do not have the property of self-adjoint.

The paper proposes a new method of solving non-local problems for the heat equation

$$u_t(x,t) - u_{xx}(x,t) = f(x,t), \tag{1}$$

satisfying the initial condition

$$u(x,0) = \phi(x), \quad 0 \le x \le 1, \tag{2}$$

and the boundary conditions

$$\begin{cases} u_x(0,t) - u_x(1,t) + au(0,t) + bu(1,t) = 0, \quad cu(0,t) + (1+c)u(1,t) = 0. \\ \\ cu(0,t) + (1+c)u(1,t) = 0. \end{cases} \tag{3}$$

with finite difference method. The main important feature of these problems is their non-self-adjointness. This non-self-adjointness causes major difficulties in their analytical and numerical solving. The problems, which boundary conditions do not possess strong regularity, are less studied. The scope of study of the paper justifies possibility of building a stable difference scheme with weights for abovementioned type of problems.

**Keywords:** heat equation, initial boundary problems, numerical method, difference scheme.

**2010 Mathematics Subject Classification:** 35P05, 35E20, 39A14, 39A05

## Estimations of the best $M-$ term approximations of functions in the Lorentz space with constructive methods

Gabdolla AKISHEV

*Department of Mathematics and Information Technology,*
*Buketov Karaganda State University*
*E-mail: akishev@ksu.kz*

**Abstract:** V.N. Temlyakov [1,2] developed a constructive method of estimation of the best $n-$ term approximations functions from Nikol'skii-Besov's classes $S^{\bar{r}}_{p,\tau}B$ in the space $L_q(I^m)$ , $1 < p < q < \infty$. In talk considers $L^*_{\bar{q},\bar{\theta}}(I^m)$ – the Lorentz space of periodic functions of many variables with the anisotropic norm $\|f\|^*_{\bar{q},\bar{\theta}}$. $e_n(f)_{\bar{q},\bar{\theta}}$ - the best $n$-term approximation of $f$. $e_n(F)_{\bar{q},\bar{\theta}} = \sup_{f\in F} e_n(f)_{\bar{q},\bar{\theta}}$.

Suppose $\rho(\bar{s}) = \left\{\overline{k} = (k_1,\dots,k_m) \in \mathbb{Z}^m : \quad 2^{s_j-1} \le |k_j| < 2^{s_j}, j = 1,\dots,m\right\}$, $\delta_{\bar{s}}(f,\overline{x}) = \sum\limits_{\overline{n}\in\rho(\bar{s})} a_{\overline{n}}(f)e^{i\langle\overline{n},\overline{x}\rangle}$, where $\langle\bar{y},\bar{x}\rangle = \sum\limits_{j=1}^{m} y_j x_j$, $s_j = 1,2,\dots$. For a function $f \in L_1(I^m)$ suppose $f_l(\bar{x}) = \sum\limits_{\bar{s}:\langle\bar{s},\bar{\gamma}'\rangle=l} \delta_{\bar{s}}(f,\bar{x})$, $\bar{x} \in I^m$, $l \in \mathbb{N}\cap\{0\} = \mathbb{N}_0$, where $\bar{\gamma}' = (\gamma_1',\dots,\gamma_m')$, $\gamma_j' = (\frac{1}{q_j}-\frac{1}{p}))/(\frac{1}{q_1}-\frac{1}{p})$. We consider the class $W^{a,b}_{\bar{q},\theta} = \left\{f \in L_1(I^m) : \|f_l\|^*_{\bar{q},\theta} \le 2^{-la}\bar{l}^{(\nu-1)b}\right\}$, where $a > 0, b > 0$, $\bar{l} = \max\{1,l\}$ and let us introduce the notation $\|f\|_{W^{a,b}_{\bar{q},\theta}} = \sup\limits_{l\in\mathbb{N}_0} \|f_l\|^*_{\bar{q},\theta} 2^{la}\bar{l}^{-(\nu-1)b}$. In the case $q_1 = \dots = q_m = \theta$ this class considered V.N. Temlyakov [1]. The main results:

**Theorem**. Let $1 < q_m \le \dots \le q_{\nu+1} < q_\nu = \dots = q_1 < p < 2$, $1 < \theta < p$.

1. If $a > \frac{1}{q_1}+\frac{1}{\theta}-\frac{2}{p}$, then $e_n\left(W^{a,b}_{\bar{q},\theta}\right)_p \asymp n^{-a+\frac{1}{q_1}-\frac{1}{p}}(\log n)^{(\nu-1)(a+b+\frac{2}{p}-\frac{1}{q_1}-\frac{1}{\theta})}$.
2. If $\frac{1}{q_1}-\frac{1}{p} < a < \frac{1}{q_1}+\frac{1}{\theta}-\frac{2}{p}$, then $e_n\left(W^{a,b}_{\bar{q},\theta}\right)_p \asymp n^{-a+\frac{1}{q_1}-\frac{1}{p}}(\log n)^{(\nu-1)b}$.
3. If $a = \frac{1}{q_1}+\frac{1}{\theta}-\frac{2}{p}$, then $e_n\left(W^{a,b}_{\bar{q},\theta}\right)_p \asymp n^{-(\frac{1}{\theta}-\frac{1}{p})}(\log n)^{(\nu-1)b}(\log\log n)^{\frac{1}{\theta}}$.

**Keywords:** Lorentz space, $M$-term approximations, constructive method

**2010 Mathematics Subject Classification:** 41A10, 41A25

**References:**

[1] Temlyakov V.N., "Constructive sparse trigonometric approximation for functions with small mixed smoothness", *arXiv: 1503.00282v1 [NA]*, 30 p., 2015.
[2] Temlyakov V.N., "Constructive sparse trigonometric approximation and other problems for functions with mixed smoothness", *Sb. Math.*, Vol. 206, No 11, pp. 131-160, 2015.

## The maximum principle for the Navier-Stokes equations

Abdigali SH. AKYSH

*Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: akysh41@mail.ru*

**Abstract:** New connections were established between extreme values of the velocity, the density of kinetic energy and the pressure of the problem for the Navier-Stokes equations in domain $Q = (0, T] \times \Omega$:

$$\frac{\partial \mathbf{U}}{\partial t} - \mu \Delta \mathbf{U} + (\mathbf{U}, \nabla)\mathbf{U} + \nabla P = \mathbf{f},\ \mathrm{div}\mathbf{U} = 0, \tag{1}$$

$$\mathbf{U}(0, \mathbf{x}) = \mathbf{\Phi}(\mathbf{x}),\ \mathbf{x} \in \Omega \subset R_3;\ \mathbf{U}(t, \mathbf{x}) = 0,\ \mathbf{x} \in \partial\Omega. \tag{2}$$

Validity of the maximum principle was shown for problem (1, (2) using these connections:

$$\|\mathbf{U}\|_{\mathbf{C}(\bar{Q})} \leq \|\mathbf{\Phi}\|_{\mathbf{C}(\bar{\Omega})} + T\|\mathbf{f}\|_{\mathbf{C}(\bar{Q})} \equiv const,\ \forall T < \infty.$$

On the basis of which the selected space to prove the uniqueness of the weak and the existence of strong solutions to the problem Navier-Stokes equations for the whole time.

**Keywords:** Extreme values vector of velocity, density of kinetic energy, pressure, the maximum principle for the Navier-Stokes equations

**2010 Mathematics Subject Classification:** 35Q30, 35Q35

**References:**

[1] Ladyzhenskaya O.A.,"The 6th problem of millennium: the Navier-Stokes equation, existence and smoothness", *UMN*, 58:2(350), pp. 45-78, 2003.

[2] Akysh A.Sh.,"About the new version of maximum principle of Navier-Stokes equations",*Buketov KarSU Herald*, 2(78), pp. 11–17, 2015.

[3] Akysh A.Sh.,"On one version of principle of the maximum for the Navier-Stokes equations",*In the book: Analytical and Numerical Methods for Modeling of Natural Science and Social problems(ANM-2014). Russia. Penza*, pp. 3–9, 2014.

## The correctness Dirichlet problem in a cylindrical domain for three-dimensional elliptic equations with type and order extinction

Serik A. ALDASHEV, Erlan T. KITAYBEKOV

*Kazakh National Pedagogical University after Abay, Kazakhstan,*
*E-mail: aldash51@mail.ru*

**Abstract:** Correct formulation of boundary value problems in the plane for elliptic equations by the method of the theory of analytic complex variable functions is well studied. In the study of similar issues, when the number of independent variables is greater than two, there are difficulties of a fundamental nature. Very attractive and convenient method of singular integral equations loses its power because of absence of any comprehensive theory of multidimensional singular integral equations. Previously correctness of the Dirichlet problems for degenerate three-dimensional elliptic equations has been established. Importance of study of boundary value problems for the elliptic equations with degeneration of order and type was noticed by A.V. Bitsadze. Let $D_\beta-$ be a cylindrical domain of the Euclidian space $E_3$ of points $(x_1, x_2, t)$, bounded by the cylinder $\Gamma = \{(x,t) : |x| = 1\}$, the planes $t = \beta > 0$ and $t = 0$, where $|x| -$ is the length of the vector $x = (x_1, x_2)$. Let us denote the parts of these surfaces forming the boundary $\partial D_\beta$ of the domain $D_\beta$ as $\Gamma_\beta, S_\beta, S_o$ respectively. Let us consider three-dimensional hyperbolic equations with type and order confluence in the domain $D_\beta$ :

$$(1) \qquad \sum_{i=1}^{2} k_i(t)u_{x_ix_i} - k_3(t)u_{tt} + \sum_{i=1}^{2} a_i(x,t)u_{x_i} + b(x,t)u_t + c(x,t)u = 0,$$

where $k_i(t) > 0$ when $t > 0$ and they vanish when $t = 0$, $k_i(t) \in C([0,\beta]) \cap C^2((0,\beta)), i = 1,2,3$. Equation (1) is elliptic if and along the plane there is a degeneration of its type and order. As a multi-dimensional Dirichlet problem, we consider

**Problem 1.** Find the solutions of the equation (1) in the domain $D_\beta$ from the class $C(\bar{D}_\beta) \cap C^2(D_\beta)$, which satisfy the boundary value conditions

$$u|_{S_\beta} = \varphi(x), u|_{\Gamma_\beta} = \psi(x,t), u|_{S_o} = \tau(x).$$

For smooth coefficients of the equation (1) and certain conditions on the boundary data unique solvability is proved, and an explicit form of classical solution of the problem D is obtained.

**Keywords:** correctness, problem, equation, type and order extinction

**2010 Mathematics Subject Classification:** 35R12

## Relation theoretic metrical fixed point theorems with an application

Javid ALI

*Department of Mathematics, Aligarh Muslim University Aligarh 202002, India*
*E-mail: javid@amu.ac.in*

**Abstract:** In this paper, we establish metrical relation-theoretic fixed point theorems via an implicit contractive condition which is general enough to yield a multitude of corollaries corresponding to several well known contraction conditions (e.g. Banach (Fund. Math. 3, 133-181 (1922)), Kannan (Am. Math. Mon. 76, 405-408 (1969)), Reich (Can. Math. Bull. 14, 121-124 (1971)), Bianchini (Boll. Unione Mat. Ital. 5, 103-108 (1972)), Chatterjea (C. R. Acad. Bulg. Sci. 25, 727-730 (1972)), Hardy and Rogers (Can. Math. Bull. 16, 201-206 (1973)), Ciric (Proc. Am. Math. Soc. 45, 267-273 (1974)) and several others) wherein even such corollaries are new results on their own. As simple we utilize our main results, to prove a theorem on the existence and uniqueness of the solution of an integral equation besides furnishing an illustrative example.

**Keywords:** Complete metric space, binary relations, implicit relations, fixed point.

**References:**

[1] Alam A., Imdad M., "Relation-theoretic contraction principle", J. Fixed Point Theory Appl., Vol.17, pp. 693-702, 2015.

[2] Ali J., Imdad M., "An implicit function implies several contraction conditions", Sarajevo J. Math., Vol.4(17), pp. 269-285, 2008.

[3] Berinde V., Vetro F., "Common fixed points of mappings satisfying implicit contractive conditions", Fixed Point Theory Appl., Vol.105, 2012.

[4] Nieto J.J., Lopez R.R., "Contractive mapping theorems in partially ordered sets and application to ordinary differential equations", Order, Vol.22(3), pp. 223-239, 2008.

## The Steklov problem and estimates for orthogonal polynomials

Alexander I. APTEKAREV [1],
*Keldysh Institute of Applied Mathematics RAS*
*E-mail: aallaberen@gmail.com, fstetik@gmail.com*

**Abstract:** A Steklov's problem is to obtain the bounds on the sequence of orthonormal polynomials at the support of the weight of orthogonality. In 1921 [1], V.A. Steklov made a conjecture that if weight of orthogonality is strictly positive then sequence of orthonormal polynomials (at the support of the weight) is bounded.

In 1979 [2] E.A. Rakhmanov disproved this conjecture by constructing the sequence of polynomials orthonormal with respect to a positive weight which has the logarithmical rate of growth. Then the Steklov's problem becomes: to obtain the maximal possible rate of growth for these sequences. The modern version of the Steklov's problem is intimately related with the following extremal problem. For a fixed $n \in \mathbb{N}$, find

$$M_{n,\delta} = \sup_{\sigma \in S_\delta} \|\phi_n\|_{L^\infty(\mathbb{T})},$$

where $\phi_n(z)$ is the orthonormal polynomials with respect to the measure $\sigma \in S_\delta$, and $S_\delta$ is the Steklov's class of probability measures $\sigma$ on the unit circle, such that $\sigma'(\theta) \geqslant \delta/(2\pi) > 0$ at every Lebesgue point of $\sigma$.

There is an elementary bound

$$M_n \lesssim \sqrt{n}.$$

In 1981 [3] E.A. Rakhmanov have proved:

$$M_n \gtrsim \sqrt{n}/(\ln n)^{\frac{3}{2}}.$$

In our joint paper with S.A. Denisov and D.N. Tulyakov [4] we have proved, that

$$M_n \gtrsim \sqrt{n},$$

i.e. the elementary bound is sharp.

In our talk we discuss the history, statement of the problem and details of the construction of a sequence of the orthonormal polynomials from the Steklov class which have the maximal possible rate of growth. We also consider the Steklov problem in $L_p$ spaces and give new upper estimates for the polynomials orthonormal with respect Muckenhoupt weights from the Steklov's class.

**References:**

[1] W. A. Stekloff, Une méthode de la solution du problême de développement des fonctions arbitraires en séries de polynômes de Tchébysheff indépendante de la théorie de fermeture. II. Izv. Rus. Ac. Sci., 1921, 281–302, 303–326.

[2] E. A. Rahmanov, On Steklov's conjecture in the theory of orthogonal polynomials, Matem. Sb., 1979, 108(150), 581–608; English translation in: Math. USSR, Sb., 1980, 36, 549–575.

[3] E. A. Rahmanov, Estimates of the growth of orthogonal polynomials whose weight is bounded away from zero, Matem. Sb., 1981, 114(156):2, 269–298; English translation in: Math. USSR, Sb., 1982, 42, 237–263.

[4] A. Aptekarev, S. Denisov, D. Tulyakov, On a problem by Steklov, Journal of AMS, 2016, 29(4), 1117–1165.

## The positivity of the difference operator with periodic conditions and its applications

Allaberen ASHYRALYEV [1], Fatih Sabahattin TETİKOGLU [2]
*Emeritus Professor, Turkey & Turkmenistan*
*E-mail: aallaberen@gmail.com, fstetik@gmail.com*

**Abstract:** In this study, the second order of approximation of the difference operator $A_h^x$ approximates the second order differential dependent operator $A^x$ defined by the formula

$$A^x u = -a(x)u_{xx}(x) + \delta u(x), \delta \geq 0, a(x) = a(x + 2\pi), x \in \mathbb{R}^1$$

with domain

$$D(A^x) = \left\{ u(x) : u(x), u'(x), u''(x), u(x) = u(x + 2\pi), x \in \mathbb{R}^1, \int_0^{2\pi} u(x)dx = 0 \right\}$$

is presented. The Green's function of the difference operator $A_h^x$ is constructed. The estimates for the Green's function are obtained. The positivity of operator $A_h^x$ in the Banach space $C(\mathbb{R}_{1h})$ is established. It is proved that for any $\alpha \in (0, \frac{1}{2})$, the norms in spaces $E_\alpha = E_\alpha(C(\mathbb{R}_{1h}), A_h^x)$ and $C^{2\alpha}(\mathbb{R}_{1h})$ are equivalent uniformly with respect to $h$. The positivity of the operator $A_h^x$ in Hölder spaces of $\mathring{C}^{2\alpha}(\mathbb{R}_{1h})$, $\alpha \in (0, \frac{1}{2})$ is proved. In applications, theorems on well-posedness for difference schemes of the approximate solution of local and nonlocal boundary value problems for elliptic differential equations are presented.

**Keywords:** positivity of difference operators, periodic boundary conditions, Green's function.

**2010 Mathematics Subject Classification:** 34K13, 34B09,

**References:**

[1] Sobolevskii P.E., "A New Method of Summation of Fourier Series Converging in C-Norm", *Semigroup Forum*, Vol.71, pp. 289-300, 2005.

[2] Ashyralyev A. , Tetikoglu F. S., "The positivity of the second order difference operator with periodic conditions in Hölder spaces and its applications" *International Conference on Analysis and Applied Mathematics (ICAAM 2014), AIP Conf. Proc,* doi: 10.1063/1.4893857, Vol.1611, pp. 336-343, 2014.

[3] Ashyralyev A. , Tetikoglu F. S., "A note on fractional spaces generated by the positive operator with periodic conditions and applications" *Boundary Value Problems,* Vol.2015(31), doi:10.1186/s13661-015-0293-9, pp. 1-17, 2015.

# On the unique solvability of a nonlocal problem with integral conditions for the system of partial differential equations of second order

Anar ASSANOVA

*Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: anarasanova@list.ru; assanova@math.kz*

**Abstract:** The nonlocal boundary value problem with integral conditions for the system of hyperbolic equations second order is considered. Algorithm of finding approximate solution to researching problem is constructed and the its convergence is proved. The sufficient conditions of the unique solvability to nonlocal problem are established in the terms of initial data.

In this work, on $\Omega = [0,T] \times [0,\omega]$ the nonlocal boundary value problem with integral conditions for the system of hyperbolic equations is considered

$$\frac{\partial^2 u}{\partial x \partial t} = A(t,x)\frac{\partial u}{\partial x} + B(t,x)\frac{\partial u}{\partial t} + C(t,x)u + f(t,x), \tag{1}$$

$$P(t)u(t,\theta) + \int_0^\theta K(t,x)u(t,x)dx = \psi(t), \qquad t \in [0,T], \tag{2}$$

$$S(x)u(\eta,x) + \int_0^\eta M(t,x)u(t,x)dt = \varphi(x), \qquad x \in [0,\omega], \tag{3}$$

where $u(t,x) = col(u_1(t,x),...,u_n(t,x))$ is unknown function, the $(n \times n)$ matrices $A(t,x)$, $B(t,x)$, $C(t,x)$, $n$-vector function $f(t,x)$ are continuous on $\Omega$, the $(n \times n)$ matrices $K(t,x)$, $M(t,x)$ are continuously differentiable on $\Omega$ by $t$, $x$, respectively, the $(n \times n)$ matrices $P(t)$ and $S(x)$, the $n$-vector functions $\psi(t)$ and $\varphi(x)$ are continuously differentiable on $[0,T]$, $[0,\omega]$, respectively, and $0 \le \eta \le T$, $0 \le \theta \le \omega$.

The conditions of classical solvability to problem (1)–(3) are established by method of introduction additional functional parameters [1-2].

**Keywords:** hyperbolic equation, nonlocal problem, integral condition

**2010 Mathematics Subject Classification:** 35L51, 35L53

**References:**

[1] Asanova A.T., Dzhumabaev D.S., "Unique solvability of the boundary value problem for systems of hyperbolic equations with data on the characteristics ", *Comp. Math. and Math. Phys.*, Vol.42, No.11, pp. 1609-1621, 2002.

[2] Asanova A.T., Dzhumabaev D.S., "Well-posedness of nonlocal boundary value problems with integral condition for the system of hyperbolic equations", *J. of Math. Anal. and Appl.*, Vol.402, No.1, pp. 167-178, 2013.

## Asymptotical estimation of boundary value problems for singularly perturbed integro-differential equations

Nilupar ATAKHAN

*Department of Differential equations and Control theory, Al-Farabi Kazakh National University, Almaty, Kazakhstan*
*E-mail: atakhan-nilupar@mail.ru*

**Abstract:** We consider the two-point boundary value problem for the singularly perturbed higher order linear integro-differential equation. The constructive formulae and asymptotic estimations for solutions and their derivatives are obtained.

**Keywords:** singularly perturbation, integro-differential equations, small parameter, asymptotic method, asymptotic expansion, initial jump

**References:**

[1] L. Schlesinger *Uber asymptotische darstellungen der losungen linearer differential systeme als funktionen eines parameters*, Mathematische Annalen, **63(3)**, 277-300 (1907).

[2] G. D. Birkhoff *On the asymptotic character of the solutions of certain linear differential equations containing a parameter*, Transactions of the American Mathematical Society, **9(2)**,219-231 (1908).

[3] P. Noaillon *Developpements asymptotiques dans les equations differentielles lineaires a parametre variable*, Mem. Soc. Sci. Liege, **3(11)**, p.197 (1912).

[4] W. Wasow *Singular perturbations of boundary value problems for nonlinear differential equations of the second order*, Comm. On Pure and Appl. Math., 9, 93-113 (1956).

[5] A. N. Tihonov *O sistemah differencial'nyh uravnenij, soderzhashhih parametry*, Matematicheskij sbornik, **27(69)**, 147-156 (1950), (in Russian).

[6] M. I. Vishik, L. A. Lyusternik *Regular degeneration and boundary layer for linear differential equations with small parameter multiplying the highest derivatives*, Usp.Mat. Nauk, **12**, 3-122 (1957), (In Russian), Amer. Math. Soc. Transl., **20(2)**, 239-364 (1962).

[7] M. I. Vishik, L. A. Lyusternik *On the initial jump for non-linear differential equations containing a small parameter*, Doklady Akademii Nauk SSSR, **132(6)**, 1242-1245 (1960), (in Russian).

[8] K. A. Kasymov *Ob asimptotike reshenija zadachi Koshi s bol'shimi nachal'nymi uslovijami dlja nelinejnogo obyknovennogo differencial'nogo uravnenija, soderzhashhego malyj parametr*, Uspehi matematicheskih nauk, **17(5)**, 187-188 (1962).

[9] K. A. Kasymov *O zadache s nachal'nym skachkom dlja nelinejnyh sistem differencial'nyh uravnenij, soderzhashhih malyj parametr*, DAN SSSR., **179(2)**, 275-278 (1968), (in Russian).

[10] M. K. Dauylbayev, N. Atakhan *The initial jumps of solutions and integral terms in singular BVP of linear higher order integro-differential equations*, Miskolc Mathematical Notes, **16(2)**, 747-761 (2015).

## Boundary value problems for loaded hyperbolic equations

Anatoly ATTAEV

*Institute of Applied Mathematics and Automation, Nalchik, Russia*
*E-mail: attaev.anatoly@yandex.ru*

**Abstract:** The paper discusses some of the distinctive features, related to the formulation and study of boundary value problems for the loaded hyperbolic equations

$$u_{xx} - u_{yy} = \lambda u(x_0, y) \tag{1}$$

where $\lambda$ and $x_0$ are given real constants. Along with the setting of the Cauchy problem we investigate domain of dependency, influence and definition of solution $u(x, y)$ for equation (1) by the Cauchy datas in the whole space, as well as in its bounded subdomain. The Goursat problem and the Darboux problem in nonlocal setting are formulated and investigated. Some of these results has already been announced in works [1] and [2] in the case of the spectrally loaded equation [3] of the form

$$u_{xx} - u_{yy} = \lambda u_{yy}(x_0, y).$$

**Keywords:** loaded wave equation, Goursat problem, Darboux problem

**2010 Mathematics Subject Classification:** 35L05, 35L20

**References:**

[1] Attaev, A.Kh., "The boundary control problem for significantly loaded string oscillation equation", *Reports of the Adyghe (Circassian) International Academy of Sciences*, Vol. 13, No. 1, pp.30–35, 2011.

[2] Attaev, A.Kh., "An analog of the Darboux problem for essentially loaded vibration string equation", *Abstracts of the International Conference on Actual Problems of Mathematics and Informatics. Azerbaijan*, pp. 19–20, 2013.

[3] Dzhenaliev, M.T.; Ramazanov, M.I., "On a boundary value problem for a spectrally loaded heat operator", *Differ. Equ.*, Vol. 43, No. 4, pp. 513–524, 2007.

## On the boundedness of the periodic Hilbert transform on generalized periodic Morrey spaces

Sh.BAITUYAKOV [1], Zh. BAITUYAKOVA [2], M. ILYASOVA [3]

[1] *Department of Mechanics and mathematics, L.N. Gumilyov Eurasian National University, Astana, Kazakhstan*
*E-mail: baituyakova.zhzh@yandex.ru*

[2] *Kazakh University of Technology and Business, Astana, Kazakhstan*
*E-mail: baituyakov.sh@yandex.ru*

[3] *Department of Mechanics and mathematics, L.N. Gumilyov Eurasian National University, Astana, Kazakhstan*
*E-mail: meruert.ilyasova@yandex.ru*

**Abstract:** The Hilbert transform is one of the most important operators in the field of signal theory. Mainly, the importance of the transform is due to its property to extend real functions into analytic functions. This property certainly induces a vast number of applications, especially in signal theory, and obviously the Hilbert transform is not merely of interest for mathematicians. This paper dedicated to the investigation of boundedness of periodic Hilbert transform on generalized periodic Morrey spaces. First, we give definitions of Morrey space and generalized Morrey space [1], [2], [3]. Then we prove the boundedness of periodic Hilbert transform on generalized periodic Morrey spaces. The boundedness of Hilbert transform on (nonperiodic)generalized Morrey space we refer to [4]. As a consequence, given uniform estimates of the operator norms of the partial sums of Fourier series on generalized periodic Morrey spaces.

**Keywords:** Hilbert transform, partial sums of Fourier series

**2010 Mathematics Subject Classification:** 44A15, 42A24

**References:**

[1] M. Sautbekova, W. Sickel, "Strong summability of Fourier series and Morrey spaces, *Analysis Math.*, 40, pp. 31-62, 2014.

[2] T. Mizuhara,"Boundedness of some classical operators on generalized Morrey spaces, *Harmonic Anal.*, 4, pp.183-189, 1991.

[3] E.Nakai, "Hardy-Littlewood maximal operator, singular integral operators and the Riesz potential on generalized Morrey spaces, *Math. Nachr.*, 166, pp. 95-103, 1994.

[4] D.Hakim, E. Nakai, Y. Sawano, "Generalized fractional maximal operators and vector-valued inequalities on generalized Orlicz-Morrey spaces, *Rev.Mat.Complut*, DOI 10.1007/s13163-015-0178-6, 2015.

## Recovery of partial differential operators on classes of periodic functions with mixed smoothness

Sholpan A. BALGIMBAYEVA [1]

[1] *Department of Function Theory, Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan*
*E-mail: balgimbayeva@math.kz*

**Abstract:** We consider the problem of optimal linear recovery for mixed partial differential operator $\mathcal{A}$ on the unit ball $\mathrm{SB}^r_{p\theta}(\mathbb{T}^n)$ of the Nikol'skii–Besov space of periodic functions with mixed smoothness.

We find error bounds sharp in order for optimal linear recovery of operator $\mathcal{A}$ on class $\mathrm{SB}^r_{p\theta}(\mathbb{T}^n)$. As information $I^{\delta}_m(f)$ about the functions $f$ from class $\mathrm{SB}^r_{p\theta}(\mathbb{T}^n)$ we shall use Fourier coefficients with numbers from step "hyperbolic" cross.

As the linear method using the information about Fourier coefficients, we shall consider action of the mixed partial differential operator $\mathcal{A}$ on special "private" sum of decomposition on system (type as wavelets) trigonometrical polynoms.

**Keywords:** Nikol'skii–Besov function space, mixed smoothness, trigonometric system, recovery of differential operator

**2010 Mathematics Subject Classification:** 42A10

## Domain of the composition of some triangles in the space of $p$-summable sequences, $(0 < p \leq 1)$

Feyzi BAŞAR

*Prof.Dr., Turkey*
*E-mail: feyzibasar@gmail.com*

**Abstract:** Let $0 < p \leq 1$. In the present paper, referring Ng and Lee [1] and following Altay and Başar [2, 3] and Başar [4] we introduce the space $\widetilde{\ell}_p$ as the domain of the triangle $\widetilde{B}$ which is the composition of $E_1$, $C_1$ and $\Delta$ of the Euler, Cesàro means of order one and the backward difference matrix. Furthermore, we give an inclusion relation concerning the space $\widetilde{\ell}_p$ and compute the alpha-, beta- and gamma-duals of the space $\widetilde{\ell}_p$ and construct its basis. We characterize the classes $(\widetilde{\ell}_p : \ell_\infty)$, $(\widetilde{\ell}_p : c)$ and $(\widetilde{\ell}_p : c_0)$ of infinite matrices, and give the characterization of some other classes of matrix transformations from the space $\widetilde{\ell}_p$ to the Euler, Riesz, difference, etc., sequence spaces, by means of a basic lemma.

**Keywords:** domain of a triangle matrix, sequence space, $BK$-space, the $\alpha$-, $\beta$- and $\gamma$-duals, Schauder basis, matrix transformations.

**2010 Mathematics Subject Classification:** 46A45, 46B45, 46A35

**References:**

[1] Ng P. -N., Lee P. -Y., "Cesàro sequences spaces of non-absolute type", *Comment. Math. Prace Mat.* Vol. 20, No.2, pp. 429–433, 1978.

[2] Altay B., Başar, F., "Some Euler sequence spaces of non-absolute type", *Ukrainian Math. J.* Vol. 57, No. 1, pp. 1–17, 2005.

[3] Altay B., Başar, F., "The matrix domain and the fine spectrum of the difference operator $\Delta$ on the sequence space $\ell_p$, $(0 < p < 1)$", *Commun. Math. Anal.* Vol. 2, No. 2, pp. 1–11, 2007.

[4] Başar F., "Domain of the composition of some triangles in the space of $p$-summable sequences", *AIP Conference Proceedings*, 1611, pp. 348–356, 2014.

## On some spaces of $q-$bounded double sequences

Feyzi BAŞAR [1], Hüsamettin ÇAPAN [2]

[1] *Prof. Dr., Turkey*

*E-mail: feyzibasar@gmail.com, husamettincapan@gmail.com*

**Abstract:** Let $0 < q < \infty$ and $\vartheta \in \{p, bp, r\}$. By $\mathcal{M}_u$, $\mathcal{C}_\vartheta$, $\mathcal{C}_{\vartheta 0}$ and $\mathcal{L}_q$; we denote the spaces of all bounded, $\vartheta-$convergent, $\vartheta-$null and absolutely $q-$summable double sequences, respectively (see e.g. [1–3]). In [4], Altay and Başar have defined the spaces $\mathcal{BS}$, $\mathcal{CS}_\vartheta$ and $\mathcal{CS}_{\vartheta 0}$ as the domain of four-dimensional summation matrix $S$ in the spaces $\mathcal{M}_u$, $\mathcal{C}_\vartheta$ and $\mathcal{C}_{\vartheta 0}$; respectively. Besides these, by using four-dimensional backward difference matrix $\Delta$ they have introduced the space $\mathcal{BV}$ of all bounded variation double sequences. As a continuation of [4]; in this study, we investigate the spaces $\mathcal{LS}_q$ and $\mathcal{BV}_q$ consisting of all double sequences whose $S$- and $\Delta$-transforms are in the space $\mathcal{L}_q$, respectively.

**Keywords:** summability theory, double sequence spaces, dual spaces, matrix transformations.

**2010 Mathematics Subject Classification:** 46A45, 40C05

**References:**

[1] Pringsheim, A., "Zur Theorie der zweifach unendlichen Zahlenfolgen", *Math. Z.* Vol. 53, pp. 289–321, 1900.

[2] Hardy, G.H., "On the convergence of certain multiple series", *Proc. Cambridge Philos. Soc.* Vol. 19, pp. 86–95, 1916–1919.

[3] Móricz, F., "Extensions of the spaces $c$ and $c_0$ from single to double sequences", *Acta Math. Hungar.* Vol. 57, pp. 129–136, 1991.

[4] Altay, B., Başar, F., "Some new spaces of double sequences", *J. Math. Anal. Appl.* Vol. 309, pp. 70–90, 2005.

---

## Fourier widths of some function classes associated with $m-$multiple Haar system

Dauren BAZARKHANOV

*Department of Function Theory, Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan*

*E-mail: dauren.mirza@gmail.com*

**Abstract:** In the communication we obtain estimates sharp in order for Fourier widths of the Nikol'skii – Besov and Lizorkin – Triebel type classes associated with $m-$multiple Haar system in the space $L_r([0,1]^k)$ for a number of relations between the parameters of classes and space.

**Keywords:** $m-$multiple Haar system, Fourier width, the Nikol'skii – Besov type space, the Lizorkin – Triebel type space

**2010 Mathematics Subject Classification:** 42A10

## Noncommutative Holder-type inequalities

Turdebek N. BEKJAN[1]

[1] Department of Fundamental Mathematics, L. N. Gumilyov Eurasian National University, Àstana, Kazakhstan
E-mail: bekjant@yahoo.com

**Abstract:** We generalize some known Holder-type inequalities of matrix to the measurable operators associated with semi-finite von Neumann algebra $\mathcal{M}$ case.

In [4], Horn and Zhan have shown that if $p,\ q$ and $r$ are positive real numbers such that $\frac{1}{p}+\frac{1}{q}=1$, then

$$|||A^*B|^r||| \leq |||A|^{pr}|||^{\frac{1}{p}}\,|||B|^{qr}|||^{\frac{1}{q}} \tag{1}$$

for $A, B \in B(H)$ and for unitarily invariant norms. Hiai and Zhan [2] proved that if $A,\ B,\ C$ and $D$ in $B(H)$, then

$$2^{|\frac{1}{p}-\frac{1}{2}|}|||\,|C^*B+D^*B|\,||| \leq |||\,|A|^p+|B|^p|||^{\frac{1}{p}}\,|||\,|C|^q+|D|^q|||^{\frac{1}{q}} \tag{2}$$

for all positive real numbers $p$ and $q$ such that $\frac{1}{p}+\frac{1}{q}=1$.

In this talk, we will use the concept of uniform Hardy-Littlewood majorization studied by Kalton and Sukochev [5], and their main result, to generalize the Holder type inequalities (1) and (2) for $\tau$-measurable operators associated with semi-finite von Neumann algebra $\mathcal{M}$ and for symmetric Banach spaces norm.

**Keywords:** first, quadratic residue codes

**2010 Mathematics Subject Classification:** 94B05, 94B15

**References:**

[1] Bekjan T. N., and Ospanov M. N., "Holder-type inequalities of measurable operators", to appear Positivity.

[2] Hiai F., and Zhan X., "Inequalities involving unitarily invariant norms and operator monotone functions", *Lin. Algebra Appl.*, Vol. 341, pp. 151-169, 2002.

[3] Horn R.A., and Mathias R., "Cauchy-Schwarz inequality associated with positive semi definite matrices", *Lin. Algebra Appl.*, Vol. 142, pp. 63-82, 1990.

[4] Horn R.A., and Zhan X., "Inequalities for C-S semi norms and Lieb functions", *Lin. Algebra Appl.*, Vol. 291, pp. 103-113, 1999.

[5] Kalton N. J., and Sukochev F. A., "Symmetric norms and spaces of operators", *J. Reine Angew. Math.*, Vol. 621, pp. 81-121, 2008.

## Interpolation properties of anisotropic Nikol'skii-Besov $B^{\alpha\mathbf{q}}_{\mathbf{pr}}(\mathbb{T}^{\mathbf{d}})$ spaces and embedding theorems

K.A. BEKMAGANBETOV [1], YE. TOLEUGAZY [2]

[1] *M.V. Lomonosov MSU, Kazakhstan Branch*
*E-mail: bekmaganbetov-ka@yandex.ru*
[2] *L.N. Gumilyov Eurasian National University*
*erzhan1996@yandex.ru*

**Abstract:** We study interpolation properties of anisotropic of Nikol'skii–Besov spaces $B^{\alpha\mathbf{q}}_{\mathbf{pr}}(\mathbb{T}^{\mathbf{d}})$ and we obtain limit embedding theorems for anisotropic Nikol'skii-Besov and Lorentz's spaces.

In article we get interpolations theorems for anisotropic Nikol'skii–Besov $B^{\alpha\mathbf{q}}_{\mathbf{pr}}(\mathbb{T}^{\mathbf{d}})$ spaces. Apply its we proofed the embedding of the different metrics, limits theorems for anisotropic Nicol'skii–Besov spaces and the anisotropic Lorentz spaces.

**Theorem 1.** Let $\mathbf{1} < \mathbf{p} = (p_1, \dots, p_n) < \mathbf{q} = (q_1, \dots, q_n) < \infty$, $\mathbf{1} \le \mathbf{r} = (r_1, \dots, r_n), \tau = (\tau_1, \dots, \tau_n) \le \infty$. Then for $\alpha = (\mathbf{1/p - 1/q})\mathbf{d}$ the embedding holds

$$B^{\alpha\tau}_{\mathbf{pr}}(\mathbb{T}^{\mathbf{d}}) \hookrightarrow L_{\mathbf{q}\tau}(\mathbb{T}^{\mathbf{d}}).$$

**Theorem 2.** Let $\mathbf{1} < \mathbf{q} = (q_1, \dots, q_n) < \mathbf{p} = (p_1, \dots, p_n) < \infty$, $\mathbf{1} \le \mathbf{r} = (r_1, \dots, r_n), \tau = (\tau_1, \dots, \tau_n) \le \infty$. Then for $\alpha = (\mathbf{1/p - 1/q})\mathbf{d}$ takes place embedding

$$L_{\mathbf{q}\tau}(\mathbb{T}^{\mathbf{d}}) \hookrightarrow B^{\alpha\tau}_{\mathbf{pr}}(\mathbb{T}^{\mathbf{d}}).$$

The research was supported by the Committee of Science of the Ministry of Education and Science of Republic of Kazakhstan (Grant 0816/GF4).

**References:**

[1] Nursultanov E.D., On the coefficients of multiple Fourier series from $L_p$-spaces. (Russian) // Izvestiya: Mathematics, Vol.64: 1, pp. 93-120, 2000.

[2] Bergh J., and Löfström J., Interpolation spaces. An introduction. Grundlehren der Mathematischen Wissenschaften, vol. 223, Springer-Verlag, Berlin – Hiedelburg – New York 1976.

[3] Bazarkhanov D.B., Various representations and equivalent normalizations of Nicol'skii-Besov and Lizorkin-Tribel spaces of the generalized mixed smoothness. Dokl. Akad. nauk. Vol.402, No.3, pp. 298-302.

[4] Bekmaganbetov K.A., Order of Approximation of Besov Classes in the Metric of Anisotropic Lorentz Spaces // Methods of Fourier Analysis and Approximation theory. Series: Applied and Numerical Harmonical Analysis, Birkhäuser, pp. 149-159, 2016.

[5] Tribel H., Interpolation theory, function spaces, differential operators, North-Holland Math. Labrary, Vol.18, North-Holland, Amsterdam – New York – Oxford, 1978.

[6] Bekmaganbetov K.A., and Nursultanov E.D., Embedding theorems for anisotropic Besov spaces $B^{\alpha\mathbf{q}}_{\mathbf{pr}}([0, 2\pi)^n)$ // Izvestiya: Mathematics, Vol.73(4), 655, pp. 3-16, 2009.

[7] Bekmaganbetov K.A., Nursultanov E.D., The method of multi-parameter interpolation and imbedding theorems of Besov spaces $B^{\vec{\alpha}}_{\vec{p}}([0, 2\pi))$ // Analysis Mathematica, Vol.24, Issue 4, pp. 262-263, 1998.

[8] Bekmaganbetov K.A., Nursultanov E.D., On interpolation and embedding theorems for the Spaces $B^{\sigma q}_{p\tau}(\Omega)$ // Mathematical Notes, Vol.84, Issue 5-6, pp. 733-736, 2008.

## Weighted inequality of Hardy type

Sh. BILALL [1], S. Kh. SHALGINBAYEVA [2]

[1] *Institute of Mathematics and mathematical modeling of MES RK, Almaty, Kazakhstan*
*E-mail: bilal44@mail.ru*
[2] *Kazakh Ablai Khan University of International Relations and World Languages, Almaty, Kazakhstan*
*E-mail:*

**Abstract:** This work is an extension of the results of article. The necessary and sufficient conditions on weight functions $\rho, \nu, r$ are obtained, for which the following inequality holds true

$$\|rf\|_q = c\left(\left\|\rho f'\right\|_p + \|\nu f\|_p\right) \tag{3}$$

for $p = q = \infty$. And for the smallest constant two-sided estimates are found, which are of the same order. The criterion of compactness of embedding operator, that corresponds to (1) is provided. Let us note that (1) becomes a generalized Hardy inequality (in differential form) as $\nu \equiv 0$. For $\rho \equiv 1, \quad 1 = p \le q < \infty$, the inequality (1) was examined by K.T. Mynbayev, for $\rho \equiv 1, \quad p = q = 2$ , by E.T. Sawyer, for $1 < p \le q < \infty$ by M. Otelbaev and R.Oynarov, and for $1 \le p < q < \infty, \quad 1 \le q < p < \infty$, in general situations by R. Oynarov. Similar inequalities for matrix operators are considered in the works of J. Tasmagambetova and A. Temirkhanova.

We introduce the following parameters describing the local behavior of the weighting functions:

$$\omega(x,y) = \sup\left\{d > 0 : \int_{x-d}^{x} \rho^{-1}(\tau)d\tau \le \int_{x}^{x+y} \rho^{-1}(\tau)d\tau, (x-d, x+y] \subset J\right\},$$

$$d^+(x) = \sup\left\{d > 0 : \sup_{x-\omega(x,d)\le t\le x+d} \upsilon(t) \int_{x-\omega(x,d)}^{x+d} \rho^{-1}(\tau)\, d\tau \le 1, \right.$$
$$\left. [x, x+d) \subset J \right\},$$

$$d^-(x) = \omega\left(x, d^+(x)\right).$$

**Keywords:** additive inequality, matrix operator, bounded sequence.

**2010 Mathematics Subject Classification:** 42A10

## On the compactness of set in generalized Morrey spaces

Nurzhan BOKAYEV [1], Victor BURENKOV [2], Dauren MATIN [1]

[1] *Department of Mechanics and Mathematics, L.N. Gumilyov Eurasian National University, Astana, Kazakhstan*
*E-mail: bokayev2011@yandex.ru, d.matin@mail.ru*
[2] *RUDN, Moscow, Russia*
*E-mail: burenkov@cardiff.ac.uk*

**Abstract:** In this paper, we present sufficient conditions for strongly pre-compact sets in the generalized Morrey spaces $M_p^w$.

Let $1 \le p \le \infty$, $w(x)$ measurable non-negative function on $(0,\infty)$. Generalized Morrey space $M_p^{w(R)}$ is defined as the set of functions $f \in L_1^{loc}(R)$ with a finite quasi-norm: $\|f\|_{M_p^w} \equiv \sup\limits_{t\in R^n, r>0} w(r) \left( \int\limits_{B(t,r)} |f(x)|^p \, dx \right)^{\frac{1}{p}} < \infty$, where $B(t,r)$ is a ball with center at the point $t$ and with radius $r$. For $f \in L_1^{loc}(R)$ and $\alpha > 0$ suppose $(M_\alpha f)(x) = \frac{1}{|B(x,\alpha)|} \int\limits_{B(x,\alpha)} f(y)dy$. We denote by $\Omega_p$ the set of all functions which are non-negative , measurable on $\Omega_p$ and for some $t_1, t_2 > 0$: $\sup\limits_{t_1 \le r < \infty} (w(r)) < \infty$, $\sup\limits_{0<r\le t_2} \left( w(r) r^{\frac{1}{p}} \right) < \infty$.

**Theorem** Suppose that $1 \le p \le \infty$ and $w \in \Omega_p$. Suppose the subset $S$ in $M_p^w$ satisfies the following conditions:

$$\sup_{f\in S} \|f\|_{M_p^w} < \infty, \ \lim_{u\to 0} \sup_{f\in S} \|f(\cdot+u) - f(\cdot)\|_{M_p^w} = 0, \ \lim_{r\to\infty} \sup_{f\in S} \left\| f\chi_{C\,B(0,r)} \right\|_{M_p^w} = 0.$$

Then $S$ is strongly pre-compact set in $M_p^w(R)$.

From the proved theorem for the case of $w(r) = r^{-\lambda}$ follows a well-known result for the Morrey space $M_p^\lambda$ [3] and in the case of $\lambda = 0$ this is well-known Frechet-Kolmogorov theorem.

**Keywords:** generalized Morrey spaces, compactness

**2010 Mathematics Subject Classification:** 42B20, 42B99

**References:**

[1] Chen Y., Ding Y., Wang X. Compactness of Commutatus for Singular Integrals on Morrey Spaces. *Canad.J.Math.* Vol.64(2), pp. 257-281, 2012.

## A variation on lacunary statistical quasi-Cauchy sequences

Huseyin CAKALLI [1], Mikail ET [2], Hacer SENGUL [3]

[1] *Department of Mathematics, Maltepe University, İstanbul, Turkey*
*E-mail: huseyincakalli@maltepe.edu.tr, hcakalli@gmail.com*
[2] *Department of Mathematics, Firat University, Elazig, Turkey*
*E-mail: met@firat.edu.tr, mikailet68@gmail.com*
[3] *Department of Mathematics, Siirt University, Siirt, Turkey*
*E-mail: hacersengul@siirt.edu.tr, hacer.sengul@hotmail.com*

**Abstract:** We introduce a concept of ideal lacunary statistical quasi-Cauchyness of order $\alpha$ of a sequence of real numbers in the sense that a sequence $(x_k)$ of points in $\mathbf{R}$ is called $I-$lacunary statistically quasi-Cauchy of order $\alpha$, if $\{r \in \mathbf{N} : \frac{1}{h_r^{\alpha}}|\{k \in I_r : |\Delta x_k| \geq \varepsilon\}| \geq \delta\} \in I$ for each $\varepsilon > 0$ and for each $\delta > 0$, where an ideal $I$ is a family of subsets of positive integers $\mathbf{N}$ which is closed under taking finite unions and subsets of its elements. The main purpose of this paper is to investigate ideal lacunary statistical ward continuity of order $\alpha$, where a function $f$ is called $I-$ lacunary statistically ward continuous of order $\alpha$ if it preserves $I$-lacunary statistically quasi-Cauchy sequences of order $\alpha$, i.e. $(f(x_n))$ is a $S_{\theta}^{\alpha}(I)$-quasi-Cauchy sequence whenever $(x_n)$ is.

**Keywords:** sequences, series, ideal convergence, continuity
**2010 Mathematics Subject Classification:** 40A05, 40A35, 26A15

**References:**

[1] Burton D., and Coleman J., "Quasi-Cauchy Sequences", *Amer. Math. Monthly*, Vol. 117, No.5, pp. 328-333, 2010.

[2] Çakalli H., "A variation on ward continuity", *Filomat*, Vol. 8, pp. 1545-1549, 2013.

[3] Çakalli H., and Hazarika B., *Ideal Quasi-Cauchy sequences*, *J. Inequal. Appl.*, pp. 2012:234, 2012. doi:10.1186/1029-242X-2012-234

[4] Cakalli H., "On G-continuity", *Comput. Math. Appl.*, Vol.61, pp. 313-318, 2011.

[5] Das P., Savas E., "On $I$-statistically pre-Cauchy sequences", *Taiwanese J. Math.*, Vol.18, no. 1, pp. 115-126, 2014.

[6] Das P., Savas E., Ghosal Sanjoy Kr., "On generalizations of certain summability methods using ideals", *Appl. Math. Lett.*, Vol.24, no. 9, pp. 1509-1514, 2011.

[7] Kostyrko P., Šalàt T., and Wilczyński W., "$I$-convergence", *Real Anal. Exchange* , Vol.26, No.2, pp. 669-686, 2000-2001.

[8] Sengul H., and Et M., "On lacunary statistical convergence of order alpha", *Acta Mathematica Scientia*, Vol.34, No.2, pp. 473-482, 2014.

## Ideal statistically quasi Cauchy sequences

Ekrem SAVAS [1], Huseyin CAKALLI [2]

[1] *Department of Mathematics, Istanbul Commerce University, İstanbul, Turkey*
*E-mail: ekremsavas@yahoo.com*
[2] *Department of Mathematics, Maltepe University, İstanbul, Turkey*
*E-mail: huseyincakalli@maltepe.edu.tr; hcakalli@gmail.com*

**Abstract:** An ideal $I$ is a family of subsets of positive integers $\mathbf{N}$ which is closed under taking finite unions and subsets of its elements. A sequence $(x_k)$ of real numbers is said to be $S(I)$-statistically convergent to a real number $L$, if for each $\varepsilon > 0$ and for each $\delta > 0$ the set $\{n \in \mathbf{N} : \frac{1}{n}|\{k \leq n : |x_k - L| \geq \varepsilon\}| \geq \delta\}$ belongs to $I$. We introduce $S(I)$-statistically ward compactness of a subset of $\mathbf{R}$, the set of real numbers, and $S(I)$-statistically ward continuity of a real function in the senses that a subset $E$ of $\mathbf{R}$ is $S(I)$-statistically ward compact if any sequence of points in $E$ has an $S(I)$-statistically quasi-Cauchy subsequence, and a real function is $S(I)$-statistically ward continuous if it preserves $S(I)$-statistically quasi-Cauchy sequences where a sequence $(x_k)$ is called to be $S(I)$-statistically quasi-Cauchy when $(\Delta x_k)$ is $S(I)$-statistically convergent to 0. We obtain results related to $S(I)$-statistically ward continuity, $S(I)$-statistically ward compactness, $N_\theta$-ward continuity, and slowly oscillating continuity.

**Keywords:** sequences, series, ideal convergence, continuity
**2010 Mathematics Subject Classification:** 40A05, 40A35, 26A15

**References:**

[1] Burton D., and Coleman J., "Quasi-Cauchy Sequences", *Amer. Math. Monthly*, Vol. 117, No.5, pp. 328-333, 2010.
[2] Çakalli H., "Statistical-quasi-Cauchy sequences", *Math. Comput. Modelling*, Vol. 54, No.5-6, pp. 1620-1624, 2011.
[3] Çakalli H., "A variation on ward continuity", *Filomat*, Vol. 8, pp. 1545-1549, 2013.
[4] Çakalli H., and Hazarika B., *Ideal Quasi-Cauchy sequences*, *J. Inequal. Appl.*, pp. 2012:234, 2012. doi:10.1186/1029-242X-2012-234
[5] Cakalli H., "On G-continuity", *Comput. Math. Appl.*, Vol.61, pp. 313-318, 2011.
[6] Das P., Savas E., "On $I$-statistically pre-Cauchy sequences", *Taiwanese J. Math.*, Vol.18, no. 1, pp. 115-126, 2014.
[7] Das P., Savas E., Ghosal Sanjoy Kr., "On generalizations of certain summability methods using ideals", *Appl. Math. Lett.*, Vol.24, no. 9, pp. 1509-1514, 2011.
[8] Fast H., "Sur la convergence statistique", Colloq. Math.", Vol.2, pp. 241-244, 1951.
[9] Kostyrko P., Šalàt T., and Wilczyński W., "$I$-convergence", *Real Anal. Exchange* , Vol.26, No.2, pp. 669-686, 2000-2001.

# Solvability of the Dirichlet problem for the $p$-Laplacian on stratified sets

Nurlan DAIRBEKOV [1], Oleg PENKIN [2], Lyazzat SARYBEKOVA [3]

*School of Mathematics and Cybernetics, Kazakh-British Technical University, Almaty, Kazakhstan*
*E-mail: nurlan.dairbekov@gmail.com* [1]
*E-mail: o.m.penkin@gmail.com* [2]
*E-mail: lsarybekova@yandex.ru* [3]

**Abstract:** A stratified set is defined to be a connected subset $\Omega$ of $\mathbb{R}^d$ which consists of a finite collection $S$ of submanifolds $\sigma_{kj}$, called strata. The first subscript in $\sigma_{kj}$ shows the dimension of a stratum, while the second serves to list strata of the same dimension. So

$$\Omega = \bigcup_{\sigma_{kj} \in S} \sigma_{kj}.$$

We assume each $\sigma_{kj}$ to have compact closure in $\mathbb{R}^d$ and assume also that strata adhere to one another regularly like cells in a CW-complex. The set $\Omega$ is regarded as the union $\Omega_0 \bigcup \partial\Omega_0$, where $\Omega_0$ is an open connected subset of $\Omega$ (the topology on $\Omega$ is induced by the standard topology of $\mathbb{R}^d$) composed of strata and such that the closure $\overline{\Omega}_0$ coincides with $\Omega$. The difference $\partial\Omega_0 = \Omega \setminus \Omega_0$ is the boundary of $\Omega_0$ in the induced topology.

A stratified measure $\mu$ on $\Omega$ is defined as

$$\mu(\omega) = \sum_{\sigma_{kj} \in S} \mu_k(\sigma_{kj} \cap \omega)$$

for naturally defined mesurable sets $\omega \subset \Omega$. Here $\mu_k$ is the measure on $\sigma_{kj}$ related to the Riemannian metric on $\sigma_{kj}$ induced by the Euclidean metric of $\mathbb{R}^d$.

For tangent vector fields $\vec{F}$ on $\Omega_0$, which can be defined quite naturally, we can consider an analog $\nabla \cdot \vec{F}$ of classical divergence, defined as the density of the flux of $\vec{F}$ with respect to $\mu$. For functions $u : \Omega \to \mathbb{R}$, we then can consider an analog of the Laplace–Beltrami operator, $\Delta u = \nabla \cdot \nabla u$, with $\nabla u$ being a specially-defined gradient.

In our talk we discuss the solvability of the Dirichlet problem for the $p$-Laplacian on a stratified set:

$$\nabla(|\nabla u|^{p-2}\nabla u) = 0, \tag{4}$$

$$u\big|_{\partial\Omega_0} = \phi. \tag{5}$$

Equation (4) is the Euler-Lagrange equation for the variational problem

$$\Phi(u) = \int_{\Omega_0} |\nabla u|^p d\mu \to \min,$$

where the minimum is taken over all continuous functions on $\Omega$ that are continuously differentiable on every stratum of $\Omega_0$ and whose boundary values are given by $\phi$.

We show that Problem (4),(5) has a weak solution in a suitable Sobolev space $W_\mu^{1,p}(\Omega_0)$ determined by the norm $\|u\|_{W_\mu^{1,p}(\Omega_0)} = \left( \int\limits_{\Omega_0} |\nabla u|^p \, d\mu \right)^{\frac{1}{p}}$.

The key point in the proof is played by an analog of the Sobolev inequality,

$$\int\limits_{\Omega} |f|^q \, d\mu \le C \int\limits_{\Omega_0} |\nabla f|^p \, d\mu,$$

where the parameter $q$ depends on $p$ and on the structure of $\Omega$. The latter dependence is rather complicated and, rouphly speaking, shows how tightly the strata are bound to one another.

**Keywords:** stratified set, $p$-Laplacian, Sobolev inequality

**2010 Mathematics Subject Classification:** 35J25, 35B45

**References:**

[1] Gavrilov A., Nicaise S., Penkin O.M. "Poincares inequality on the stratified sets and applications", *Progress in nonlinear differential equations and their applications*, Vol. 55, pp. 195–213, 2003 Birkhauser Verlag Basel/Switzerland

[2] Nicaise S., Penkin O.M. "Poincare Perrons method for the Dirichlet problem on stratified sets", *J. Math. Anal. Appl.*, Vol. 296, No.2, pp. 504–520, 2004.

# Asymptotic estimates of the solution of the Cauchy problem for singularly perturbed integro-differential equations

Muratkhan DAUYLBAYEV [1], Aziza MIRZAKULOVA [2]
*Al-Farabi Kazakh National University, Almaty, Kazakhstan*
*E-mail:* [1] *dmk57@mail.ru,* [2] *aziza.mirzakulova@mail.ru*

**Abstract:** This report is devoted to the Cauchy problem for the integro-differential equation with a small parameter in the highest derivatives:

$$\sum_{r=1}^{m} \varepsilon^r A_{n+r}(t) y^{(n+r)} + \sum_{k=0}^{n} A_k(t) y^{(k)} = F(t) + \int_0^1 \sum_{i=0}^{l+1} H_i(t,x) y^{(i)}(x,\varepsilon) dx \tag{6}$$

with the initial conditions

$$y^{(i)}(0,\varepsilon) = \alpha_{i+1}, \quad i = \overline{0, n+m-1}. \tag{7}$$

Here $\varepsilon > 0$ is small parameter and $l = \text{fix}\{0, 1, \ldots, n-1\}$.

A similar problem for differential equations considered in [1].

In this paper we consider the case when the roots $\mu_i(t)$, $i = \overline{1,m}$ of the additional characteristic equation

$$\mu^m + A_{n+m-1}(t)\mu^{m-1} + \ldots + A_{n+1}(t)\mu + A_n(t) = 0$$

have negative real parts.

Asymptotic estimates for the solution of the problem (1) and (2) and of the difference between the solutions of the original singularly perturbed problem (1) and (2) and a corresponding unperturbed problem are obtained.

**Keywords:** Singularly perturbations, small parameter, integro-differential equations, asymptotic estimates

**2010 Mathematics Subject Classification:** 34E15, 34K26, 45J05

**References:**

[1] Kassymov K.A., Nurgabyl D.N. "The asymptotic behavior of solutions of the linear singularly perturbed common undivided boundary value problems with the initial jump", *Ukrainian mathematical journal*, Vol. 55, No. 11, pp. 1496–1508, 2003.

## Weighted Sobolev spaces, quasielliptic operators and equations not solvable with respect to the highest-order derivative

Gennadii DEMIDENKO

*Sobolev Institute of Mathematics, Novosibirsk, Russia,*
*Novosibirsk State University, Novosibirsk, Russia*
*E-mail: demidenk@math.nsc.ru*

**Abstract:** In the paper some classes of quasielliptic operators $L(D_x)$ are considered in the whole space $\mathbb{R}^n$. In particular, they include homogeneous elliptic operators, elliptic and parabolic operators in the sense of Petrovskii, elliptic operators in the sense of Douglis – Nirenberg, operators of parabolic type with "opposite time directions" and etc. [1, 2]. We introduce special scales of weighted Sobolev spaces $W^l_{p,\sigma}$ and study their properties. We establish isomorphism theorems for the quasielliptic operators These results give us known isomorphism theorems for homogeneous elliptic operators and a number of new isomorphism theorems for matrix elliptic and parabolic operators; in particular, for the stationary Navier–Stokes operator.

Using the isomorphism theorems, we establish unique solvability of initial value problems for some classes of partial differential equations not solvable with respect to the highest-order derivative

$$L(D_x)D_t^m u + \sum_{k=0}^{m-1} L_{m-k}(D_x)D_t^k u = f(t,x).$$

In particular, we consider equations and systems of Sobolev type, pseudoparabolic and pseudohyperbolic equations [3, 4]. The author was supported by the Program of the Presidium of the Russian Academy of Sciences (project no. 0314-2015-0011).

**Keywords:** quasielliptic operators, weighted Sobolev spaces, isomorphism theorems, Sobolev type equations

**2010 Mathematics Subject Classification:** 35H30, 35K70, 35L82

**References:**

[1] Demidenko G.V., "Quasielliptic operators and Sobolev type equations. I, II", *Sib. Math. J.*, Vol. 49, No. 5, pp. 842–851, 2008; Vol. 50, No. 5, pp. 838–845, 2009.

[2] Demidenko G.V., "Matrix quasi-elliptic operators in $\mathbb{R}^n$", *Dokl. Math.*, Vol. 81, No. 2, pp. 244–247, 2010.

[3] Demidenko G.V., Uspenskii S.V., Partial differential equations and systems not solvable with respect to the highest-order derivative, Marcel Dekker, New York, Basel, 2003.

[4] Demidenko G.V., "Solvability conditions of the Cauchy problem for pseudohyperbolic equations", *Sib. Math. J.*, Vol. 56, No. 6, pp. 1028–1041, 2015.

# The stabilization rate of a solution to the Cauchy problem for parabolic equation with lower order coefficients

Vasilii N. DENISOV

*Lomonosov Moscow State University, Russia*
*E-mail: vdenisov2008@yandex.ru*

**Abstract:** In the half-space $D = R^N \times (0, \infty), N \geqslant 3$, consider the Cauchy problem for a non divergent parabolic equation.

$$L_1 u \equiv Lu + (b, \nabla u) + c(x,t)u - u_t = 0, \quad (x,t) \in D, \tag{1}$$

$$u(x,0) = u_0(x), \quad x \in R^N, \tag{2}$$

where
$$Lu = \sum_{i,k=1}^{N} a_{ik}(x,t) u_{x_i x_k}, \quad (b, \nabla u) = \sum_{i=1}^{N} b_i(x,t) u_{x_i}. \tag{3}$$

In this report we study sufficient conditions on the lower order coefficients of a parabolic equation guaranteeing the power rate of the uniform stabilization to zero of the solution to the Cauchy problem on every compact in $R^N$ and for any bounded initial function.

We assume that following conditions are satisfied:

$$\exists k_0^2 > 0 : \ k_0^2 = \inf_{D, |\xi|=1} \sum_{i,k=1}^{N} a_{ik}(x,t)\xi_i\xi_k; \ \exists k_1^2 > 0 : \ k_1^2 = \sup_{D, |\xi|=1} \sum_{i,k=1}^{N} a_{ik}(x,t)\xi_i\xi_k.$$

The coefficients $b_i(x,t) (i = 1, ..., N)$ satisfy *condition* $(B)$; i.e., there exist constant $B > 0$ such as

$$\sup_{(x,t)\in D} (1+r) \sum_{i=1}^{N} |b_i(x,t)| = B < k_0^2 N, \quad \text{where } r = \sqrt{x_1^2 + \cdots + x_N^2}.$$

The coefficient $c(x,t)$ satisfies *condition (C)*, i.e. exist $\alpha > 0$ for which

$$c(x,t) \leqslant a_\alpha(r) = -\alpha^2 \min(1, r^{-2}).$$

Let $\lambda_1(\alpha) = (2 - S + \sqrt{D_1})/2, \ D_1 = (2-S)^2 + 4\overline{\alpha}^2, \ \overline{\alpha} = \alpha/k_1.$

**Theorem.** *Suppose that* $(B)$ *and* $(C)$ *with*

$$\alpha^2 > k_1^2(S-1), \ S = \left(k_1^2(N-1) + k_0^2 + B\right)/k_0^2,$$

*hold. Then* $\lim\limits_{t\to\infty} t^{\nu_n} u(x,t) = 0$, $\nu_n = \lambda_1(\alpha)/(2 + \frac{1}{n})$ *for any* $n = 1, 2, ...,$ *uniformly with respect to* $x$ *on any compact* $K$ *in* $R^N$.

**Keywords:** Cauchy problem, non-divergent equation, stabilization

**2010 Mathematics Subject Classification:** 35K15, 35K20, 35B40

# On the solvability of a nonlocal boundary value problem for the Laplace operator with opposite flows at the part of the boundary

Gulnar DILDABEK [1], Isabek ORAZOV [2]

[1,2] *Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail:* [1] *dildabek.g@gmail.com,* [2] *aijan0973@mail.ru*

**Abstract:** In the present paper we investigate a nonlocal boundary problem for the Laplace equation in a half-disk, with opposite flows at the part of the boundary. Our goal is to find a function $u(r,\theta) \in C^0(\bar{D}) \cap C^2(D)$ satisfying in $D$ the equation

$$\Delta u = 0$$

with the boundary conditions

$$u(1,\theta) = f(\theta),\ 0 \le \theta \le \pi,$$
$$u(r,0) = 0, r \in [0,1],$$
$$\frac{\partial u}{\partial \theta}(r,0) = -\frac{\partial u}{\partial \theta}(r,\pi) + \alpha u(r,\pi), \quad r \in (0,1)$$

where $D = \{(r,\theta) : \quad 0 < r < 1, 0 < \theta < \pi\}$; $\alpha < 0$; $f(\theta) \in C^2[0,\pi]$, $f(0) = 0$, $f'(0) = -f'(\pi) + \alpha f(\pi)$.

The difference of this problem is the impossibility of direct applying of the Fourier method (separation of variables). Because the corresponding spectral problem for the ordinary differential equation has the system of eigenfunctions not forming a basis. Throughout this note we mainly use techniques from our work [1]. A special system of functions based on these eigenfunctions is constructed. This system has already formed the basis. This new basis is used for solving of the nonlocal boundary value problem. The existence and the uniqueness of the classical solution of the problem are proved.

**Keywords:** Laplace equation, basis, eigenfunctions, nonlocal boundary value problem

**2010 Mathematics Subject Classification:** 33C10, 34B30, 35J, 35P10

**References:**

[1] Dildabek G., Tengayeva A., and Sadybekov M.A. "On the solvability of a nonlocal boundary problem for the Helmholtz operator in a half-disk", *Contemporary Analysis and Applied Mathematics*. Vol. 2, No. 2, pp. 305–313, 2014.

# About the correctness to some nonlocal boundary value problem for the equation of the mixed type of the first kind and the second order in space

Sirojiddin DJAMALOV [1]

[1] *Institute of Mathematics, National University of Uzbekistan, Uzbekistan*
*E-mail: siroj63@mail.ru*

**Abstract:** As it is known the Dirichlet problem for the equation of the mixed type of the first sort is incorrect. Naturally there is a problem: whether it is impossible to substitute statements of the problem of Dirichlet other conditions enveloping all boundary which ensure a problem correctness? For the first time, such problems have been offered and studied in the works T.S.Kalmenov's. In the present work for the equation of the mixed type of the first kind, the second order in space unambiguously solvability and smoothness of the generalized solution, of some nonlocal boundary value problem with constant coefficients from Sobolev spaces is studied.

Let $\Omega = \prod_{i=1}^{n} (\alpha_i, \beta_i) \in R^n$. $n-$ a measured parallelepiped.

In the area $Q = \Omega \times (0, T)$ we consider a differential equation of the second order

$$(1) \quad Lu = K(x)\, u_{tt} - \sum_{i,j=1}^{n} \frac{\partial}{\partial\, x_i} \left(a_{i,j}(x)\, u_{x_j}\right) + \alpha\,(x,t)\, u_t + c(x,t)\, u = f(x,t),$$

where, $x_1 K(x) > 0$ at $x_1 \neq 0$ and thus $x_1 \in (\alpha_1, \beta_1)$, $\alpha_1 < 0 < \beta_1$.

We assume, that coefficients of equation (1) -are smooth enough functions and let the condition following is satisfied: $a_{i,j}\xi_i\xi_j \geq a_0|\xi|^2$, where,

$$a_{i,j}(x) = a_{j,i}(x);\ a_0 - const > 0,\ \xi \in R^n; |\xi|^2 = \sum_{i=1}^{n} \xi_i^2.$$

**Problem.** To find a generalized solution of equation (1) from the Sobolev space $W_2^l(Q)$, ($2 \leq l$- is integer), satisfying to nonlocal boundary conditions

$$(2) \qquad \gamma D_t^p\, u|_{t=0} = D_t^p\, u|_{t=T}$$

$$(3) \qquad \eta D_{x_i}^p\, u|_{x_i=\alpha_i} = D_{x_i}^p\, u|_{x_i=\beta_i},$$

at $p = 0, 1$; here $\gamma$ *and* $\eta - const \neq 0$, where $D_t^p u = \frac{\partial^{\,p} u}{\partial t^{\,p}}$ $D_t^0\, u = u$.

**Keywords:** second order mixed type equation of the first kind,nonlocal boundary value problem with constant coefficients, Sobolev spaces,unique solvability,existence of solution, smoothness of the generalized solution.

**2010 Mathematics Subject Classification:** 35M10

## Solution of the linearized nonregular free boundary problem for the parabolic equations

Zhanat DZHOBULAEVA

*Institute of mathematics and mathematical modeling, Àlmaty, Kazakhstan*
*E-mail: zhanat-78@mail.ru*

**Abstract:** We study linearized free boundary problem for the system of the parabolic equations with two small parameters at the principle terms in the boundaries conditions. The original nonlinear free boundary problem describes the process of phase transition (melting, solidification) of substance with admixture of unknown concentration.

There are proved in the Hölder spaces the existence, uniqueness and uniform with respect to the small parameters coercive estimates of the solution to the linearized free boundary problem.

These results are on the basis of the solution of the nonlinear singular perturbed free boundary problem for the system of the parabolic equations.

**Keywords:** Hölder space, parabolic equation, boundary value problem, small parameters, existence, uniqueness, coercive estimates

**2010 Mathematics Subject Classification:** 35K51, 35B20, 35B65

## Properties of an isolated solution to a nonlinear boundary value problem with parameters

Dulat S.DZHUMABAEV, Bayan B. MINGLIBAYEVA

*Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: dzhumabaev@list.ru; bayan_math@mail.ru*

**Abstract:** A nonlinear two-point boundary value problem for ordinary differential equation with parameters is studied. Necessary and sufficient conditions for the existence of an isolated solution are obtained. The conditions of continuity dependence of isolated solution are established.

We consider the nonlinear two-point boundary value problem with parameter for system of ordinary differential equations

$$\frac{dx}{dt} = f(t, x, \mu), \qquad t \in (0, T), \qquad x \in R^n, \qquad \mu \in R^m \tag{1}$$

$$g(\mu, x(0), x(T)) = 0, \tag{2}$$

where $f : [0, T] \times R^n \times R^m \to R^n$, $g : R^m \times R^n \times R^n \to R^{n+m}$ are continuous.

The essentially nonlinear problems usually have several, moreover, infinitely many solutions. Isolation is one of the most important properties of solutions to nonlinear problems. Continuous dependence of solution on the change of considered differential equations and boundary conditions is very important in applications for the adequate describing of processes studied. This property is also necessary while constructing the approximate and numerical methods for finding the solutions. However, in general, the isolated solution considered as an isolated element of the set of solutions to problem (1), (2) does not have that property. Therefore, it is considered an "isolated" in a narrow sense solutions to problem (1), (2).

**Keywords:** nonlinear boundary value problem with parameters, "isolated" solution.

**2010 Mathematics Subject Classification:** 34A34, 34B08

**References:**

[1] Dzhumabaev D.S., and Temesheva S.M. "A parametrization method for solving nonlinear two-point boundary value problems ", *Comp. Math. and Math. Phys.*, Vol.47, No.1, pp. 37-61, 2007.

## Unique solvability of an inverse problem for a semilinear equation with final overdetermination

Ali Ugur SAZAKLIOGLU [1], Abdullah Said ERDOGAN [2], Allaberen ASHYRALYEV [3]

[1] *University of Turkish Aeronautical Association, Turkey*
*E-mail: ausazaklioglu@thk.edu.tr*
[2] *E-mail: aserdogan@gmail.com*
[3] *Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: aashyr@yahoo.com*

**Abstract:** This study is devoted to the investigation of unique solvability of the inverse problem governed by a semilinear equation subject to a final overdetermination

$$\begin{cases} \frac{du}{dt} + Au(t) = f(t, u(t)) + p, \ 0 < t < T, \\ \\ u(0) = \varphi, \ u(T) = \psi \end{cases} \tag{4}$$

in an arbitrary Banach space $E$. Here, $A$ is a linear operator acting in $E$, with domain $D(A)$. Assume that $-A$ is the generator of the analytic semigroup $\exp\{-tA\}$ with an exponentially decreasing norm

$$\left\| e^{-tA} \right\|_{E \to E} \leq M e^{-\delta t}, \ t \left\| Ae^{-tA} \right\|_{E \to E} \leq M, \ t \geq 0. \tag{5}$$

For the numerical solution of this problem the first order of accuracy Rothe difference scheme is proposed. The existence and uniqueness results for this difference scheme is given. As an application, an inverse problem governed by a semilinear parabolic equation subject to a final overdetermination is considered. Throughout this note we mainly use techniques from [1, 2].

**Keywords:** Existence and uniqueness, semilinear equation, inverse problem, final overdetermination, difference scheme

**2010 Mathematics Subject Classification:** 65L12, 31A25, 35K58, 34A12

**References:**

[1] A. Ashyralyev, and P. E. Sobolevskii, *New Difference Schemes for Partial Differential Equations*, Birkhäuser, Basel, Boston, Berlin, 2004.
[2] A. Ashyralyev, "A survey of results in the theory of fractional spaces generated by positive operators", *TWMS Journal of Pure and Applied Mathematics,* Vol. 6, pp. 129–157, 2015.

## Green's function of a heat problem with a periodic boundary condition

Nurzhan E. ERZHANOV [1]

[1] *Institute of Mathematics and Mathematical Modeling, Kazakhstan,*
[1] *Regional Social and Innovative University, Kazakhstan*
*E-mail: imanbaevnur@mail.ru*

**Abstract:** In the paper a non-local initial-boundary value problem for a non-homogeneous one-dimensional heat equation is considered. The domain under consideration is a rectangle. The classical initial condition with respect to $t$ is put. A non-local periodic boundary condition with respect to a spatial variable $x$ is put. It is well-known that a solution of problem can be constructed in the form of convergent orthonormal series according to eigenfunctions of a spectral problem for an operator of multiple differentiation with periodic boundary conditions. Therefore Green's function can be also written in the form of an infinite series with respect to trigonometric functions (Fourier series). For classical first and second initial-boundary value problems there also exists a second representation of the Green's function by Jacobi function. In this paper we find the representation of the Green's function of the non-local initial-boundary value problem with periodic boundary conditions in the form of series according to exponents.

Throughout this note we mainly use techniques from [3-5].

**Keywords:** heat equation, initial-boundary value problems, periodic boundary condition, Green's function

**2010 Mathematics Subject Classification:** 5C15, 35E15, 35K05, 35K20

**References:**

[1] Kal'menov T.Sh., Sadybekov M.A., "Dirichlet problem and nonlocal boundary-value-problems for the wave-equation", *Differential Equations*. Vol.26, No.1, pp. 55–59, 1990.

[2] Kalmenov T.Sh., Shaldanbaev A.S., "On a criterion of solvability of the inverse problem of heat conduction", *Journal of Inverse and Ill-Posed Problems*. Vol.18, No.5, pp. 471–492, 2010.

[3] Orazov I., Sadybekov M.A., "On a class of problems of determining the temperature and density of heat sources given initial and final temperature", *Siberian Mathematical Journal*. Vol.53, No.1, pp. 146–151, 2012.

# On the properties of operators with the semi-power type lacunas

Leonid FALALEEV [1]

[1] *Institute of Mathematics and Mathematical Modeling, Àlmaty, Kazakhstan*
*E-mail: v_gulmira@mail.ru*

**Abstract:** For linear matrix operators

$$u_N(f,\Lambda,X)=\frac{a_0}{2}+\sum_{k=1}^{N}\lambda_k^{(n)}(a_k\cos kx+b_k\sin kx),$$

($a_k$, $b_k$ are Fourier coefficients of $f(x)$), constructed with respect to subsequences of the Fourier sums $S_{n_k}$ with the lacunas $n_k=[\frac{k^\gamma}{\ln^\beta(k+1)}]$, $k=1,2,...,N$, a sufficient condition for boundedness of norms $\|u_N\|_{C_{2\pi}}$

$$(1)\qquad \|u_N\|\le c\{n_N^{\frac{1}{\gamma}}\sum_{k=0}^{N}(\Delta\lambda_k)^2\ln^{\beta/\gamma}(k+1)[1+(N-k)^{1-1/\gamma}k^{1/\gamma-1}]\},$$

is given. $N$ is the number of non-zero differences $\Delta\lambda_k=\lambda_{n_k}-\lambda_{n_k+1}$, $c>0$, $c=const.$ $[y]$ is the integer part of $y$. $\beta\ge 0$, $\gamma\ge 1$. $\lambda_k^n\equiv\lambda_k$ are limited multipliers of summation.

Using (1) approximative properties of the discrete analog of biharmonic operators $D(f,r,x)$ are investigated. In particular the precise estimate

$$(2)\quad \sup_{f\in C^1}\|f(x)-D(f,r,x)\|_{C_{2\pi}}=O\big((\gamma-\beta)\sqrt{(1-r)\ln^{\beta/\gamma}\frac{1}{1-r}}\big),\ r\to 1-.$$

is obtained in the class of differentiable functions $C^1$.

If $\gamma=1$, $\beta=0$ then (2) coincides with the estimate of the work [1].

**Keywords:** semi-power type lacunas, biharmonic operator

**2010 Mathematics Subject Classification:** 42A24

**References:**

[1] Falaleev L.P., Vasilina G.K. "On biharmonic operators rarefied by lacunas *VIII Int. symp. "Fourier series and their applications"*. Rostov-on-Don, pp. 28-34, 2014.

## Initial value problem for a class of fractional order inhomogeneous equations in Banach spaces

Vladimir E. FEDOROV [1], Roman R. NAZHIMOV [2], Dmitriy M. GORDIEVSKIKH [3]

[1] *Laboratory of Quantum Topology, Mathematical Analysis Department, Chelyabinsk State University, Chelyabinsk, Russia*
*E-mail: kar@csu.ru*
[2] *Mathematical Analysis Department, Chelyabinsk State University, Chelyabinsk, Russia*
*E-mail: goldenboy454@mail.ru*
[3] *Shadrinsk State Pedagogical University, Shadrinsk, Kurgan Region, Russia*
*E-mail: dmitriy_g90@mail.ru*

**Abstract:** The fractional order linear inhomogeneous equation in Banach space with the Riemann–Liouville derivative, and with a bounded operator at the unknown function endowed by initial conditions for fractional derivatives of a solution is studied. Theorem on unique solvability of the problem is proved. As application of the obtained result an initial boundary value problem for Barenblatt–Zheltov–Kochina equation with the Riemann–Liouville fractional derivative with respect to time is considered. Unique solvability conditions and a form of the problem solution are found in terms of the Fourier series with respect to the eigenfunctions of the Laplace operator.

Famous works (see [1] and others) concerning related problems do not contain similar results. Note papers [2, 3], where analogous problems are solved in the case of the Caputo fractional derivative, correspondently, with initial conditions for the integer derivatives of a solution.

**Keywords:** Riemann–Liouville fractional derivative, Riemann–Liouville fractional integral, Mittag-Leffler function, initial boundary value problem, partial differential equation

**2010 Mathematics Subject Classification:** 35R11, 34A08

**References:**

[1] Bajlekova E.G. *Fractional Evolution Equations in Banach Spaces*, PhD thesis, University Press Facilities, Eindhoven University of Technology, 2001.

[2] Fedorov V.E., Gordievskikh D.M. "Resolving operators of degenerate evolution equations with fractional derivative with respect to time", *Russian Mathematics*, Vol. 59, pp. 60–70, 2015.

[3] Fedorov V.E., Gordievskikh D.M., Plekhanova M.V., "Equations in Banach spaces with a degenerate operator under a fractional derivative", *Differential Equations*, Vol. 51, pp. 1360–1368, 2015.

## On an illposed model of oscillations of a flat plate with a variety of mounts on opposite sides

Ulzada ISKAKOVA

*Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: iskakova@math.kz*

**Abstract:** We consider a model case of stationary vibrations of a thin flat plate, one side of which is embedded, the opposite side is free, and the sides are freely leaned. In mathematical modeling there is a local boundary value problem for the biharmonic equation in a rectangular domain. Boundary conditions are given on all boundary of the domain.

**Problem** $C$. *Find in* $D = \{(x, y) : 0 < x < \pi, 0 < y < l\}$ *a solution to the biharmonic equation*

$$\triangle^2 u \equiv u_{xxxx}(x, y) + 2u_{xxyy}(x, y) + u_{yyyy}(x, y) = 0, \quad (x, y) \in D,$$

*satisfying boundary conditions in the first spatial variable* $x$*:*

$$u|_{x=0} = 0, \ \triangle u|_{x=0} = 0; \quad u|_{x=\pi} = 0, \ \triangle u|_{x=\pi} = 0; \quad 0 \le y \le l;$$

*and boundary conditions in the second spatial variable* $y$*:*

$$u|_{y=0} = \varphi_1(x), \ \frac{\partial u}{\partial y}\Big|_{y=0} = \varphi_2(x), \quad 0 \le x \le \pi;$$

$$\triangle u|_{y=l} = \psi_1(x), \quad \frac{\partial \triangle u}{\partial y}\Big|_{y=l} = \psi_2(x), \ \ 0 \le x \le \pi.$$

We show that the considered problem is self-adjoint. Herewith the problem is ill-posed. We show that the stability of solution to the problem is disturbed. Necessary and sufficient conditions of existence of the problem solution are found. Spaces of the ill-posedness of the considered problem are constructed.

Throughout this note we mainly use techniques from our works [1-2].

**Keywords:** oscillations, thin flat plate, biharmonic equation, boundary value problem, ill-posed problem

**2010 Mathematics Subject Classification:** 35J40, 35P20

**References:**

[1] Kal'menov T.Sh., Iskakova U.A., "A method for solving the Cauchy problem for the Laplace equation", *Doklady Mathematics*, Vol. 78, No. 3, pp. 874–876, 2008.
[2] Kal'menov T.Sh., Iskakova U.A., "Criterion for the strong solvability of the mixed Cauchy problem for the Laplace equation", *Differential Equations*, Vol. 45, No. 10, pp. 1460–1466, 2009.

# Approximation properties of some summation methods in the Smirnov classes with variable exponent

Daniyal ISRAFILOV [1,2], Ahmet TESTICI [1]

[1] *Department of Mathematics, Balikesir University, Balikesir, Turkey*
*E-mail: mdaniyal@balikesir.edu.tr, testiciahmet@hotmail.com*
[2] *Institute of Mathematics and Mechanics Baku, Azerbaijan*

**Abstract:** Let $G \subset \mathbb{C}$ be a finite domain in the complex plane, bounded by a rectifiable Jordan curve $\Gamma$. The variable exponent Lebesgue spaces $L^{p(\cdot)}(\Gamma)$ for a given Lebesgue measurable variable exponent $p(z) \geq 1$ on $\Gamma$ we define as the set of Lebesgue measurable functions $f$, such that $\int_\Gamma |f(z)|^{p(z)} |dz| < \infty$. The function

$$\|f\|_{L^{p(\cdot)}(\Gamma)} := \inf \left\{ \lambda > 0 : \int_\Gamma |f(z)/\lambda|^{p(z)} |dz| \leq 1 \right\}$$

defines a norm on $L^{p(\cdot)}(\Gamma)$.

Given Lebesgue measurable function $p(\cdot) : \Gamma \to [1, \infty)$ we define the variable exponent Smirnov classes of analytic functions in $G$ as $E^{p(\cdot)}(G) := \left\{ f \in E^1(G) : f \in L^{p(\cdot)}(\Gamma) \right\}$.

Each function $f \in E^{p(\cdot)}(G)$, has the non-tangential limits almost everywhere $(a.e)$ on $\Gamma$ and hence if we define $\|f\|_{E^{p(\cdot)}(G} := \|f\|_{L^{p(\cdot)}(\Gamma)}$, then the space $E^{p(\cdot)}(G)$ is also a normed space of analytic functions in $G$.

In this work we continue our investigations [1,2], on the approximation problems in the variable exponent Smirnov classes $E^{p(\cdot)}(G)$. Namely, we study the approximation properties of the different approximation aggregates and obtain the appropriate estimations in term of the higher modulus of smoothness for a given function $f \in E^{p(\cdot)}(G)$.

**Keywords:** Variable exponent Smirnov classes, Faber series, De Vallée Poussin means, Jackson means

**2010 Mathematics Subject Classification:** 30E10, 41A10

**Acknowledgement** This work was supported by TÜBITAK grant 114F422: "*Approximation Problems in the Variable Exponent Lebesgue Spaces*".

**References:**

[1] Guven A., and Israfilov D., "Trigonometric Approximation in Generalized Lebesgue Spaces $L^{p(x)}$", *Journal of Math. Inequalities*, Vol.4, No.2, pp. 285-299, 2010.

[2] Israfilov D., Testici A., "Approximation in Smirnov classes with variable exponent", *Complex Variable and elliptic equations*, Vol.60, No.9, pp. 1243-1253, 2015.

## Algorithm of finding of a solution bounded on stripe of linear hyperbolic equation with the mixed derivative

Symbat KABDRAKHOVA

*Al-Farabi Kazakh National University, Kazakhstan*
*E-mail: S-Kabdrachova@mail.ru*

**Abstract:** A semiperiodic boundary value problem is examined for linear hyperbolic equation with two independent variables. Euler's modification broken method is offered for construction of a solution bounded on a stripe. The conditions of existence of a solution bounded on a stripe are received.

We consider a linear hyperbolic equation with two independent variables

$$\frac{\partial^2 u}{\partial x \partial t} = A(x,t)\frac{\partial u}{\partial x} + B(x,t)\frac{\partial u}{\partial t} + C(x,t)u + f(x,t), \tag{1}$$

where $A(x,t), B(x,t), C(x,t)$ are functions continuous on $\overline{\Omega}$, $f : \overline{\Omega} \times R^3 \to R$, is function continuous on $t \in \mathbb{R}$ at fixed $x \in [0,\omega]$ and uniformly relatively $t \in \mathbb{R}$ continuous on $x \in [0,\omega]$.

The solution of equation (1) is investigated satisfying to the following conditions

$$u(0,t) = \psi(t), \quad t \in \mathbb{R} = (-\infty,\infty), \quad \frac{\partial u(x,t)}{\partial x} \in C^*(\overline{\Omega},\mathbb{R}), \tag{2}$$

where $C^*(\overline{\Omega},\mathbb{R})$ a space of bounded functions, the continuously differentiable function $\psi(t)$ is bounded on $\mathbb{R}$ together with the derivative $\dot{\psi}(t)$. The problem (1), (2) we will reduce to the equivalent problem, entering new unknown functions

$$v(x,t) = \frac{\partial u(x,t)}{\partial x},\ w(x,t) = \frac{\partial u(x,t)}{\partial t}.$$

For finding bounded on a stripe solution we apply Euler's modification broken method [1] and we get a problem for the systems ordinary differential equations. Unlocal boundary value problems are built for equation (1) on a finite area, allowing with set to approximate narrowing exactness on this area of bounded on the stripe solution.

**Keywords:** linear hyperbolic equation, boundary value problem, Euler's modification broken method, solution bounded on a stripe

**2010 Mathematics Subject Classification:** 34A30, 34B05, 35L20

**References:**

[1] Dzhumabaev D.S. "Approximation of a bounded solution of linear ordinary differential equation by solutions of two point boundary value problems", *JVM.*, No.3, pp. 388-404, 1990.

## On boundary value problem for the fourth order mixed type equation with fractional derivative

Bakhtiyor J. KADIRKULOV[1], Jambul A. SERIKBAEV [2]

[1] *Tashkent State Institute of Oriental Studies, Uzbekistan*
[2] *Scientific Research Institute of Mathematics and Mechanics, Kazakh National University, Kazakhstan*

**Abstract:** In the present work we investigate nonlocal problem for the fourth order mixed type equations.

Let $\Omega = \{(x,t) : 0 < x < 1, -p < t < q\}, \Omega^+ = \Omega \cap (t > 0), \Omega^- = \Omega \cap (t < 0)$, where $p, q > 0$. In the domain $\Omega$ we consider equation

$$Lu(x,t) = f(x,t) \tag{1}$$

where

$$Lu(x,t) = \begin{cases} \frac{\partial^4 u}{\partial x^4} + {}_C D^{\alpha}_{0t} u & t > 0, \\ \frac{\partial^4 u}{\partial x^4} + \frac{\partial^2 u}{\partial t^2}, & t < 0, \end{cases}$$

${}_C D^{\alpha}_{0t} u$ is the Caputo fractional operator of the order $\alpha \in (0, 1]$

**Problem A.** Find a function $u(x,t)$, which is:

1) continuous in $\overline{\Omega}$, together with derivatives appearing in boundary conditions;

2) satisfies equation (1) in domains $\Omega^+ \cup \Omega^-$;

3) satisfies boundary conditions

$$u(0,t) = u_{xx}(1,t) = 0, u_x(1,t) = u_x(0,t), u_{xxx}(1,t) = u_{xxx}(0,t), -p \le t \le q, \tag{2}$$

$$u(x,-p) = 0, 0 \le x \le 1; \tag{3}$$

4) satisfies gluing condition

$${}_C D^{\alpha}_{0t} u(x,+0) = \frac{\partial u(x,-0)}{\partial t}, 0 < x < 1. \tag{4}$$

The main result of this study is the following theorem.

**Theorem.** Let $f(x,t) \in C^{4,0}_{x,t}(\bar{\Omega})$, $\frac{\partial^5 f(x,t)}{\partial x^5} \in L_2(\Omega)$,

$f(0,t) = f_{xx}(1,t) = f_{xxx}(0,t) = 0, f_x(1,t) = f_x(0,t), f_{xxx}(1,t) = f_{xxx}(0,t)$, and $\Delta_n = \lambda_n^2 \sin \lambda_n^2 p + \cos \lambda_n^2 p \neq 0$ at $n = 1, 2, ...$

Then the solution of Problem A exists and unique.

## On boundary value problem for the fourth order mixed type equation with fractional derivative

Bakhtiyor KADIRKULOV [1], Jambul SERIKBAEV [2]

[1] *Tashkent State Institute of Oriental Studies, Uzbekistan*
*E-mail: berdyshev@mail.ru*

[2] *ScRI of Math. and Mech, Al Farabi Kazakh National University, Kazakhstan*

**Abstract:** Many problems in diffusion and dynamical processes, electrochemistry, biosciences, signal processing, system control theory lead to differential equations of fractional order. In the present work we investigate nonlocal problem for the fourth order mixed type equations.

Let $\Omega = \{(x,t) : 0 < x < 1, -p < t < q\}, \Omega^+ = \Omega \cap (t > 0), \Omega^- = \Omega \cap (t < 0)$, where $p, q > 0$. In the domain $\Omega$ we consider equation

$$Lu(x,t) = f(x,t) \tag{1}$$

where

$$Lu(x,t) = \begin{cases} \frac{\partial^4 u}{\partial x^4} +_C D^{\alpha}_{0t} u & t > 0, \\ \frac{\partial^4 u}{\partial x^4} + \frac{\partial^2 u}{\partial t^2}, & t < 0, \end{cases}$$

${}_C D^{\alpha}_{0t} u$ is the Caputo fractional operator of the order $\alpha \in (0,1]$.

**Problem A.** Find a function $u(x,t)$, which is:
1) continuous in $\overline{\Omega}$, together with derivatives appearing in boundary conditions;
2) satisfies equation (1) in domains $\Omega^+ \cup \Omega^-$;
3) satisfies boundary conditions

$$u(0,t) = u_{xx}(1,t) = 0, u_x(1,t) = u_x(0,t), u_{xxx}(1,t) = u_{xxx}(0,t), -p \le t \le q, \tag{2}$$

$$u(x,-p) = 0, 0 \le x \le 1; \tag{3}$$

4) satisfies gluing condition

$${}_C D^{\alpha}_{0t} u(x,+0) = \frac{\partial u(x,-0)}{\partial t}, 0 < x < 1. \tag{4}$$

The main result of this study is the following theorem.

**Theorem 1.** *Let* $f(x,t) \in C^{4,0}_{x,t}(\bar{\Omega})$, $\frac{\partial^5 f(x,t)}{\partial x^5} \in L_2(\Omega)$,

$$f(0,t) = f_{xx}(1,t) = f_{xxx}(0,t) = 0,$$

$$f_x(1,t) = f_x(0,t), f_{xxx}(1,t) = f_{xxx}(0,t),$$

*and* $\Delta_n = \lambda_n^2 \sin \lambda_n^2 p + \cos \lambda_n^2 p \neq 0$ *at* $n = 1, 2, \ldots$ *Then the solution of Problem A exists and unique.*

**Keywords:** mixed type equation, Caputo fractional derivative, Mittag-Leftler function.

**2010 Mathematics Subject Classification:** 35R11, 35M12, 35M32

**References:**

[1] Kilbas A.A., Srivastava H.M., and Trujillo, J.J., *Theory and applications of fractional differential equations*, Elsevier Science & Tech, 2006.

## Asymptotics solutions of a singularly perturbed integro-differential problem with rapidly oscillating coefficients

Burkhan KALIMBETOV [1], Dinmukhambet SAPAKOV [1]

[1] *Department of Mathematics, H.A. Yasawi International Kazakh-Turkish University, Turkestan, Kazakhstan*
*E-mail: bkalimbetov@mail.ru, sapakov@mail.ru*

**Abstract:** In this paper we consider the Cauchy problem for a singularly perturbed integro-differential systems with rapidly oscillating coefficients:
(1)

$$\varepsilon\dot{z} - A(t)z - \varepsilon\varphi(t)cos\frac{2\beta(t)}{\varepsilon}Bz - \int\limits_{t_0}^{t} k(s)z(s,\varepsilon)ds = h(t),\ z(t_0,\varepsilon) = z^0,\ t \in [0,T],$$

where $z(t,\varepsilon) = \{u(t,\varepsilon), \vartheta(t,\varepsilon)\}$ is an unknown vector-function, $z^0 = \{u^0, \vartheta^0\}$ is a known constant vector, $h(t) = \{h_1(t), h_2(t)\}$ are known vector-function, $A(t)$, $k(t)$, $B$ are given matrices, $\varphi(t)$, $\beta(t) > 0$ are known functions, $\varepsilon > 0$ is a small parameter. It is required to construct the main term of asymptotics of a solution of (1) at $\varepsilon \to +0$.

A particular case of problem (1) (with $\beta(t) = \beta \cdot t, \beta = const, h(t) \equiv 0, k(t) \equiv 0$), describing the phenomenon of parametric amplification, been considered in [1–3]. We will illustrate the application of the regularization method [4, 5] for obtaining the principal term of the asymptotic behavior of the solution of problem (1).

**Keywords:** Integro-differential system, small parameter, asymptotic solution

**References:**

[1] Daletskii, Yu.L, *Reports AS USSR,* Vol. 143 no. 5, pp. 1026-1029, 1962.
[2] Feshenko, S.F., Shkil, N.I. and Nikolenko, L.D., Asymptotic methods in the theory of linear differential equations, Kiev, Naukova dumka, 1966.
[3] Lomov, S.A. and Lomov, I.S., Fundamentals the mathematical theory of the boundary layer. M.: Publishing house of MSU, 2011.
[4] Ryzhikh, A.D., Proceedings MPEI, Vol. 357, pp. 92-94, 1978.
[5] Shkil, N.I., Asymptotic methods in differential equations, Kiev, Naukova dumka, 1971.

## On a problem of the Frankl type for an equation of the mixed parabolic-hyperbolic type

Tynysbek KAL'MENOV [1], Makhmud SADYBEKOV [2]

[1] *Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: kalmenov@math.kz*
[2] *Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: sadybekov@math.kz*

**Abstract:** In the paper a new non-local boundary value problem for an equation of the parabolic-hyperbolic type is formulated. This equation is of the first kind, that is, the line of type change is not a characteristic of the equation. The suggested new non-local condition binds points on boundaries of the parabolic and hyperbolic parts of the domain with each other. This problem is generalization of the well-known problems of Frankl type. Unlike the existing publications of the other authors related to the theme, in the suggested formulation of the problem the hyperbolic part of the domain coincides with a characteristic triangle. Unique solvability of the formulated problem is proved in the sense of classical and strong solutions.

Let $\Omega \subset R^2$ be a finite domain bounded for $y > 0$ by the segments $AA_0$, $A_0B_0$, $B_0B$, $A = (0,1)$, $B_0 = (1,1)$, $B = (1,0)$ and for $y < 0$ by the characteristics $AC: \ x + y = 0$ and $BC: \ x - y = 1$ of an equation of the mixed parabolic-hyperbolic type

$$Lu = \left\{ \begin{array}{c} u_x - u_{yy},\ y > 0 \\ \\ u_{xx} - u_{yy},\ y < 0 \end{array} \right\} = f(x,y). \tag{1}$$

Problem $F$. *Find a solution to Eq.*(1) *satisfying classical boundary conditions*

$$u|_{AA_0} = 0, \quad u_y|_{A_0B_0} = 0, \tag{2}$$

*and a non-local boundary condition*

$$\alpha u\left(\theta_0\left(t\right)\right) + \beta u\left(\theta_1\left(t\right)\right) = \gamma u\left(\theta\left(t\right)\right), \quad 0 \le t \le 1, \tag{3}$$

*where* $\theta\left(t\right) = (t,1)$, $\theta_0\left(t\right) = \left(\frac{t}{2}, -\frac{t}{2}\right)$, $\theta_1\left(t\right) = \left(\frac{t+1}{2}, \frac{t-1}{2}\right)$; $\alpha$, $\beta$ *and* $\gamma$ *are given numbers.*

**Keywords:** non-local boundary conditions, equation of the parabolic-hyperbolic type, Green's function, Frankl problem

**2010 Mathematics Subject Classification:** 35M10, 35M12

## Weighted multiplicative estimate for norm of discrete Hardy operator

Aigerim KALYBAY [1], Saltanat SHALGINBAYEVA [2]

[1] *KIMEP University, Almaty, Kazakhstan*
*E-mail: kalybay@kimep.kz*
[2] *Kazakh Ablai Khan University of International Relations and World Languages, Almaty, Kazakhstan*
*E-mail: salta_sinar@mail.ru*

**Abstract:** Let $f \geq 0$ be a sequence of real numbers $f = \{f_i\}_{i=1}^{\infty}$ with non-negative terms.

Let $v > 0$, $u \geq 0$, and $w \geq 0$ be weight sequences. Let $A$ be a matrix operator in the form:

$$(Af)_i = \sum_{j=1}^{i} a_{i,j} f_j, \quad i \geq 1,$$

where $a_{i,j} \geq 0$ for $i \geq j \geq 1$ and $a_{i,j} = 0$ for $i < j$.

We consider the following weighted multiplicative inequality:

$$\|uPf\|_q \leq C\|vf\|_p^{\alpha} \, \|wAf\|_r^{1-\alpha}, \quad f \geq 0, \quad 0 \leq \alpha \leq 1, \tag{4}$$

where $(Pf)_i = \sum\limits_{j=1}^{i} f_j$ is the discrete Hardy operator and $\|\cdot\|_q$ is the standard norm of the space $l_q$.

When $\alpha = 1$ inequality (1) is a weighted estimate of the discrete Hardy operator, which is well-studied for all values of the parameters $0 < p, q < \infty$ and has numerous applications (see [1,2]). Multiplicative inequalities are also of great importance and have applications to different problems of Analysis. Some of applications can be found in [3].

When $\alpha = 0$ from (1) we get the inequality that is of independent interest.

**Keywords:** multiplicative inequality, Hardy-type inequality, matrix operator, sequence

**2010 Mathematics Subject Classification:** 26D10, 26D15, 39B82

**References:**

[1] Hardy G.H., Littlewood J.E., Pólya G., "Inequalities", *Cambridge Univ. Press*, 1934.

[2] Kufner A., Maligranda L., Persson L.E., "The Hardy inequality. About its history and some related results", *Vydavatelský servis*, Pilsen, 2007.

[3] Magaril-Il'yaev G.G., Tikhomirov V.M., "Convex Analysis: Theory and Applications", *American Mathematical Society*, 2003.

## Weighted additive estimate for norm of discrete Hardy operator

Aigerim KALYBAY [1], Saltanat SHALGINBAYEVA [2]

[1] *KIMEP University, Almaty, Kazakhstan*
*E-mail: kalybay@kimep.kz*
[2] *Kazakh Ablai Khan University of International Relations and World Languages, Almaty, Kazakhstan*
*E-mail: salta_sinar@mail.ru*

**Abstract:** Let $f \geq 0$ be a sequence of real numbers $f = \{f_i\}_{i=1}^{\infty}$ with non-negative terms.

Let $v > 0$, $u \geq 0$, and $w \geq 0$ be weight sequences. Let $P$ be a discrete Hardy operator $(Pf)_i = \sum\limits_{j=1}^{i} f_j$, and $A$ be a matrix operator in the form $(Af)_i = \sum\limits_{j=1}^{i} a_{i,j} f_j$, $i \geq 1$, where $a_{i,j} \geq 0$ for $i \geq j \geq 1$ and $a_{i,j} = 0$ for $i < j$.

We consider the following weighted additive estimate:

$$\|uPf\|_q \leq C\left(\|vf\|_p + \|wAf\|_r\right), \quad f \geq 0, \tag{5}$$

where $\|\cdot\|_q$ is the standard norm of the space $l_q$.

In papers [1–3] under some conditions on the elements $(a_{i,j})$ the authors have found necessary and sufficient conditions for the validity of the inequality:

$$\|uAf\|_q \leq C\left(\|vf\|_p + \|wPf\|_p\right), \quad f \geq 0,$$

where $1 < p, q < \infty$.

Moreover, continuous analogue of inequality (1) has been studied in paper [**?**] for $A \equiv P$, $r = p$ and $1 < p \leq q < \infty$.

**Keywords:** additive inequality, Hardy-type inequality, matrix operator, sequence

**2010 Mathematics Subject Classification:** 26D10, 26D15, 39B82

**References:**

[1] Taspaganbetova Z., Temirkhanova A., "Boundedness and compactness of a class of matrix operators", *Math. Journal*, Vol.2, No.4, pp. 73–85, 2011.

[2] Taspaganbetova Z., Temirkhanova A., "Boundedness of matrix operators in weighted spaces of sequences and their apllications", *Ann. Funct. Anal.*, Vol.1, No.2, pp. 114–127, 2011.

[3] Temirkhanova A., "An additive estimate of a class of matrix operators", *Ph.D. thesis, Luleå University of Technology*, Luleå, 2015.

## Variations on strongly lacunary quasi Cauchy sequences

Huseyin KAPLAN [1], Huseyin CAKALLI [2]

[1] *Nigde University, Department of Mathematics, Nigde, Turkey*
*E-mail: hkaplan@nigde.edu.tr*
[2] *Maltepe University, Division of Mathematics, İstanbul, Turkey*
*E-mail: hcakalli@gmail.com*

**Abstract:** In this paper $N_\theta^\alpha(p)$-ward continuity is introduced in the sense that a real valued function $f$ defined on a subset $A$ of $\mathbf{R}$, the set of real numbers is $N_\theta^\alpha(p)$-ward continuous if $(f(x_k))$ is $N_\theta^\alpha(p)$-quasi-Cauchy whenever $(x_k)$ is $N_\theta^\alpha(p)$-quasi-Cauchy sequence of points in $A$, where a sequence $(x_k)$ is called $N_\theta^\alpha(p)$-quasi-Cauchy if $\lim_{r\to\infty} \frac{1}{h_r^\alpha} \sum_{k\in I_r} |\Delta x_k|^p = 0$, where $\Delta x_k = x_{k+1} - x_k$ for each positive integer $k$, $p$ is a constant positive integer, $\alpha$ is a constant in $]0,1]$, $I_r = (k_{r-1}, k_r]$, and $\theta = (k_r)$ is a lacunary sequence, which is an increasing sequence of positive integers such that $k_0 \neq 0$, and $h_r : k_r - k_{r-1} \to \infty$.

**Keywords:** Sequences, series, strongly lacunary convergence, continuity

**2010 Mathematics Subject Classification:** 40A05, 40A35, 26A15

**References:**

[1] Burton D., and Coleman J., "Quasi-Cauchy Sequences", *Amer. Math. Monthly*, Vol. 117, No.5, pp. 328-333, 2010.

[2] Cakalli H., "Forward continuity", *J.Comput.Anal.Appl.*, Vol.13, No.2, pp. 225-230, 2011.

[3] Caserta A., and Kocinac Ljubisa D.R., "On statistical exhaustiveness", *Appl. Math. Lett.*, Vol.25, No.10, pp. 1447-1451, 2012.

[4] Cakalli H., "N-theta-Ward continuity", *Abstract and Applied Analysis*, Vol.2012, Article ID 680456, (2012), 8 pp. doi:10.1155/2012/680456.

[5] Et M., and Sengul H.,"Some Cesaro-Type Summability Spaces of Order alpha and Lacunary Statistical Convergence of Order alpha",*Filomat* Vol.28, No.2, pp.1593-1602, 2014.

[6] Çakalli H., and Kaplan H., "A study on N-theta quasi-Cauchy sequences, *Abstr. Appl. Anal.*, Vol.2013, Article ID 836970, 4 pp. 2013. doi:10.1155/2013/836970

[7] Canak I., and Dik M., "New types of continuities", *Abstr. Appl. Anal.*, Vol.2010, Article ID 258980, 6 p., 2010.

[8] Cakalli H., "Sequential definitions of compactness", *Appl. Math. Lett.*, Vol.21, No.6, pp. 594-598, 2008.

[9] Connor J., and Grosse-Erdmann K.G., "Sequential definitions of continuity for real functions", *Rocky Mountain J. Math.*, Vol.33, No.1, pp. 93-121, 2003.

[10] Cakalli H., "On G-continuity", *Comput. Math. Appl.*, Vol.61, pp. 313-318, 2011.

[11] Mucuk O., Sahan T., "On G-Sequential Continuity", *Filomat*, Vol.28, No.6, pp. 1181-1189, 2014.

[12] Sengul H., and Et M., "On lacunary statistical convergence of order alpha", *Acta Mathematica Scientia*, Vol.34, No.2, pp. 473-482, 2014.

## On an Uniqueness and Correct Solvability of the Biharmonic Boundary Value Problem

Valery V. KARACHIK [1], Berikbol T. TOREBEK [2]

[1] *South Ural State University, Chelyabinsk, Russia*
*E-mail: valerykarachik@mail.ru*
[2] *Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan*
*E-mail: torebek@math.kz*

**Abstract:** In this paper a generalized third boundary value problem for the biharmonic equation:

$$\Delta^2 u = f(x), \quad x \in S,$$

in the unit ball $S = \{x \in \mathbb{R}^n : |x| < 1\}$ with boundary operators up to third order containing normal derivatives: $\frac{\partial}{\partial \nu}$ and Laplacian: $\Delta$

$$\begin{cases} a_{00}u + a_{01}\frac{\partial}{\partial \nu}u + a_{02}\Delta u\Big|_{\partial S} = \varphi_1(s), \quad s \in \partial S, \\ \\ a_{11}\frac{\partial}{\partial \nu}u + a_{12}\Delta u + a_{13}\frac{\partial}{\partial \nu}\Delta u\Big|_{\partial S} = \varphi_2(s), \quad s \in \partial S, \end{cases}$$

is investigated.

The existence and uniqueness theorems are proved.

Some particular cases of general problem are considered.

**Keywords:** biharmonic equation, boundary value problems, Laplace operator, harmonic polynomials

**2010 Mathematics Subject Classification:** 35J05, 35J25, 26A33.

## On Generalized Quasi-Convex Bounded Sequences

Mahmut KARAKUŞ [1]

[1] *Department of Mathematics, Yüzüncü Yıl University, Van, TURKEY*
*E-mail: matfonks@gmail.com*

**Abstract:** The space of all sequences $a = (a_k)$ for which $||a||_q = \sum_k k|\Delta^2 a_k| + \sup_k |a_k| < \infty$ is denoted by $q$ [2]. Here, $\Delta a_k = a_k - a_{k+1}$ and $\Delta^m a_k = \Delta(\Delta^{m-1}a_k) = \Delta^{m-1}a_k - \Delta^{m-1}a_{k+1}$ with $\Delta^0 a_k = a_k$, $m \geq 1$. If $a = (a_k) \in q$ then $k\Delta a_k \to 0 \ \ (k \to \infty)$ and $q \subset bv$, the space of all sequences of bounded-variation, since $\sum_k |\Delta a_k| \leq \sum_k k|\Delta^2 a_k|$. In the study we shall give a generalization of quasi-convex bounded sequences.

**Keywords:** $FK$ spaces, $\beta-$ and $\gamma-$ duals, topological sequence spaces.

**2010 Mathematics Subject Classification:** 46A45, 40G05.

**References:**

[1] Bosanquet J., Note on convergence and summability factors, *J. London Math. Soc.* 20, pp. 39-48, 1945.

[2] Buntinas M., Convergent and bounded Cesro sections in FK-spaces, *Math. Zeitschr.*121 , pp. 191-200, 1971.

[3] Dağadur I., $C_\lambda$ -semiconservative FK spaces, *Ukrainian Math. J.* 64(7): pp. 1036-1039, 2012.

[4] Wilansky A., *Summability Through Functional Analysis*, North-Holland, Amsterdam, 1984.

---

## On $AK(S)$ and $AB(S)$ Properties of a $K$-Space

Mahmut KARAKUŞ [1]

[1] *Department of Mathematics, Yüzüncü Yıl University, Van, TURKEY*
*E-mail: matfonks@gmail.com*

**Abstract:** A typical sum $s$ on a K- space $S$ has often the representation,

$$s(z) = \lim_{\gamma\in\Gamma} \sum_k u_{\gamma k} z_k, \ z = (z_k) \in S \tag{6}$$

where $\Gamma$ is a directed index set and $u_\gamma = (u_{\gamma k}) \in \phi$, the space of finitely non-zero sequences, for each $\gamma \in \Gamma$. Let a K-space $S$ be equipped with a sum (6). Then, for each $x = (x_k)$ and $\gamma \in \Gamma$, the sequence $P_\gamma(x) = \sum_k u_{\gamma k} x_k \delta^k, \ \ (\gamma \in \Gamma)$ is called the $\gamma$th $S$-section of $x$ [2]. Here $\delta^k$ is the sequence whose $k$th component is 1 all the others are 0.

If $\lambda \supset \phi$ is a $K$ space, then Boos and Leiger defined the spaces $\lambda_{AB(S)}$ and $\lambda_{AK(S)}$ in [2] as

$$\lambda_{AB(S)} = \{x \in \omega | (P_\gamma(x))_{\gamma\in\Gamma} \text{ is a bounded net in } \lambda\},$$

and

$$\lambda_{AK(S)} = \{x \in (\lambda_{AB(S)} \cap \lambda) | \lim_\gamma P_\gamma(x) \text{ exists in } \lambda\}.$$

In this work, we investigate some properties of these spaces and give some theorems related to the duals.

**Keywords:** K- spaces, $n$-th section of a sequence, $\beta$-, $\gamma$-, $f$- duality.

**2010 Mathematics Subject Classification:** Primary: 40H05, 46A45, Secondary: 40G99, 40C05.

**References:**

[1] Boos J., Leiger T., Dual pairs of sequence spaces, *Int. J. Math. Math. Sci.*, Vol. 28, pp. 9-23, 2001.

[2] Boos J., Leiger T., Dual pairs of sequence spaces II, *Proc. Estonian Acad. Sci. Phys. Math.*, Vol.51, pp. 3-17, 2002.

[3] Boos J., Leiger T., Dual pairs of sequence spaces III, *J. Math. Anal. Appl.*, Vol.324, pp. 1213-1227, 2006.

[4] Boos J., Classical and Modern Methods in Summability, *Oxford University Press. New York, Oxford,* 2000.

[5] Buntinas M., Convergent and bounded CesÃ ro sections in FK-spaces, *Math. Zeitschr.*, Vol.121, pp. 191-200, 1971.

[6] Buntinas M., On Toeplitz sections in sequence spaces, *Math. Proc. Camb. Phil. Soc.*, Vol.78, pp. 451-460, 1975.

[7] Ruckle W. H., An abstract concept of the sum of a numerical series, *Can. J. Math.*, Vol.22, pp. 863-874, 1970.

---

## On $q^\lambda$ and $q_0^\lambda$ Invariant Sequence Spaces

Mahmut KARAKUŞ [1]

[1] *Department of Mathematics, Yüzüncü Yıl University, Van, TURKEY*
*E-mail: matfonks@gmail.com*

**Abstract:** Invariant sequence spaces are very helpful for investigations of the duality of sequence spaces. For instance, if the sequence space $X$ satisfies the condition $\ell_\infty.X = X$ then its $\alpha-$, $\beta$- and $\gamma$- duals are same [4]. Garling [1] investigated $B$- and $B_0$- invariant sequence spaces and Buntinas [2] introduced and investigated $q$- and $q_0$- invariant sequence spaces and recently, Grosse- Erdmann [3] studied on $\ell_1$ invariant sequence spaces.

In this work, we define $q^\lambda$ and $q_0^\lambda$ invariant sequence spaces, $X$ with $q^\lambda.X = X$ and $q_0^\lambda.X = X$, respectively. and give some related theorems.

**Keywords:** K- spaces, $\lambda$-boundedness and $\lambda$-convergence of a sequence, $\beta$-, $\gamma$-, $f$- duality.

**2010 Mathematics Subject Classification:** Primary: 40H05, 46A45, Secondary: 40G99, 40C05.

**References:**

[1] Garling D. J. H., *On topological sequence spaces*, Proc. Camb. Phil. Soc., Vol.63, pp. 997-1019, 1967.

[2] Buntinas M., *Convergent and bounded CesÃ ro sections in FK-spaces*, Math. Zeitschr., Vol.121, pp. 191-200, 1971.

[3] Grosse-Erdmann K.G., *On $\ell_1$ invariant sequence spaces*, J. Math. Anal. Appl., Vol.262, pp. 112-132, 2001.

[4] Boos J., Classical and Modern Methods in Summability, *Oxford University Press. New York, Oxford,* 2000.

---

## Solution of a Cauchy problem for iterated generalized two-axially symmetric equation of hyperbolic type

Shakhobiddin T. KARIMOV [1]

[1] *Faculty of Mathematics and Physics, Ferghana State University, Ferghana-150100, Republic of Uzbekistan*
*E-mail: shkarimov09@rambler.ru*

**Abstract:** In the work an explicit solution of Cauchy problem for iterated generalized two-axially symmetric equations of hyperbolic type was constructed.

In this work, we considered an analogue of the Cauchy problem in a domain $\Omega = \{(x, y) : 0 < y < x\}$ finding a classical solution $u(x, y) \in C^{2m-1}(\bar{\Omega})$ of the equation

$$[h^{\lambda}_{\beta,\alpha}]^m(u) \equiv \left( \frac{\partial^2}{\partial y^2} - \frac{\partial^2}{\partial x^2} - \frac{2\alpha}{x}\frac{\partial}{\partial x} + \frac{2\beta}{y}\frac{\partial}{\partial y} + \lambda^2 \right)^m u(x, y) = 0, \tag{1}$$

satisfying the initial conditions

$$\left.\frac{\partial^{2k} u}{\partial y^{2k}}\right|_{y=0} = \varphi_k(x),\ x \geqslant 0; \quad \left.\frac{\partial^{2k+1} u}{\partial y^{2k+1}}\right|_{y=0} = 0,\ x \geqslant 0,\ k = \overline{0, m-1}; \tag{2}$$

where $\alpha, \beta, \lambda \in R,\ m \in N$, and $0 < \alpha < 1,\ 0 < \beta < 1/2$, $\varphi_k(x)$, $(k = \overline{0, m-1})$ are given smooth functions.

By applying two-dimensional generalized Erdelyi-Kober operator [1], the explicit formula of a solution of the problem {(1), (2)} was received

$$u(x, y) = \sum_{n=0}^{m-1} \frac{2^{-2n} y^{1-2\beta}}{n!\Gamma(\beta + n)} \int_{x-y}^{x+y} f_n(s) \left(\frac{s}{x}\right)^{\alpha} \frac{\Xi_2(\alpha, 1-\alpha; \beta+n; \sigma, \omega)}{[y^2 - (x-s)^2]^{1-\beta-n}} ds,$$

where $f_n(x) = \sum\limits_{j=0}^{n} (-1)^j C_n^j [B^x_{\alpha-1/2}]^j \tilde{\varphi}_{n-j}(x)$, $\tilde{\varphi}_n(x) = \sum\limits_{j=0}^{n} \gamma_n C_n^j \lambda^{2(n-j)} \varphi_j(x)$, $\gamma_n = \Gamma((2n+1)/2+\beta)/\Gamma((2n+1)/2)$, $\sigma = [y^2-(s-x)^2]/(4xs)$, $\omega = -(\lambda^2/4)[y^2-(s-x)^2]$, $\Xi_2(\alpha, 1-\alpha; \beta+n; \sigma, \omega)$ is Humbert hypergeometric function.

**Keywords:** Erdélyi-Kober operator, two-axially symmetric equations

**2010 Mathematics Subject Classification:** 26A33, 35L10

**References:**

[1] Karimov Sh.T., "Multidimensional generalized Erdélyi-Kober operator and its application to solving Cauchy problems for differential equations with singular coefficients", // Fract. Calc. Appl. Anal., Vol. 18, No 4, pp. 845-86, 2015.

## On the sufficient condition for embedding of generalized Lizorkin-Triebel spaces

Gulden KARSHYGINA[1], Alua IGENBERLINA[1], Zauresh SULEYMENOVA[1]

[1] *Department of Mechanics and Mathematics, Eurasian National University, Astana, Kazakhstan*

*E-mail: karshygina84@mail.ru, igenberlina@mail.ru, zr-s2012@yandex.kz*

**Abstract:** In this paper the sufficient conditions for the embedding of generalized Lizorkin–Triebel spaces $F^{\varphi}_{p\theta}$ are given.

Denote by U the set of all continuous, increasing on $[0, 2\pi]$ functions such that $\varphi(0) = 0$ and for some $\alpha > 0$ $\varphi(t)t^{-\alpha}$ almost increases. Let $1 \le p, \theta \le \infty$ and $\varphi \in U$. We say that a function $f$ belongs to generalized Lizorkin–Triebel spaces $F^{\varphi}_{p\theta}$ if $f \in L^p(R)$ and

$$\|f\|_{F^{\varphi}_{p,\theta}} = \|f\|_p + \left\| \left\{ \sum_{k=0}^{\infty} \left[ \varphi^{-1}(2^{-k}) \left| S_{2^{k+1}} f - S_{2^k} f \right| \right]^{\theta} \right\}^{\frac{1}{\theta}} \right\|_{L_p} < \infty,$$

where $S_k f(x)$ partial sum Fourier series. If $\varphi(x) = x^r$, $r > 0$ this space coincides with the space $F^r_{p\theta}$ [1].

**Theorem 2.** *Let* $1 \le p \le p' \le \infty$, $1 \le \theta, \theta' < \infty$. *Let* $\varphi, \psi \in U$.

1) *for* $\theta' < \theta$ *if* $\sum_{s=0}^{\infty} \left[ \varphi(2^{-s}) \psi^{-1}(2^{-s}) 2^{\frac{1}{p'}} \right]^{\frac{\theta'\theta}{\theta - \theta'}} < \infty$,

2) *for* $\theta' \ge \theta$ *if* $\sup\limits_{0<t\le 1} \left[ \varphi(t)\psi(t)^{-1} t^{\kappa} \right] < \infty$,

*where* $\kappa = \frac{1}{p} - \frac{1}{p'}$ . *Then the embedding* $F^{\varphi}_{p\theta} \hookrightarrow F^{\psi}_{p'\theta'}$ *holds.*

In the case of generalized Nikol'skii – Besov spaces similar embedding was considered in [2].

**Keywords:** Lizorkin-Triebel spaces, embedding

**References:**

[1] Triebel H., Theory of function spaces , 450 p., 1983.
[2] Goldman M. L., "On embedding for generalized Holder classes , *Math. notes*, Vol.12, No.3, pp. 325-336, 1972.

# About one special boundary value problem for multidimensional parabolic integro-differential equation

Ermek KHAIRULLIN [1]

[1] *Department of Mathematics, Kazah National Research Technical University, Almaty, Kazakhstan*
*E-mail: khairullin_42_42@mail.ru*

**Abstract:** In this paper we consider a special boundary value problem for multidimensional parabolic integro-differential equation with boundary conditions that contains as a boundary condition containing derivatives of order higher than the order of the equation [1,2,4]. The solution is sought in the form of a thermal potential of a double layer. Shows lemma of finding the limits of the derivatives of the unknown function in the neighborhood of the hyperplane. Using the boundary condition and lemma obtained integral-differential equation (IDE) of parabolic operators, whre an unknown function under the integral contains higher-order space variables derivatives. IDE is reduced to a singular integral equation (SIE), when an unknown function in the spatial variables satisfies the Holder.The characteristic part is solved in the class of distribution function using method of transformation of Fourier-Laplace. Found an algebraic condition for the transition to the classical generalized solution. Integral equation of the resolvent for the characteristic part of SIE is obtained. Integro-differential equation is reduced to the Volterra-Fredholm type integral equation of the second kind by method of regularization. It is shown that the solution of SIE is a solution of IDE. Obtain a theorem on the solvability of the boundary value problem of multidimensional parabolic integro-differential equation, when a known function of the spatial variables belongs to the Holder class and satisfies the solvability conditions.

**Keywords:** Multidimensional parabolic integro-differential equation, singular integral equation, regularization

**References:**

[1] Tihonov A.N. O kraevyh uslovijah, soderzhashhih proizvodnye porjadka, prevyshajushhego porjadok uravnenija. *Matem. sbornik.* 1950, V.26(68), N 1, S. 35-56(in Russ).

[2] Kim E.I. Ob usloviyah razreshimosti odnoj granichnoj zadachi uravnenija teploprovodnosti, *Dokl. AN SSSR.*1961, V.140, N.3, S.553-556 (in Russ).

[3] Khairullin E.M. Reshenie odnogo klassa mnogomernyh singulyarnyh integral ' nyh uravnenij, *Vestnik KazNTU* - Almaty, 2010. - N. 5 ( 81 ) - S.236-240(in Russ).

[4] Khairullin E.M., Tulesheva G.A., Lukpanova L.H. "O kraevyh uslovijah dlja parabolicheskogo uravnenija v poluprostranstve, prevyshajuschih porjadok uravnenija", *Trudy II Mezhdunarodnoj nauchno-prakticheskoj konferentsii Informatsionno-innovatsionnyje tehnologii: integratsija nauki, obrazovanija i biznesa,* T.II. Almaty, 2011. S.433-437(in Russ).

[5] Khairullin E.M., Khalbaeva ZH.A./ Reshenie tretej kraevoj zadachi dlya uravneniya teploprovodnosti s podvizhnymi granicami, *Trudy Rol' i mesto molodyh uchenyh v realizacii novoj ehkonomicheskoj politiki Kazahstana mezhdunarodnyh Satpaevskih chtenij.* Almaty, 2015. - TV 4. - S. 450-456(in Russ).

## About solvability of some boundary value problems for Poisson equation in the ball

M.KOSHANOVA [1], K.I.USMANOV [2], B.TURMETOV [3]

[1,2,3] *Akhmet Yasawi University, 29 Sattarkhanov street, 161200 Turkestan, Kazakhstan*
*E-mail: maira_koshanova@mail,y_kairat@mail.ru*
[3] *Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: turmetovbh@mail.ru*

**Abstract:** In the paper we study properties of some integral - differential operators of fractional order. As an application of the properties of these operators for Poisson equation we examine questions on solvability of a fractional analogue of Neumann problem and analogues of periodic boundary value problems for circular domains. The exact conditions for solvability of these problems are found

**Keywords:** fractional derivative, Hadamard operator, Poisson equation, Neumann problem, periodic problem

**2010 Mathematics Subject Classification:** 35J15, 35J25

---

## On weak solutions of the Dirichlet problem for the Tricomi equation

Vladimir V. KOVRIZHKIN

*Omsk State University, Russia*
*E-mail: rudin09@rambler.ru*

**Abstract:** Let $Tu \equiv yu_{xx} + u_{yy}$ — formal differential operator. $\Delta$ — curvilinear triangle bounded by segment $AB$ of $x$-axis, arc $BC$ of the characteristic $x_B = x + \frac{2}{3}|y|^{3/2}$ and smooth curve $AC$ which have not characteristic directions. Also $\Delta$ is convex with respect to characteristics of $T$.

Consider operator $\Lambda'$ defined on functions $u \in C^2(\overline{\Delta})$, $u|_{AB \cup AC} = 0$ by equality: $\Lambda' u = Tu$. Let $\Lambda$ is closure of $\Lambda'$ in $L_2(\Delta)$. Weak solution of the Dirichlet problem for the Tricomi equation $Tv = g$, $g \in L_2$ is $v \in L_2$ such that $\langle v, \Lambda u \rangle = \langle g, u \rangle \ \forall u \in D(\Lambda)$. From the definition follow that $v$ is the weak solution of the Dirichlet problem for the Tricomi equation $Tv = g$ if and only if $v \in D(\Lambda^*)$.

**Theorem.** Any weak solution of the Dirichlet problem belongs to $W_2^1(\Delta \backslash O_A)$ where $O_A$ any neighbourhood of point $A$. Moreover there is orthogonal decomposition of $L_2(\Delta)$ to the sum $H_1 \oplus H_2$ such that for $g \in H_1$ corresponding $v \in W_2^1(\Delta)$ and for $g \in H_2$ $v \notin W_2^1(\Delta)$.

**Keywords:** Dirichlet boundary problem, Tricomi equation

**2010 Mathematics Subject Classification:** 94B05, 94B15

**References:**

[1] Kovrizhkin V. V., "On weak solutions of the Darboux problem", *Diff. Equations*, Vol. 8, pp. 68–75, 1972.

## Global solvability in some models related to equations of Sobolev type

Aleksandr KOZHANOV

*Sobolev Institute of Mathematics,*
*Siberian Branch of the Russian Academy of Sciences, Novosibirsk, Russia*
*E-mail: kozhanov@math.nsc.ru*

**Abstract:** In the report there are results on global solvability of initial-boundary value problems for

— higher-order Sobolev-type strongly nonlinear equations (pseudoparabolic equations) appearing in viscoelasticity;

— nonlinear analogs of the Boussinesq equations of long waves appearing in modeling a line of transmissions containing nonlinear elements (for example, diodes, capacity of which is a nonlinear function of voltage);

— linear and nonlinear integro-differential equations of viscoelasticity containing degenerate equations.

Some properties of solutions are also additionally studied.

**Keywords:** Global solvability, pseudoparabolic equation, equations of Sobolev type, Boussinesq equation, integro-differential equations.

**2010 Mathematics Subject Classification:** 35S05, 35K55

**References:**

[1] Amirov Sh., Kozhanov A.I., "Global Solvability of Initial Boundary-Value Problems for Nonlinear Analogs of the Boussinesq Equation", *Mat. Zametki*, Vol. 99, No. 2, pp. 171–180, 2016.

[2] Kozhanov A.I., "Inverse problems for determining boundary regimes for some equations of Sobolev type", *Vestnik YuUrGU. Ser. Mat. Model. Progr.* , Vol. 9, No. 2, pp. 37-45, 2016.

## Oscillation and nonoscillation of half-linear second order differential equation with alternating potential

Leili KUSSAINOVA[1], Bakhytty KOSHKAROVA[2]

*L.N. Gumilyov Eurasian national university, Kazakhstan*
*E-mail:* [1] *leili2006@mail.ru,* [2] *b-koshkarova@yandex.kz*

**Abstract:** In this work we study the questions of oscillation and nonoscillation of the following half–linear second order differential equation

$$(HL) \qquad \left(\rho(t)\left|y'(t)\right|^{p-2}y'(t)\right)' + q(t)\left|y(t)\right|^{p-2}y(t) = 0, \;\; t \geq 0,$$

on the interval $I = (a, \infty)$, $-\infty \leq a < \infty$, when $1 < p < \infty$, $p \neq 2$. Here $\rho(t) > 0$ and $q(t)$ continuous functions on $I$.

The results of investigations of the properties and methods for the equation (HL) up to year 2005 are exposed in the book by Došlý and Řehák [1].

For researching properties of equation (HL) we use the variational method, and we establish the conditions of oscillation and nonoscillation of the equation (HL), when the function $q(t)$ changes the sign on the interval $I$.

**Keywords:** oscillation, nonoscillation, half-linear equation, alternating potential, variational method

**2010 Mathematics Subject Classification:** 34C10

**References:**

[1] Došlý O., Řehák P., "Half–linear differential equations", *IEEE Math. Studies, North-Holland*, Vol.202, 2005.

## Soliton deformations of some minimal surfaces

Damir KURMANBAEV

*Department of Mechanics and Mathematics, al-Farabi Kazakh national university, Almaty, Kazakhstan*
*E-mail: kurmanbaev.damir@gmail.com*

**Abstract:** Modified Veselov-Novikov equation was introduced in [1]. A method for constructing exact solutions of this equation by Mutard transformations was proposed in [2] and in [3] was obtained a geometric interpretation of Mutard transformations. It is given by the solution of the Dirac equation and three real constants. Any solution of the Dirac equation defines a surface in three-dimensional Euclidean space, given up to shifts by Weierstrass representation[4]. Fixing the three constants, we completely fix the surface. On this surface are given a conformal parameter, and the potential of the Dirac operator U is the potential representation of the surface. Applying inversion to this surface with its center at the origin, we obtain a new surface with the same conformal parameter and new potential. In this paper applied this transformations for mVN equation on the examples of higher order Enneper's surface, catenoid and helicoid and constructed soliton deformations [5] of this surfaces.

**Keywords:** Dirac operator, modified Veselov-Novikov equation, Mutard transformations, Enneper's surface,catenoid, helicoid,deformation of surface, inversion.

**2010 Mathematics Subject Classification:** 35Q51, 49Q05

**References:**

[1] Bogdanov L.V., "Veselov-Novikov equation as a natural two-diensional generalization of the Korteweg-de Vries equation", *Theor. Math. Phys.*, Vol. 70, pp.309–314, 1987.

[2] Delong Yu, Q.P. Liu, and Shikun Wang, "Darboux transformation for the modified Veselov-Novikov equation",*J. of Physics A 35*, pp. 3779-3785, 2001.

[3] I.A.Taimanov, "The Moutard transformations of two-dimensional Dirac operators and Mebius geometry",*Mat.Zametki*, Vol. 97, Issue. 1, pp. 129-141, 2015.

[4] I. A. Taimanov, "Two-dimensional Dirac operator and the theory of surfaces", *Russian Math. Surveys*, Vol.61, No1, pp. 79-159, 2006.

[5] Konopelchenko B.G., "Induced surfaces and their integrable dynamics", *Stud. Appl. Math.*, Vol.96, No.1, pp. 9-51, 1996.

# Necessary and sufficient condition that function belong to the Nikolskii-Morrey space $H^{(r)}M^{\lambda}_{p\,x_1}$

Nurgul KYDYRMINA [1]

[1] *Institute of Applied Mathematics, Karaganda, Kazakhstan*
*E-mail: nurgul-k@mail.ru*

**Abstract:** In this paper there were established the necessary and sufficient conditions that the function belongs to the Nikol'skii–Morrey space $H^{(r)}M^{\lambda}_{p\,x_1}$.

**Definition 1.** *Let $0 < p \le +\infty$, $0 \le \lambda \le \frac{n}{p}$. Denote by $M^{\lambda}_{p}(\mathbb{R}^n)$ the space of all real-valued function $f$, measurable on $\mathbb{R}^n$, such that*

$$\|f\|_{M^{\lambda}_{p}(\mathbb{R}^n)} = \sup_{\substack{x\in\mathbb{R}^n \\ \rho>0}} \|\rho^{-\lambda}\|f\|_{L_p(B(x,\rho))}\|_{L_\infty(0,\infty)} < \infty.$$

**Definition 2.** *We say measurable function $f(x_1,\dots,x_n)$ belongs to the class $H^{(r)}M^{\lambda}_{p\,x_1}(B)$ $(r>0,\ 1\le p\le+\infty)$, if it satisfies the following condition.*

*Assume that $r=\rho+\alpha$, where $\rho$ is integer and $0<\alpha\le 1$. Let for almost all $x_2,\dots,x_n$ exists the partial derivative $\frac{\partial^\rho f}{\partial x_1^\rho}$ defined almost everywhere on $\mathbb{R}^n$ that belong to the Morrey space $M^{\lambda}_{p}(\mathbb{R}^n)$ and such that for all $h$*

$$\left\| \frac{\partial^\rho f(x_1+h,x_2,\dots,x_n)}{\partial x_1^\rho} + 2\frac{\partial^\rho f(x_1,\dots,x_n)}{\partial x_1^\rho} + \frac{\partial^\rho f(x_1-h,x_2,\dots,x_n)}{\partial x_1^\rho} \right\|_{M^{\lambda}_{p}(\mathbb{R}^n)} \le B|h|^\alpha.$$

**Definition 3.** *Denote by $A_{\nu\,x_1}(f;M^{\lambda}_{p}(\mathbb{R}^n))$ the best approximation of function $f\in M^{\lambda}_{p}(\mathbb{R}^n)$ by entire functions $g_\nu$ of exponential type $\nu$ in metrics of the space $M^{\lambda}_{p}(\mathbb{R}^n)$, i.e.*

$$A_{\nu\,x_1}(f;M^{\lambda}_{p}(\mathbb{R}^n)) = \inf_{g_\nu} \|f-g_\nu\|_{M^{\lambda}_{p}(\mathbb{R}^n)}.$$

**Theorem 4.** *Function $f\in M^{\lambda}_{p}(\mathbb{R}^n)$ belongs to the class $H^{(r)}M^{\lambda}_{p\,x_1}$ (with some constant $B$), if and only if there exists the constant $K$ such that*

$$A_{\nu,x_1}(f;M^{\lambda}_{p}(\mathbb{R}^n)) < \frac{K}{\nu^r},$$

*for all $\nu\ge 1$ or for all $\nu$ that run over geometric increasing progression $\nu=a^k$ $(a>1,\ \ k=0,1,\dots)$ .*

This work was supported by the Scientific Committee of the Ministry of Education and Science of the Republic of Kazakhstan, grant No. 1777/GF4.

**Keywords:** Morrey space, Nikolskii-Morrey space, best approximation, entire functions of exponential type

**2010 Mathematics Subject Classification:** 41A50, 46E35

## Cauchy problem for a nonlinear degenerate parabolic system not in divergence form

Alisher MATYAKUBOV [1]

[1] *Department of Applied Mathematics and Computer Analysis, National university of Uzbekistan, Tashkent, Uzbekistan*
*E-mail: almasa@list.ru*

**Abstract:** In this paper, we consider the following Cauchy problem to a nonlinear degenerate parabolic system not in divergence form:

$$\begin{array}{l} \frac{\partial u}{\partial t} = v^{m_1} \nabla \left( \left| \nabla u^k \right|^{p-2} \nabla u^n \right), \\ \frac{\partial v}{\partial t} = u^{m_2} \nabla \left( \left| \nabla v^k \right|^{p-2} \nabla v^n \right), \end{array} \quad t > 0, \ x \in R^N,$$

$$u|_{t=0} = u_0(x) \geq 0, \quad v|_{t=0} = v_0(x) \geq 0, \ \forall x \in R^N$$

where $p$, $m_1$, $m_2$, $k$, $n$ - are positive constants, $N \geq 1$.

This system can be used to describe the development of multiple groups in the dynamics of biological groups, where $u$, $v$ are the densities of the different groups (see [1]). In paper [1], Gao et al. considered parabolic systems not in divergence form in the case $n = 1$ and $p = 2$, with source and null Dirichlet boundary conditions. The local existence and uniqueness of classical solution are proved. Moreover, it is proved that all solutions exist globally with homogeneous Dirichlet boundary condition. Porous medium equations with local sources or with nonlocal sources subjected to nonlocal boundary conditions were studied (see [3, 4]). They discussed the conditions of existence and blow-up.

In this paper, by using the comparison principle [2] and the standard equation method [2], we will obtain a global solvability of solutions of Cauchy problem and using the method of nonlinear splitting, we can construct an asymptotic representation of the solution as $t \to \infty$. We obtain the estimates of the solution of Cauchy problem. Results of computational experiments shows, that the self - similar solutions are very appropriate approximation and the iterative method basing on the method Picard is effective for the solution of nonlinear problems and leads to the nonlinear effects if we will use as initial approximation the solutions of self similar equations constructed by the method of nonlinear splitting and by the method of standard equation.

**Keywords:** nonlinear system, non-divergence form, global solutions, asymptotic behavior of solutions

**2010 Mathematics Subject Classification:** 35B40, 35B51, 35C06

**References:**

[1] Gao Y., Meng Q., Guo Y., "Study of properties of solutions for quasilinear parabolic systems", *MATEC Web of Conferences*, Vol.61, No.1, pp. 1-4, 2016.

[2] Aripov M., "Standard equation's methods for solutions to nonlinear problems", *Tashkent: Fan*, 138 p., 1988.

[3] Zhong G., Tian L., "Blow-up for a parabolic system with nonlocal sources and nonlocal boundary conditions", *Boundary Value Problems*, Vol.2015, No.61, 2015.

[4] Chen Y., "TGlobal blow-up for a localized quasilinear parabolic system with nonlocal boundary conditions", *Appl. Anal.*, Vol.7, pp. 1495-1510, 2013.

[5] Samarskii A., Galaktionov V., Kurdyumov S., Mikhailov A., "Blow-up in quasi-linear parabolic equations", *Berlin: De Gruyter Expositions in Mathematics*, Vol.19, 538 p., 2011.

---

## On separability of a differential operator of non-classical type in an unbounded domain

Mussakan MURATBEKOV, Sabit IGISINOV, Abai SHYRAKBAEV

*Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: musahan_m@mail.ru*

**Abstract:** Review of literature on the equations of nonclassical type can be found in [1–3]. Properties of the semi periodic Dirichlet problem for degenerate equations were studied in [4–6].

In these studies we investigated the solvability and smoothness of solutions of boundary value problems for nonclassical equations without degeneration

$$Lu = -\frac{\partial^2 u}{\partial y^2} - \sum_{k=0}^{s} R_k(y)\frac{\partial^{2k+1} u}{\partial x^{2k+1}} + \sum_{k=0}^{m}(-1)^k C_k(y)\frac{\partial^{2k} u}{\partial x^{2k}}$$

defined on a set $C_o^\infty(\Omega)$, where $\Omega = \{(x,y) : -\infty < x < +\infty, 0 < y < 1\}$, $C_o^\infty(\Omega)$ is the set of infinitely differentiable functions satisfying the condition:

$$u(x,0) = u(x,1) = 0,$$

and compactly supported with respect to $x$.

The conditions existence of a solution and the separability of the operator were found during the study.

**Keywords:** non-classical type operator, resolvent, separability, an unbounded domain

**2010 Mathematics Subject Classification:** 02.30.Jr, 02.60.Lj

**References:**

[1] Dubinskii, U.A., "On some differential-operator equations of arbitrary order, *Math.Sbornik*, Vol. 90, pp.1–22, 1973.

[2] Romanko, V.K., "Solvability of boundary value problems for operator differential equations of high order, *Differential Equations*, Vol. 14, pp.1081–1092, 1978.

[3] Yurchuk, N.I., "On boundary problems for equations with operators of the form $\frac{d^{2m+1}}{dt^{2m+1}} + A$, *Differential Equations*, Vol. 10, pp.950–952, 1974.

[4] Kal'menov, T.Sh., Otelbaev, M., "On the smoothness of solutions of a class of degenerate elliptic equations, *Differential Equations*, Vol. 13, pp.1244–1255, 1977.

[5] Muratbekov,M.B., "Coercive estimates for a higher order differential operator, *Differential Equations*, Vol. 17, pp.893–901, 1981.

[6] Muratbekov,M.B., Ospanov, K.N., Muratbekov, M.M., "Coercive solvability of odd-order differential equations and its application, *Doklady Mathematics*, Vol. 82, pp.1–3, 2010.

## On spectral properties of a Shrödinger operator with a negative parameter

Mussakan MURATBEKOV, Madi Muratbekov

*Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: musahan_m@mail.ru*

**Abstract:** In the paper a Shrödinger operator with a negative parameter

$$-\Delta + (-t^2 + ita(x) + c(x)) \tag{1}$$

is studied in the space $L_2(\mathbb{R}^n)$, where $t$ is a parameter and $-\infty < t < \infty, i^2 = -1$. $\mathbb{R}^n$ is an n-dimensional Euclidean space $x \in \mathbb{R}^n$ $(x = (x_1, x_2, ..., x_n))$.

It is well known that when $t = 0$ the spectral properties of the Schrödinger operator $\Delta + c(x)$ are strongly dependent on the behavior of $c(x)$ on infinity. In this case, the spectral characteristics of the Schrödinger operator are well studied.

The discreteness of the spectrum and the estimates of approximate numbers (s-numbers) of a Schrödinger operator

$$-\Delta + q_1(x) + iq_2(x), (q_1(x) \geq 0, q_2(x) \geq 0)$$

have been studied.

In this paper such questions as:

a) the existence of the resolvent; b) the discreteness of spectrum; c) the estimates of the distribution function of singular values ($s$-values) will be studied for the operator (1).

Review of the literature shows that these questions for the Schrödinger operator with a negative parameter are insufficiently studied.

**Keywords:** resolvent, spectrum, singular values, Shrödinger operator

**2010 Mathematics Subject Classification:** 02.30.Jr, 02.30.Sa

## On solvability of a class of nonlinear two-point boundary value problems

Ergashboy MUKHAMADIEV, Alizhon NAIMOV

*Department of information systems and technologies,*
*Vologda State University, Russia*
*E-mail: emuhamadiev@rambler.ru, nan67@rambler.ru*

**Abstract:** We study the solvability of a class of nonlinear two-point boundary value problems for systems of ordinary second-order differential equations on the plane. On the basis of properties of the leading nonlinear terms, we prove a criterion for the solvability of boundary value problems under arbitrary perturbations in a given set by using methods for the computation of the winding number of vector fields.

In this work, we study the solvability of nonlinear two-point boundary value problems of the form

$$z''(t) = \overline{z'(t)}^m + f(t, z(t), z'(t)), \quad z = x + iy \in \mathbf{C}, \quad 0 < t < 1, \tag{1}$$

$$z'(0) = A_0(z(0)) + h_0(z), \quad z'(1) = A_1(z(0)) + h_1(z). \tag{2}$$

Here $m$ is an integer larger than unity, $\mathbf{C}$ is the complex plane, $\overline{z} = x - iy$, and the mappings $f : [0,1] \times \mathbf{C}^2 \mapsto \mathbf{C}$ and $A_0, A_1 : \mathbf{C} \mapsto \mathbf{C}$ are continuous and satisfy the conditions:

$$A_j(\lambda z) \equiv \lambda A_j(z) \quad \text{for any } \lambda \geq 0, j = 0, 1, \tag{3}$$

$$\max_{0 \leq t \leq 1} |f(t, z, w|(|z| + |w|)^{-m} \to 0 \quad \text{as } |z| + |w| \to \infty. \tag{4}$$

The mappings $h_0$ and $h_1$ act continuously from $C^1([0,1]; \mathbf{C})$ into $\mathbf{C}$, where $C^1([0,1]; \mathbf{C})$ is the space of complex-valued functions continuously differentiable on the interval $[0,1]$, and satisfy the conditions

$$|h_j(z)|/||z||_{C^1} \to 0 \quad \text{as } ||z||_{C^1} \to \infty, j = 0, 1. \tag{5}$$

For this class of boundary value problems, we study the solvability on the basis of an a priori estimate of solutions with the use of methods for the computation of the winding number of vector fields [1]. This problem was earlier considered in [2, 3, 4] and is of interest from the viewpoint of the application of methods of nonlinear analysis and the development of investigation methods for nonlinear two-point boundary value problems.

The research was supported in part by the Russian Foundation for Basic Research (projects no. 15-01-04713a, 16-01-00150a).

**Keywords:** nonlinear two-point boundary value problem, the winding number of vector fields.

**2010 Mathematics Subject Classification:** 34B15

**References:**

[1] Krasnosel'skii, M.A. and Zabreiko, P.P., Geometricheskie metody nelineinogo analiza (Geometric Methods of Nonlinear Analysis), Moscow: Nauka, 1975.

[2] Naimov, A.N., A Remark on the Theory of Two-Point Boundary Value Problems for Systems of Ordinary Diferential Equations, Dokl. Akad. Nauk Resp. Tadjikistan, 1998, vol. 41, no. 9, pp. 30—34.

[3] Naimov, A.N., On the Computation of the Winding Number of a Completely Continuous Vector Field, Dokl. Akad. Nauk Resp. Tadjikistan, 1998, vol. 41, no. 10, pp. 56—61.
[4] Naimov, A.N., On the Solvability of the Third Nonlinear Two-Point Boundary Value Problem on the Plane, Difer. Uravn., 2002, vol. 38, no. 1, pp. 132—133.

---

## Convergence of kernel density estimators on compact sets

Kairat MYNBAEV [1,2,3]

[1] *International School of Economics, Kazakh-British Technical University, Almaty, Kazakhstan*
*E-mail: kairat_mynbayev@yahoo.com*
[2] *Institute of Mathematics and Mathematical Modelling, Almaty, Kazakhstan*
[3] *Kazakh National University, Almaty, Kazakhstan*

**Abstract:** In many statistical applications, one has to estimate the rate of convergence in probability of a kernel density estimator. The usual estimate on compact sets is of order $\left(\frac{\log n}{nh}\right)^{1/2}$ where $n$ is the sample size, $h > 0$ is a bandwidth that depends on $n$ and $h \to 0$, $nh \to \infty$ [1]. We improve this estimate by replacing $\log n$ with a function of $n$ which grows slower than $\log n$. For example, one can take the $k$th iterated log, for any natural $k$. The idea is illustrated in estimating a density in the one-dimensional case and can be used in many other setups.

Let $K$ be a kernel, that is a function on $R$ satisfying $\int_R K(t)dt = 1$. By $X_1, ..., X_n$ we denote independent observations from a density $f$. A kernel density estimate of $f(x)$ is defined by $\hat{f}(x) = \frac{1}{nh}\sum_{i=1}^{n} K\left(\frac{x-X_i}{h}\right)$.

**Theorem 1.** Let the kernel $K$ satisfy the Lipschitz condition on $R$, be bounded and let $\int_R K^2(t)(1+|t|)dt < \infty$. Assume that the density $f$ also satisfies the Lipschitz condition on $R$ and is bounded. Fix a compact $G \subset R$ and let $f$ be positive on $G$. Suppose a function $\psi(n)$ is such that $\psi(n) \to \infty$ as $n \to \infty$ and $\limsup_{n\to\infty} \frac{n^2}{\psi(n)} < \infty$, $\liminf \frac{nh}{\psi(n)} > 0$. Then

$$\sup_{x\in G}\left|\hat{f}(x) - E\hat{f}(x)\right| = O_p\left(\left(\frac{\psi(n)}{nh}\right)^{1/2}\right).$$

**Keywords:** kernel density estimator, convergence in probability.

**2010 Mathematics Subject Classification:** 62F12, 62G07, 62G20

**References:**

[1] Martins-Filho C., Yao F., Torero M. "High-Order Conditional Quantile Estimation Based on Nonparametric Models of Regression", *Econometric Reviews*, Vol.34, pp. 907-958, 2015.

# Existence and nonexistence of positive solutions for singular $n$th-order three-point nonhomogeneous boundary value problem

Mostepha NACERI [1], Amir ELHAFFAF [2]

[1] *Economics,Commercial and Management Sciences,Preparatory School of Oran-Algeria*
*E-mail: nacerimostepha@yahoo.fr*
[2] *Mathematics Faculty Of Science,Oran University,BP1524,Es-Senia Algeria*
*E-mail: elhaffaf1@yahoo.com*

**Abstract:** In this article, we consider the boundary value problem $u^{(n)}(t) + f(t, u(t)) = 0$, $0 < t < 1$, subject to the boundary conditions $u(0) = u'(0) = 0, ....., u^{(n-3)}(0) = u^{(n-2)}(0) = 0$ and $u^{(n-2)}(1) - \alpha u^{(n-2)}(\eta) = \lambda$. In the setting, $0 < \eta < 1$ and $\alpha \in [0, \frac{1}{\eta})$ are constants and $\lambda \in [0, +\infty)$ is parameter. By placing certain restrictions on the nonlinear term $f$, we prof the existence and nonexistence of at least one positive solution to the boundary value problem with the use of the Krasnosel'skii fixed point theorem. The novelty in our setting lies in the fact that $f(t, u)$ may be singular at $t = 0$ and $t = 1$. We conclude with examples illustrating our results obtained in this paper.

**References:**

[1] Agarwal R.P., Oregan D., and Wong P.J.Y., *Positive solutions of differential difference and integral equations,* Kluwer Academic Publishers, Dordercht, 1999.

[2] Agarwal R.P., *On fourth-order boundary value problems arising in beam analysis*, Differ. Integral. Equ. 2, pp. 91-110, 1989.

[3] Bai Z., Wang H., *Positive solutions of some nonlinear fourth-order beam equations*, J. Math. Anal. Appl. 270 p., 357-368, 2002.

[4] Bai Z., *Existence of solution for some third-order boundary value problems*, Elect J.Diff.Eq, Vol.25, pp. 1-6, 2008.

[5] Baxley J.V., Haywood L., *Multiple positive solutions of nonlinear boundary value problems dynamics of continuous discrete and impulsive systems*, Dynam. Cont. Discrete. Impuls, Systems Ser.A, Vol.10, pp. 157-168, 2003.

[6] Davis A.R., Karageorghis A., and Phillips T.N., *Spectral Galerkin methods for the primary two-point boundary value problem in modelling viscoelastic flows*, Int. J.Names Engng, Vol.26, pp. 647-662, 1988.

[7] Deimling K., *Nonlinear functional analysis*, Springer New York, 1985.

[8] El-Shahed M., and Al-Mezel S., *Positive solutions for boundary value problem of fifth-order differential equations*, Inter. Math. Forum, 4, Vol.33, pp. 1635-1640, 2009.

[9] Graef J.R., Qian C., and Yang B., *A three point boundary value problem for nonlinear fourth-order differential equation*, J. Math. Anal. Appl, Vol.287, pp. 217-233, 2003.

[10] Guo L., Sun J., and Zhao Y., *Existence of positive solution for nonlinear third-order three point boundary value problem*, Nonlinear Anal, Vol.68, pp. 3151-3158, 2008.

[11] Gu D., and Lakshmikantham V., *Nonlinear problems in abstract cones*, Academic Press, San Diego, 1988.

[12] Guillopé G., Saut J. C., *Existence results for the flow of Viscoelastic fluids with a differential constitutive law*, Nonlinear Anal, Vol.15 (9), pp. 849-869, 1990.

[13] Ji C., O'Regan D., Yan B., and Agarwal R. P., *Non existence and existence of positive solutions for second-Order singular three point boundary value problems with derivate dependent and sign-changing nonlinearities*, J.Appl. Math. Comput., Vol.36, pp. 61-87, 2011.

[14] Krasnosel'skii M. A., *Positive solutions of operators equations*, Noordhoff, Groningen, 1964.

[15] Kosmatov N., *Countably many solutions of a fourth-order boundary value problem*, Elect.J. Qualit. Theory. Diff. Equ, Vol.12, pp. 1-15, 2004.

[16] Odda S.N., *Positive solutions for nth-order differential equation under some condition,* Appl. Math. Vol.6., pp. 232-239, 2011.

[17] O'Regan D., *Solvability of some fourth(and higher) order singular boundary value problems*, J. Math. Anal. Appl , Vol.161, pp. 78-116, 1991.

[18] Sun Y., *Positive solutions for third-order three point nonhomogeneous boundary value problems*, Appl. Math. lett. Vol.22, pp. 45-51, 2009.

[19] Sun Y., Liu L., Zhang J., and Agarwal R. P., *Positive solutions of singular three-point boundary value problems for second-order differential equations*, J.Comput. Appl. Math., Vol.230, pp. 738-750, 2009.

[20] Sun Y., Zhu C., *Existence of positive solution for singular fourth-order three-point boundary value problems*, Advances in Difference; Equations., Vol.51, pp. 1-6, 2013. doi: 10.1186/1687-1847-2013-51.

[21] Yang B., *Positive solutions for a fourth-order boundary value problem*, Elect. J. Qualit. Theory. Diff. Equ. 3, pp. 1-17, 2005.

---

## Riesz Lemma and Bernstein Polynomials deformations of some minimal surfaces

Igor NOVIKOV

*Voronezh, Russia*
*E-mail: igor.nvkv@gmail.com*

**Abstract:** The talk focuses on the possibility to organize the taking of non-negative trigonometric roots of polynomials converging to a piecewise-analytic function to save convergence. The most interesting for the author is the case when the trigonometric polynomials are determined using Bernstein polynomials approximating a piecewise linear function. This issue is related to the construction of compactly supported wavelets that preserves the localization with increasing smoothness.

## Summability of the Fourier coefficients of functions from anisotropic Lorentz space

Erlan NURSULTANOV [1], Nazerke TLEUKHANOVA [2]

[1] *L.N. Gumilyov Eurasian National University, Kazakhstan, and Moscow State University, Kazakh Branch, Kazakhstan*
*E-mail: er-nurs@yandex.ru*
[2] *L.N. Gumilyov Eurasian National University*
*E-mail: tleukhanova@rambler.ru*

**Abstract:** Let $1 < p < \infty$ and $\Phi = \{\phi\}_{k=1}^{\infty}$ be the orthonormal system bounded in total, $f \in L_p[0,1], f \overset{\text{a.e.}}{=} \sum_{k=1}^{\infty} a_k \phi_k(x)$. In a case of $1 < p < 2$ the task about properties of summability of Fourier coefficients $\{a_k\}_{k=0}^{\infty}$ can be solved by classical Hardy-Littlewood-Polya inequality [1]

$$\sum_{k=1}^{\infty} k^{p-2} |a_k|^p \le c \|f\|_{L_p}^p. \tag{6}$$

For the case when orthonormal system $\Phi = \{\phi\}_{k=1}^{\infty}$ is regular and $2 \le p < \infty$, according to inequality, proved by E.D. Nursultanov ( [2]), it follows

$$\sum_{k=1}^{\infty} k^{p-2} |\bar{a_k}|^p \le c \|f\|_{L_p}^p, \tag{7}$$

here $\bar{a_k} = \frac{1}{k} \left| \sum_{m=1}^{k} a_m \right|$.

These inequalities are also valid in the multidimensional case (see e.g. [3]). For function from anisotropic spaces with a vector parameter $\mathbf{p} = (p_1, ..., p_n)$ , it is necessary to consider a problem of summability of Fourier coefficients when some parameters $p_i$ more than 2, and others less or equal to 2.This article is devoted to the problem.

**Keywords:** Anisotropic Lorentz space, Summability of the Fourier coefficients, Hardy-Littlewood-Polya inequality

**2010 Mathematics Subject Classification:** 42B08, 42C10

**References:**

[1] Bary N.K., "Trigonometric series", Vol. I and II, Macmillan, New York, 1964.
[2] Nursultanov E.D., "On multipliers of Fourier series in a trigonometric system", *Math. Notes*, Vol. 63, No. 1–2, pp. 205–214, 1998.
[3] Nursultanov E. D., "On the coefficients of multiple Fourier series in $L_p$-spaces", *Izvestiya Rossiiskoi Akademii Nauk. Seriya Matematicheskaya*, Vol. 64, No. 1, pp. 95–122, 2000.

## Integral operators with two variable integration limits on the cone of monotone functions

Ryskul OINAROV [1], Ainur TEMIRKHANOVA [1]

[1] *L.N. Gumilyov Eurasian National University, Astana, Kazakhstan*
*E-mail: o_ryskul@mail.ru*
*E-mail: ainura-t@yandex.kz*

**Abstract:** The work is done together with A.A. Kalybay.

Let $I = (a, b)$, $-\infty \le a < b \le \infty$, and $1 < p, q < \infty$. Let $\omega$ and $v$ be non-negative and measurable functions almost everywhere finite on $I$ such that $\omega^{-q'}$, $\omega^{q}$, $v^{p'}$ and $v^{-p'}$ are locally summable on $I$.

Let $M\downarrow$ and $M\uparrow$ be sets of functions non-increasing and non-decreasing on $I$, respectively.

For the integral operators

$$K_- f(x) = \int\limits_{\alpha(x)}^{\beta(x)} K(s,x) f(s) ds \quad \text{and} \quad K_+ f(x) = \int\limits_{\alpha(x)}^{\beta(x)} K(x,s) f(s) ds$$

we consider the inequalities

$$\|\omega K_- f\|_q \le C \|v f\|_p, \quad f \in M\downarrow, \tag{1}$$

$$\|\omega K_+ f\|_q \le C \|v f\|_p, \quad f \in M\uparrow, \tag{2}$$

where $\|\cdot\|_p$ is the standard norm of the space $L_p(I)$. Moreover, the boundary functions $\alpha$ and $\beta$ satisfy the following conditions:

$(i)$ $\alpha(x)$ and $\beta(x)$ are functions differentiable and strictly increasing on $I$;

$(ii)$ $\alpha(x) < \beta(x)$ for any $x \in I$; $\lim\limits_{x\to a^+} \alpha(x) = \lim\limits_{x\to a^+} \beta(x) = a$ and $\lim\limits_{x\to b^-} \alpha(x) = \lim\limits_{x\to b^-} \beta(x) = b$.

In the case when $K(s,x) \equiv 1$ the operator $K_- f(x)$ is called the Hardy-Steklov operator. Weighted inequalities of the form (1) and (2) for the Hardy-Steklov operators have been investigated with success [1]. The similar problem for operators with kernels has remained unsolved up to now. The aim of this work is to get necessary and sufficient conditions for the validity of inequalities (1) and (2) when the kernels $K(\cdot,\cdot)$ are from a large class.

**2010 Mathematics Subject Classification:** 26D10, 39B62

**References:**

[1] Stepanov V.D., Ushakova E.P., "Hardy operator with variable limits on monotone functions", *J. Funct. Spaces Appl.*, Vol.1, No.1, pp. 1-15, 2003.

## Additive and multiplicative weighted estimates of intermediate integral operator

Ryskul OINAROV [1]

[1] *L.N. Gumilyov Eurasian National University, Astana, Kazakhstan*
*E-mail: o_ryskul@mail.ru*

**Abstract:** The work is done together with A.A. Kalybay.

We consider the inequalities

$$\|uK_\beta f\|_q \le C\left(\|\rho f\|_p + \|vKf\|_r\right), \quad f \ge 0, \tag{1}$$

$$\|uK_\beta f\|_q \le C\|\rho f\|_p^\alpha \|vKf\|_r^{1-\alpha}, \quad f \ge 0, \tag{2}$$

where $0 \le \alpha, \beta \le 1$ and $\|\cdot\|_p$ is the standard norm of the space $L_p(0,\infty)$. Moreover, $K$ and $K_\beta$ are integral operators of the form

$$Kf(x) = \int_0^\infty K(x,s)f(s)ds, \quad K_\beta f(x) = \int_0^\infty K^\beta(x,s)f(s)ds$$

with the kernel $K(\cdot,\cdot) \ge 0$ satisfying the condition

$$\exists D \ge 1 : D^{-1}\Big(K(x,t) + K(t,s)\Big) \le K(x,s) \le D\Big(K(x,t) + K(t,s)\Big)$$

for $x \ge t \ge s > 0$.

We give necessary and sufficient conditions for the validity of inequalities (1) and (2) for $1 \le \max\{p,r\} \le q \le \infty$ and $\frac{\alpha}{p} + \frac{1-\alpha}{r} \ge \frac{1}{q}$, respectively.

As an application, we obtain criteria for the inequalities

$$\|uf^{(k)}\|_q \le C\left(\|\rho f^{(n)}\|_p + \|vf\|_r\right)$$

and

$$\|uf^{(k)}\|_q \le C\|\rho f^{(n)}\|_p^\alpha \|vf\|_r^{1-\alpha}$$

to hold on the class of $n$-convex functions absolutely monotone on the interval $(0,\infty)$; in particular, when $(n=2)$ on the class of convex and twice differentiable functions, where $0 \le k < n$.

**Keywords:** multiplicative inequality, additive inequality, integral operator, kernel

**2010 Mathematics Subject Classification:** 26D10, 39B62

# Inverse coefficient problems for one-dimensional heat transfer with a preservation of medium temperature condition

Gulaiym ORALSYN

*Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: g.oralsyn@list.ru*

**Abstract:** We examine mathematical model models describing the process of heat transfer in a homogeneous bar with a prescribed law of changes of the average temperature. So there is an inverse problem for the heat equation in which together with finding the solution of the equation it is required to find unknown coefficient depending only on the time variable.

In the domain $\Omega = \{(x,t) : 0 < x < 1, 0 < t < T\}$ consider a problem on finding unknown coefficient $p(t)$ of the heat equation

$$u_t = u_{xx}(x,t) - p(t)u(x,t) + f(x,t)$$

and its solution satisfying the initial condition

$$u(x,0) = \varphi(x),\ 0 \le x \le 1,$$

the nonlocal boundary condition

$$u_x(0,t) = u_x(1,t) - \alpha u(1,t),\ \ u(0,t) = 0,\ \ 0 \le t \le T,$$

and the overdetermination conditions

$$\int_0^1 u(x,t)dx = E(t),\ E(t) \ne 0, 0 \le t \le T,$$

where $E(t) \in W_2^1(0,T)$. Here the parameter $\alpha$ is any positive number, and $f(x), \varphi(x)$ and $E(t)$ are given functions.

In [1] (for $\alpha < 0$) some of these conditions were removed and it is shown that the inverse problem has the unique generalized solution. This work is a continuation of these studies. We consider the case $\alpha > 0$.

**Keywords:** heat transfer, law of changes of the average temperature, inverse problem, heat equation

**2010 Mathematics Subject Classification:** 33C10, 34B30, 35J, 35P10

**References:**

[1] Oralsyn G., Sadybekov M.A., "An inverse coefficient problem of heat conductivity with a nonlocal Samarskii-Ionkin type condition", *AIP Conference Proceedings*, doi: 10.1063/1.4930442, Vol.1676, pp. 020016, 2015.

## Construction of mathematical models of extraction processes with non-local conditions by a spatial variable

Issabek ORAZOV [1], Sultanbek AYAZ [1]

[1] *Institute of Mathematics and Mathematical Modelling, Kazakhstan*
*E-mail: i_orazov@mail.ru, ayaz.sultanbek@gmail.com*

**Abstract:** In this paper we consider issues of constructing mathematical models of extraction processes from solid polydisperse porous materials considering the porosity of structure of particles, taking into account the connection of the residence time of fractions with particle size in the extractant, based on inverse problems of recovery of coefficients of diffusion processes under various variants of boundary conditions by a spatial variable.

We consider a mathematical model which models the extraction process of a target component from the polydispersed porous material. The suggested model is demonstrated by the example of a at solid material with bidispersed pores of different size in the form of a system of channels of macropores with micropores facing their walls. The macropores and the micropores in the material have homogeneous size. We model a case when micropores of the solid material (dispersed medium) are initially filled with an oil (dispersion phase), which is our target component. And the macropores are filled in with a pure solvent. In the process of extraction the oil diffuses from the micropore to the macropore, and then from the micropores to the external solvent volume, wherein the ratio of concentrations in the macropore and the micropore is taken in accordance with the linear law of adsorption. The well-posedness of the formulated mathematical model has been justified.

The theoretical mathematical science has deep enough advanced in solving inverse problems for diffusion processes. And besides, as a rule, the problems are researched under simplest selfadjoint boundary conditions by a spatial variable. Unlike the mentioned works we propose to consider the problems with more general boundary conditions by a spatial variable. The selfadjointness of the boundary conditions is not assumed, only requirement of their regularity by Birkhoff is sufficient. The inverse problems researched by us are directly obtained from mathematical models of technological processes.

**Keywords:** extraction processes, non-local conditions, inverse problems

**2010 Mathematics Subject Classification:** 35K15, 35P10, 35R30

## Nonlocal estimates for solutions of a singular higher order differential equation

Kordan OSPANOV [1,2]

[1] *Faculty of Mathematics and Mechanics, L.N. Gumilyov Eurasian National University, Astana, Kazakhstan*
[2] *Institute of Mathematics and Mathematical Modelling, Almaty, Kazakhstan*
*E-mail: ospanov_kn@enu.kz*

**Abstract:** We consider the linear differential equation of even order

$$L_0 y := y^{(2n)} + a_1 y^{(2n-1)} + a_2 y^{(2n-2)} + ... + a_{2n} y = f(x), \tag{1}$$

where $x \in \mathbf{R} = (-\infty, +\infty)$, and $a_k = a_k(x)$ $(k = 1, 2, ..., 2n)$ and $f \in L_2 = L_2(\mathbf{R})$ are given functions.

In this paper we investigate the equation (1) and give some sufficient conditions for its unique and coercive solvability. Similar questions were studied in many papers (see for instance the monographs [1], [2] and [3]). However, their results were obtained under the following conditions: the intermediate coefficients $a_s (s = 1, 2, ..., 2n - 1)$ of equation (1) are constant, or their growths are bounded above by some degrees of the function $|a_{2n}|$ at infinity. In the case $n = 1$, the equation (1) has been investigated in [4].

We give an estimate of norms of solution and its derivatives up to the order $2n$ of the equation (1), in the case $a_s$ $(s = 1, 2, ..., 2n - 1)$ are not bounded and $a_{2n}$ can be equal to zero.

**Keywords:** even order differential equation, coercive solvability, intermediate coefficient

**2010 Mathematics Subject Classification:** 34B40

**References:**

[1] Coddington E.A., Levinson N., "Theory of Ordinary Differential Equations", - *New York: McGraw-Hill*, 1955.

[2] Naimark M.A., "Linear Differential Operators, Vols. I, II", - *New York: Ungar*, 1968.

[3] Fedoryuk M.V., "Asymptotic analysis: linear ordinary differential equations", *Berlin: Springer*, 1993.

[4] Ospanov K., Akhmetkaliyeva R.D., "Separation and the existence theorem for second order nonlinear differential equation", *Electronic Journal of Qualitative Theory of Differential Equations*, No.66, pp. 1-12, 2012.

## Coercive solvability of degenerate system of second order difference equations

Kordan N. OSPANOV [1], Assylbek ZULKHAZHAV [1]

[1] *Faculty of Mathematics and Mechanics, L.N. Gumilyov Eurasian National University, Astana, Kazakhstan*
*E-mail: ospanov_kn@enu.kz, asilbekpin@mail.ru*

**Abstract:** We consider the following infinite system of second order difference equations:

$$l_0 y = -\Delta^{(2)} y + R\Delta y = F, \tag{1}$$

where $y = \{y_j\}_{j=-\infty}^{+\infty}$, $\Delta y = \{\Delta y_j\}_{j=-\infty}^{+\infty} = \{y_{j+1} - y_j\}_{j=-\infty}^{+\infty}$, $\Delta^{(2)} y = \{\Delta(\Delta y_j)\}_{j=-\infty}^{+\infty} = \{y_{j+1} - 2y_j + y_{j-1}\}_{j=-\infty}^{+\infty}$, $F = \{F_j\}_{j=-\infty}^{+\infty}$ and $R = (r_{i,j})_{i,j=-\infty}^{+\infty}$ is a real matrix.

Coercive solvability of the diagonal system

$$-\Delta^{(2)} y_j + \hat{R}\Delta y_j + Q y_j = F_j (\ j \in \mathbf{Z}),$$

where $\hat{R} = diag\{r_j,\ j \in \mathbf{Z}\}$, $Q = diag\{q_j,\ j \in \mathbf{Z}\}$ was discussed in [1]. In the case where oscillation of the sequence $\{r_j\}_{j=-\infty}^{j=+\infty}$ satisfies some conditions, we proved that this system is unique and coercive solvable.

In this paper, for difference system (1), we obtain some sufficient conditions of the solvability in the space of all square-summable real sequences. We also will establish the coercive estimate for its solution.

**Keywords:** difference system, solvability, coercive estimates of solutions

**2010 Mathematics Subject Classification:** 39A06

**References:**

[1] Ospanov K.N., Zulkhazhav A., "On the estimates of solutions of second order systems of difference equations", *AIP Conference Proceedings*, doi: 10.1063/1.4930497, Vol.1676, pp. 020071, 2015.

## Approximation in time of fractional equations

Sergey PISKAREV

*Lomonosov Moscow State University, Moscow, Russia*
*E-mail: piskarev@gmail.com*

**Abstract:** In this talk we continue our investigations [1] on discretization of differential equations of fractional order in time.

Recently, in [2] and [3] were considered the relation between well-posed Cauchy problems

$$v'(t) = A^l v(t) + g(t), t > 0, v(0) = u^0,$$

and

$$(\mathbf{D}_t^{1/l} u)(t) = Au(t), t > 0, u(0) = u^0.$$

Moreover, they have shown that for such kind problems with the operator $A$ which generates analytic $C_0$-semigroup one has $v(t) \equiv u(t)$ for any $t > 0$ as soon as $l = 2$ and special choice of $g(t)$.

In this talk, we would like to use such kind of relations for discretization of differential equations of fractional order in abstract spaces.

**Acknowledgement.**

Research was partially supported by grant of RFBR 15-01-00026-a.

**Keywords:** fractional differential equations, approximation

**2010 Mathematics Subject Classification:** 65N40, 35-99

**References:**

[1] Liu R., Li M., Pastor J., and Piskarev S., " On the approximation of fractional resolution families." *Differential Equations*, Vol.50(7), pp. 927-937, 2014.

[2] Keyantuo V., Lizama C., "On a connection between powers of operators and fractional Cauchy problems." *J. Evol. Equ.*, Vol.12, No.2, pp. 245-265, 2012.

[3] Li M., Chuang Ch., Li Fu-Bo., "On fractional powers of generators of fractional resolvent families." *J. Funct. Anal.* Vol.259, No.10, pp. 2702-2726, 2010.

# Sobolev type equations of time-fractional order with periodical boundary conditions

Marina PLEKHANOVA [1]

[1] *Department of Applied Mathematics, South Ural State University, Chelyabinsk, Russia*
*E-mail: mariner79@mail.ru*

**Abstract:** The main purpose of the paper is studying of the unique solvability for initial boundary value problem to the fractional differential equation

$$D_t^\alpha Lx(t) = Mx(t) + N(t, x(t), x^{(1)}(t), \dots, x^{(m-1)}(t)), \tag{1}$$

where the operators $L$ and $M$ are differential operators with respect to the spatial variables, $N$ is a nonlinear operator, $\alpha > 0$, $m \in \mathbb{N}$, $m - 1 < \alpha \leq m$, $D_t^\alpha$ is the Gerasimov–Caputo derivative. Initial conditions have the Cauchy form and boundary conditions are periodical with respect to every spatial variable on a parallelepiped. Such problems arise in many engineering and scientific disciplines as the mathematical modelling of systems and processes in the fields of physics, chemistry, aerodynamics, electrodynamics of complex medium, polymer rheology [1]. Using some results on fractional differential equations in Banach spaces from [2] the local unique solvability conditions are found for the initial boundary value problem to equation (1). General results are applied to the research of the time-fractional order Benjamin–Bona–Mahony–Burgers and Allair partial differential equations.

Equations of form (1) not solved with respect to the time derivative are called Sobolev type equations [3]. Some classes of Sobolev type fractional equations were studied in [4].

**Keywords:** Caputo fractional derivative, Sobolev type equation, nonlinear equation, initial boundary value problem, periodical boundary condition

**2010 Mathematics Subject Classification:** 35R11,34A08

**References:**

[1] Mainardi F., Spada G., Creep, relaxation and viscosity properties for basic fractional models in rheology, *The European Physics Journal, Special Topics*, Vol.193, pp. 133-160, 2011.

[2] Plekhanova M.V., Nonlinear equations with degenerate operator at fractional Caputo derivative, *Mathematical Methods in the Applied Sciences*, in press, doi: 10.1002/mma., Vol.3830, 2016.

[3] Sviridyuk G.A., Fedorov V.E., *Linear Sobolev Type Equations and Degenerate Semigroups of Operators*, Utrecht, VSP, Boston, 2003.

[4] Balachandran K., and Kiruthika S., Existence of Solutions of Abstract Fractional Integrodifferential Equations of Sobolev Type, *Comput. Math. Appl.*, Vol.64, pp. 3406-3413, 2012.

## Pohozhaev identities and applications for quasilinear equations of mixed elliptic-hyperbolic type

Nedyu POPIVANOV

*Faculty of Mathematics and Informatics, University of Sofia, Bulgaria,*
*E-mail: nedyu@fmi.uni-sofia.bg*

*Dedicated to 70th birthday of Prof. Tynysbek Kalmenov.*

**Abstract:** It is well known result of Pohozhaev (1965), that the homogeneous Dirichlet problem for semilinear elliptic equations in a bounded subset $\Omega$ of $\mathbb{R}^n$, with $n > 2$, permits only the trivial solution if the domain is star-shaped, the solution is sufficiently regular, and the power of nonlinearity $p > 2^*(n) := 2n/(n-2)$, where the latter quantity is the critical exponent in the Sobolev embedding of $H_0^1(\Omega)$ into $L^p(\Omega)$ for $p < 2^*(n)$. To the opposite of this fact, in the case $2 < p < 2^*(n)$ there exist nontrivial solutions. In the last 50 years the Pohozhaev identities and results have been used and extended for a large class of elliptic problems. Let us mention now that in [1], [2] it has been shown that the nonexistence principle in supercritical case also holds for certain two dimensional problems for the mixed elliptic-hyperbolic Gellersted operator $L$ (instead of $\Delta$), with some appropriate boundary conditions. It is also valid for a large class of such problems even in higher dimensions [3]. In dimension 2, such operators have a long-standing connection with transonic fluid flow. Of course, the critical Sobolev embedding in this case is for a suitable weighted version of $H_0^1(\Omega)$ into $L^p(\Omega)$. As usual, in the BVP for such mixed elliptic-hyperbolic Gellersted operator $L$, the boundary data are given only on the proper subset of the boundary $\partial\Omega$. To compensate the lack of a boundary condition on a part of boundary, a sharp Hardy-Sobolev inequality is used, as was first done in [1], [2] and later in [3], [4]. Some further results, already published or in progress, prepared jointly with colleagues from Italy and Norway will be also discussed.

**Keywords:** quasilinear equation, equation of mixed elliptic-hyperbolic type, Pohozhaev identities.

**2010 Mathematics Subject Classification:** 35B33, 35L80, 35M10

**References:**

[1] Lupo D., and Payne K., "Critical exponents for semi-linear equations of mixed elliptic-hyperbolic types", *Comm. Pure Appl. Math.*, Vol. 56, pp. 403–424, 2003.

[2] Lupo D., and Payne K., "Conservation laws for equations of mixed elliptic-hyperbolic type", *Duke Math. J.*, Vol. 127, pp. 251–290, 2005.

[3] Lupo D., Payne K., and Popivanov N., "Nonexistence of nontrivial solutions for supercritical equations of mixed elliptic-hyperbolic type", *Progress in Non-Linear Differential Equations and Their Applications*, Birkhuser Verlag Basel, Vol. 66, pp. 371–390, 2005.

[4] Lupo D., Payne K., and Popivanov N., "On the degenerate hyperbolic Goursat problem for linear and nonlinear equations of Tricomi type", *Nonlinear Analysis, Series A: Theory, Methods & Appl.*, Vol. 108, pp. 29–56, 2014.

## To the theory of linear ordinary differential equations of fractional order with constant coefficients

Arsen PSKHU

*Institute of Applied Mathematics and Automation, Nalchik, Russia*
*E-mail: pskhu@list.ru*

**Abstract:** Consider the equation

$$\sum_{k=1}^{m} \lambda_k \frac{d^{\sigma_k}}{dx^{\sigma_k}} u(x) = f(x),$$

where $d^{\sigma_k}/dx^{\sigma_k}$ is a fractional derivative of order $\sigma_k$ ($\sigma_k \geq 0$), $\lambda_k \in \mathbb{R}$. The fractional differentiation is given by the Dzhrbashyan-Nersesyan operator [1], associated with the sequence $\{\gamma_0^k, \gamma_1^k, \cdots, \gamma_{n(k)}^k\}$.

In the work we discuss the questions concerning the construction of the fundamental solution, the representation of the general solution and the correct form of initial conditions for the equation under consideration.

**Keywords:** fractional derivative, Dzhrbashyan-Nersesyan operator, fractional order differential equation

**2010 Mathematics Subject Classification:** 34A08

**References:**

[1] Dzhrbashyan M.M., Nersesyan A.B., "Fractional derivatives and the Cauchy problem for differential equations of fractional order", *Izv. Akad. Nauk Armenian SSR Matem.*, Vol.3, No.1, pp. 3-28, 1968 (in Russian).

---

## Some properties of linear operators in locally convex cones

Asghar RANJBARI

*Department of Pure Mathematics, Faculty of Mathematical Sciences, University of Tabriz, Tabriz, Iran*
*E-mail: ranjbari@tabrizu.ac.ir*

**Abstract:** A cone is defined to be a commutative monoid $\mathcal{P}$ together with a scalar multiplication by nonnegative real numbers satisfying the same axioms as for vector spaces. A locally convex cone is a generalization of topological vector space. The theory of locally convex cones uses an order theoretical concept or a convex quasiuniform structure to introduce a topological structure on a cone (see [2] and [3]). We verify some proprties of linear operators between two locally convex cones [1].

**Keywords:** locally convex cones, quasiuniform structures

**2010 Mathematics Subject Classification:** 46A03

**References:**

[1] Ayaseh D., Ranjbari A., Bornological locally convex cones, Matematiche (Catania), V0l.69(2), pp. 267-284, 2014.

[2] Keimel K., Roth W., Ordered cones and approximation, Vol.1517 of Lecture Notes in Mathematics, Springer-Verlag, Berlin, 1992.

[3] Roth W., Operator-valued measures and integrals for cone-valued functions, Vol.1964 of Lecture Notes in Mathematics, Springer-Verlag, Berlin, 2009.

## Some problems with nonlocal conditions for hyperbolic equations

Ludmila PULKINA

*Samara National Research University, Russia*
*E-mail: louise@samdiff.ru*

**Abstract:** In this report we present some new results concerning with nonlocal problems for hyperbolic equations. Namely, we consider a problem with nonlocal integral condition and present some methods to study it. It is well known that classical methods most widely used to prove solvability of initial-boundary problems break down as a rule when applied to nonlocal problems. In this talk, we focus on spatial nonlocal integral condition of the second kind and consider a problem: find a solution of

$$\mathcal{L}u \equiv u_{tt} - (a(x,t)u_x)_x + c(x,t)u = f(x,t) \tag{2}$$

in $Q_T = (0,l) \times (0,T)$, $l, T < \infty$, satisfying initial data

$$u(x,0) = 0, \quad u_t(x,0) = 0, \tag{3}$$

boundary condition

$$u_x(0,t) = 0, \tag{4}$$

and nonlocal condition

$$\alpha(t)u(l,t) + \int_0^l K(x)u(x,t)dx = 0. \tag{5}$$

Two methods to prove solvability of the problem will be presented in the talk. The choice of a method depends on properties of the function $\alpha(t)$ in (5). We propose a new approach which anables to prove a unique solvability of the nonlocal problem with degenerate integral condition and modification of certain known one for the case $\alpha(t) = 1$.

**Keywords:** nonlocal problem, hyperbolic equation, integral conditions dynamical boundary conditions, degenerate integral condition

**2010 Mathematics Subject Classification:** 35L10, 35L20, 35L99

**References:**

[1] Pulkina L.S. "A Nonlocal Problem for Hyperbolic Equation with Degenerate Integral Condition", *EJDE*, Vol.2016, No.193, pp. 1–12, 2016.

## To theory one class of integral equation by tube domain

Nusrat RAJABOV

*Research Institute Tajik National University, Tajikistan*
*E-mail: nusrat38@mail.ru*

**Abstract:** Let $\Omega$ denote the tube $\Omega = \{(t,z) : a < t < b, |z| < R\}$. Lower ground this cylinder denote by $D = \{t = a,\ |z| < R\}$ and lateral surface denote by $= \{a < t < b,\ |z| = R\}$, $z = x + iy$. In $\Omega$ we shall consider the following integral equation

$$\varphi(t,z) + \lambda \int_a^x \frac{\varphi(\tau,z)}{\tau - a} d\tau - \frac{\mu}{\pi} \iint_D \frac{exp[i\theta]\,\varphi(t,s)}{(R-\rho)^{\beta}(s-z)} ds -$$

$$\frac{\delta}{\pi} \int_a^x \frac{d\tau}{\tau - a} \iint_D \frac{exp[i\theta]\,\varphi(\tau,s)}{(R-\rho)^{\beta}(s-z)} ds = f(t,z), \tag{1}$$

where $\lambda$, $\mu$, $\delta$ – are given real constants, $f(t,z)$ – are given function, $\theta = \arg s$, $s = \xi + i\eta$, $ds = d\xi d\eta$, $\varphi(t,z)$ – unknoun function, $\rho^2 = \xi^2 + \eta^2$.

The solution to this equation is sought in the class of function $\varphi(t,z) \in C\left(\overline{\Omega}\right)$, $\varphi(a,z) = 0$, $\varphi(t, Re^{i\theta}) = 0$, $\theta = \arg z$ and its asymptotic behavior at $t \to a$, $r \to R$ is given by formulas

$\varphi(t,z) = 0\left[(t-a)^{\varepsilon}\right]$, $\varepsilon > 0$ at $t \to a$, $\varphi(t,z) = 0[(R-r)^{\delta_1}]$, $\delta_1 > \beta - 1$, at $t > a$.

For integral equation (1) found condition to parameters, present in kernels, at fulfilment which, the problem found solution this type integral equation reduce, to problem found two splitting system integral equation, theory which detailed investigation in author works. In this case solution integral equation (1) found in explicit form. In those cases, when general solution contain arbitrary function, found inverse formula, that is arbitrary functions found by valued solution integral equation in surface cylinder. In the case, when constants present in kernels no among themselves, the solution integral equation(1) found in form absolutely and uniformly converges generalized power series by power $(t-a)$, by infinite numbers analytic functions variable point ground this cylinder. Found invers formula, that is arbitrary analytic functions founded by valued solution integral equation and its derivative at $t = 0$. Obtained integral representation and its inverse formula use for stand and investigate Dirichlet type boundary value problems.

**Keywords:** integral equation, tube domain, two splitting system integral equation

**2010 Mathematics Subject Classification:** 45E05, 45B05

# About solutions properties of Tricomi problem for Gellerstedt equation

Alexander ROGOVOY

*Department of Computer sytems and design, South-Kazakhstan Humanitarian Institute for the Humanities, Shymkent, Kazakhstan*
*E-mail: rog2005@list.ru*

**Abstract:** Tricomi problem research has a fundamental importance for all theory of mixed type equations and various areas of application. This problem for Tricomi equation $y\,u_{xx}+u_{yy}=f(x,y)$ was originally suggested in [1]. Later more common Gellerstedt equation $(sgny)\,y^m\,u_{xx}+u_{yy}=f(x,y)$ was suggested in [2].

We consider the Tricomi problem for the Gellerstedt equation for the case in which the elliptic part of the boundary is the curve

$$\sigma_\delta=\left\{(x,y):\,\left(x-\frac{1}{2}\right)^2+\left(\frac{2}{m+2}y^{\frac{m+2}{2}}+\delta\right)^2=\frac{1}{4}+\delta^2,\,-\infty<\delta<+\infty\right\}.$$

Smooth solutions for this domain have been got in [3]. The notion of an $n$-regular solution for model Lavrent'ev-Bitsadze equation has been introduced in [4]. In present work we prove a criterion for the existence of $n$-regular solution of Tricomi problem for Gellerstedt equation.

**Keywords:** Tricomi problem, Gellerstedt equation, non continuous solutions

**2010 Mathematics Subject Classification:** 35M10, 35M12

**References:**

[1] Tricomi, F., "Sulle equazioni lineari alle derivate parziali di $2^0$ ordine di tipo misto", *Mem. Lincei*, Ser. V, XIV, fasc. VII, 1923.

[2] Gellerstedt, S., "Sur un probleme aux limites pour une equation lineaire aux derivees partielles du second ordre de type mixte", *Thesis*, Uppsala, 1935.

[3] Bazarbekov, A.B., "Smoothness of the Solution of the Tricomi Problem for Equations of Mixed Type", *Differ. Uravn.*, Vol.25, No.5, pp. 816-822, 1989.

[4] Rogovoy, A.V., "Properties of Solutions of the Tricomi Problem for the Lavrent'ev-Bitsadze Equation at Corner Points", Vol. 49, No. 12, pp. 1692-1696, 2013.

## On $I_\lambda$-statistical convergence of weight $g$

Ekrem SAVAS [1]

[1] *Istanbul Commerce University, Turkey*
*E-mail: ekremsavas@yahoo.com*

**Abstract:** In this paper, following a very recent and new approach of [1], we further generalize the recently introduced summability method of [2] and introduce the new notion namely, $I_\lambda-$statistical convergence of weight $g$,where $g:\mathbb{N}\to[0,\infty)$ is a function satisfying $g(n)\to\infty$ and $n/g(n)\nrightarrow 0$. We mainly investigate certain properties of this convergence.

**Keywords:** Ideal, filter, $\mathcal{I}$-statistical convergence of order $g$, $\mathcal{I}_\lambda-$statistical convergence

**2010 Mathematics Subject Classification:** 40A05, 40D25

**References:**

[1] Balcerzak M., Das P., Filipczak M., Swaczyna J., Generalized kinds of density and the associated ideals, to appear in Acta Math. Hungar.
[2] Savas E., Pratulananda Das, A generalized statistical convergence via ideals, *Appl. Math. Letters*, Vol.24, pp. 826-830, 2011.

# Asymptotically $\mathcal{I}_\lambda$–Statistical Equivalent Sequences of weight $g$

Ekrem SAVAS [1]

[1] *Istanbul Commerce University, Istanbul, Turkey*
*E-mail: ekremsavas@yahoo.com*

**Abstract:** This paper presents the following definition which is a natural combination of the definition for asymptotically equivalent of weight $g$, $\mathcal{I}$ -statistically limit, and $\lambda$- statistical convergence, where $g : \mathbb{N} \to [0, \infty)$ is a function satisfying $g(n) \to \infty$ and $n/g(n) \nrightarrow 0$. The two nonnegative sequences $x = (x_k)$ and $y = (y_k)$ are said to be asymptotically $\mathcal{I}^g$- statistical equivalent of weight $g$ to multiple $L$ provided that for every $\epsilon > 0$, and $\delta > 0$,

$$\{n \in \mathbb{N} : \frac{1}{g(\lambda_n)} |\{k \in I_n : |\frac{x_k}{y_k} - L| \geq \epsilon\}| \geq \delta\} \in \mathcal{I},$$

(denoted by $x \overset{S_\lambda^L(I)^g}{\sim} y$) and simply asymptotically $\mathcal{I}^g$- statistical equivalent of weight $g$ if $L = 1$. In addition, we shall also present some inclusion theorems.

**Keywords:** Asymptotical equivalent, ideal convergence, $\mathcal{I}$-statistical convergence, $\lambda$- statistical convergence, statistical convergence of weight $g$

**2010 Mathematics Subject Classification:** 40A05, 40D25

**References:**

[1] Balcerzak M., Das P., Filipczak M., Swaczyna J., Generalized kinds of density and the associated ideals, Acta Math. Hungarica 147(1), pp. 97-115, 2015.

[2] Gumus H., and Savas E., On $S_\lambda^L(I)-$Asymptotically Statistical Equivalent Sequences, Numerical Analysis and Appl. Math. ( ICNAAM- 2012), AIP conference proceeding, V. 1479, pp. 936-941, 2012.

[3] Kumar V., and Sharma A., On Asymptotically Generalized Statistical Equivalent Sequences via ideal, Tamkang J.Math. 43(3), pp. 469-478, 2012

[4] Li J., Asymptotic Equivalence of Sequences and Summability, Internat. J. Math. & Math. Sci. 20(4), pp. 749-758, 1997.

[5] Marouf M., Asymptotic equivalence and summability. Internat. J. Math. Math. Sci., Vol.16(4), pp. 755-762, 1993.

[6] Patterson R.F., On Asymptotically Statistically Equivalent Sequences, Demonstratio Math., Vol.36(1), pp. 149-153, 2003.

[7] Savaş E., On $\mathcal{I}-$ asymptotically lacunary statistical equivalent sequences, Advances in Difference Equations 2013, 2013:111 (18 April 2013).

[8] Savaş E., On asymptotically $\mathcal{I}-$statistical equivalent sequences of order $\alpha$, Indian J. Math., Special Volume Dedicated to Professor Billy E. Rhoades 56(2), 2014.

# Weak solutions for nonlinear fractional differential equations with fractional separated boundary conditions in Banach spaces

Djamila SEBA [1,2], Rebai HAMZA [2]

[1,2] *Department of Mathematics, University of Boumerdes, Algeria*
*E-mail: djam_seba@yahoo.fr*
[2] *Laboratory of Dynamical Systems, University of Sciences and Technology Houari Boumediene, Bab-Ezzouar, Algeria*
*E-mail: rebaihamza14@gmail.com*

In this paper, we consider the following boundary value problem of fractional differential equation with fractional separated Boundary Conditions of the form (1)

$$\begin{cases} {}^cD^r x(t) = f(x, x(t)), \quad t \in J = [0,\, 1],\ 1 < r \leq 2. \\ \\ \alpha_1 x(0) + \beta_1({}^cD^p x(0)) = \gamma_1, \quad \alpha_2 x(1) + \beta_2({}^cD^q x(1)) = \gamma_2,\ 0 < p < 1 \end{cases}$$

where ${}^cD^r$ is the Caputo fractional derivative of order $r$, $f : J \times E \to E$ is a given function satisfying some assumptions that will be specified later, $\alpha_i$, $\beta_i$, $\gamma_i$ $(i = 1,\, 2)$ are suitably chosen constants in $E$, with $\alpha_1 \neq 0$. and $E$ is a Banach space with norm $\|.\|$.

The topic of fractional differential equations has been of great interest for many researchers in view of its theoretical development and widespread applications in various fields of science and engineering such as physics, biophysics, chemistry, statistics,economics, blood flow, phenomena, control theory, porous media, electromagnetic, and other fields. Boundary value problems with integral boundary conditions constitute an important class of problems and arise in the mathematical modeling of various phenomena such as heat conduction, wave propagation, gravitation, chemical engineering, underground water flow, thermoelasticity, and plasma physics. They include two-point, three-point, multipoint boundary value problems.

Our investigation relies upon the method associated with the technique of measures of noncompactness and the fixed point theorem of Mönch type.

**Keywords and phrases:** Caputo fractional derivative, Riemann Liouville integral, measure of noncompactness, fixed point, Banach space.

**2010 Mathematics Subject Classification: 26A33**

# Hardy-type inequality for a fractional integral operator in $q$-analysis

S. SHAIMARDAN

*Faculty of Mathematics and Mechanics, L.N. Gumilyov Eurasian National University, Astana, Kazakhstan*
*E-mail: email@address*

**Abstract:** We consider the operator $I_{q,n}$ the following form

$$I_{q,n}f(x)=\frac{1}{\Gamma_q(n)}\int\limits_0^\infty \mathcal{X}_{(0,x]}(s)K_{n-1}(x,s)f(s)d_qs,$$

which is defined for all $x>0$ [1]. where $K_{n-1}(x,s)=(x-qs)_q^{n-1}$.

Then the $q$-analog of the two-weighted inequality for the operator $I_{q,n}$ of the form

$$\left(\int\limits_0^\infty u^r(x)\left(I_{q,n}f(x)\right)^r d_qx\right)^{\frac{1}{r}}\leq C\left(\int\limits_0^\infty v^p(x)f^p(x)d_qx\right)^{\frac{1}{p}}. \tag{1}$$

which has several applications in various fields of science. Where $C$ a positive constants independent of $f$ and $u(\cdot),v(\cdot)$ are positive real valued functions on $(0,\infty)$, i.e. weight functions.

**Theorem.** Let $1<r<p<\infty$. Then the inequality (1) holds if and only if $Q_{n-1}<\infty$ holds, where

$$\begin{aligned}Q_m^{n-1} &= \left\{\int\limits_0^\infty\left(\int\limits_0^\infty \mathcal{X}_{(0,z]}(s)K_m^{p'}(z,s)v^{-p'}(s)d_qs\right)^{\frac{p(r-1)}{p-r}}\right.\\ &\times \left(\int\limits_0^\infty \mathcal{X}_{[z,\infty)}(x)K_{n-m-1}^r(x,z)u^r(x)d_qx\right)^{\frac{p}{p-r}}\\ &\times \left.D_q\left(\int\limits_0^\infty \mathcal{X}_{(0,z]}(s)K_m^{p'}(z,s)v^{-p'}(s)d_qs\right)\right\}^{\frac{p-r}{pr}}.\end{aligned}$$

Moreover, $Q_{n-1}\approx C$, $C$ is the best constant in (1).

**Keywords:** inequalities; Hardy-type inequalities, integral operator; q-calculus; q-integral

**2010 Mathematics Subject Classification:** 26D10, 26D15, 39A13

**References:**

[1] Al-Salam, W.A., "Some fractional $q$-integrals and $q$-derivatives", *Proc. Edinb. Math. Soc*, Val. 15, pp.135–140, (1966/1967).

## On the applicability of Picard's successive approximations to the numerical solving the problem of heat transfer in domain, degenerate at the initial time

Yu.R. SHPADI, A.A. KAVOKIN, A.T. KULAKHMETOVA

*Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: yu-shpadi@yandex.ru*

**Abstract:** The following problem for the heat equation is considered:

$$\frac{\partial u}{\partial t} = a^2 \frac{\partial^2 u}{\partial x^2}, \qquad 0 < x < \alpha_0 t, \quad 0 < t < T, \tag{1}$$

$$u(0,t) = \varphi(t), \quad (2) \qquad u(\alpha(t),t) = \psi(t), \quad (3) \qquad \varphi(0) = \psi(0) = 0, \quad (4)$$

where known boundary functions $\varphi(t)$ and $\psi(t)$ are Hölder continuous:

$$|\varphi(t') - \varphi(t'')| < A|t' - t''|^\sigma, \quad |\psi(t') - \psi(t'')| < A|t' - t''|^\sigma, \quad \sigma > \frac{1}{2}. \tag{5}$$

We represent the solution of problem (1)–(5) as $u(x,t) = u_1(x,t) + W(x,t)$, where $u_1(x,t)$ and $W(x,t)$ – are double layer potentials with density $\varphi(t)$ and unknown density $\theta(t)$, respectively. Function $u(x,t)$ satisfies equation (1) and condition (2), and substituting it into equation (3) we obtain the following integral equation:

$$\theta(t) = g(t) + \int_0^t K(t,\tau)\theta(\tau)d\tau, \tag{6}$$

where $W(\alpha_0 t, t) = \int_0^t K(t,\tau)\theta(\tau)d\tau$ is the valeur principale of the potential $W(x,t)$ at $x = \alpha_0 t$, and $g(t) = u_1(\alpha_0 t, t) - \psi(t)$. It was shown in [1] that the equation (6), in general, may not have a solution when $g(t)$ is just the bounded continuous functions. Moreover, the integral operator (6) can have eigenfunction, [3]. Properties of equation (6) was studied Properties of equation (6) were studied in papers [2], [4], [5], etc.

We establish in this work the additional conditions on the boundary functions in (2), (3) for which the solution of equation (6) exist, is unique, and can be found by the Picard's method of successive approximations. For this we define the functional class $M_\beta(0,T)$which contains continuous functions $f(t)$ having estimation $|f(t)| \le M_f t^\varepsilon$, where $\varepsilon > \beta$. The following results are proved:

**Theorem 1.** If the function $g(t) \in M_{\frac{1}{2}}(0,T)$ , then the Picard's iterative process converges uniformly to the unique solution of integral equation and $\theta(t) \in M_0(0,T)$.

**Theorem 2.** The function $g(t) \in M_{\frac{1}{2}}(0,T)$ under the conditions (4),(5).

**Keywords:** Heat equation in degenerating domain, Volterra Singular integral operator, Picard's method

**2010 Mathematics Subject Classification:** 35K05, 35K20, 45D05

**References:**

[1] Kharin S.N., "The thermal processes in the electrical contacts and the related singular integral equations (rus.)", *Abstr.of PhD diss.*, Alma-Ata, p.16, 1970.

[2] Omarov T.E., Otelbaev M.O. "On a class of singular integral equations of Volterra type II (rus.)",*Mathematical investigation*, V3, Karaganda, pp.12-19, 1976.

[3] Ramazanov M.I., "Investigation of eigenvalues and eigen-functions of the singular integral equation of the Volterra second kind (rus.)", in *Proceedings of Differential equations and their application*, Alma-Ata, pp.83-90, 1979.

[4] Shpadi Y.R. "A Heat equation problems in domains with varying cross-section (rus.)" *Abstr.of PhD diss.*, Almaty, p.16, 1998.

[5] Jenaliev M.T at al. "About Dirichlet boundary value problem for the heat equation in angular domain", *Boundary value problems*, pp.1-21, 2014.

## The Hardy-Littlewood type theorems in Besov spaces with the Haar basis

Yesmukhanbet SMAILOV [1]

[1] *Institute of Applied Mathematics, Karaganda, Kazakhstan*
*E-mail: esmailov@mail.ru*

**Abstract:** In this paper there were established the coefficient necessary and sufficient conditions that the function belongs to the Besov spaces with the Haar basis.

Let $H = \{h_m(t)\}_{m=1}^{+\infty}$, $t \in [0,1]$ be the orthonormal Haar system [1].

We denote by $\|f\|_p$ the norm of function from $L_p[0,1]$, $1 \le p < +\infty$.

Let $f \in L_p[0,1]$, $1 \le p < +\infty$. Let $E_n(f)_{p\theta}$ be the best approximation of function $f$ in the metrics of Lebesgue space $L_p[0,1]$ by the Haar polynomials of order not higher than $n$.

**Definition 1.** *Let $1 \le p \le +\infty$, $1 \le \theta \le +\infty$, $r > 0$. We say that function $f \in L_p[0,1]$ belongs to the Besov space with the Haar basis, if*

$$|f; B^r_{p\theta}([0,1]; H)| = \|f\|_p + \left\{ \sum_{k=0}^{+\infty} 2^{kr\theta} E^{\theta}_{2^k}(f)_p \right\}^{\frac{1}{\theta}}, \quad 1 \le \theta < +\infty;$$

$$|f; B^r_{p\infty}([0,1]; H)| = \|f\|_p + \sup \left\{ 2^{kr} E_{2^k}(f)_p \mid k \in \mathbb{Z}^+ \right\}.$$

The following two theorems contain $GM$-monotone condition on [2] the Fourier-Haar coefficients of function $f \in B^r_{p\theta}([0,1], H)$.

**Theorem 1.** Let $f \in L[0,1]$. Let $f(x) \sim \sum\limits_{k=1}^{+\infty} a_k h_k(x)$ be its the Fourier-Haar series. If the positive sequence of the Fourier-Haar coefficients $\{a_k\} \in GM$, then for $f \in B^r_{s\theta}([0,1]; H)$, where $r > 0$, $1 < s < +\infty$, $1 \le \theta \le +\infty$, it is sufficient that for some $p$: $1 \le p < s$

$$\left\{ \sum_{k=1}^{+\infty} k^{r\theta + \theta\left(\frac{1}{2} + \frac{1}{p} - \frac{1}{s}\right) - 1} a_k^{\theta} \right\}^{\frac{1}{\theta}} < +\infty. \tag{2}$$

Moreover, the following inequality

$$|f; B^r_{s\theta}([0,1]; H)| \le A_{psr\theta} \left\{ \sum_{k=1}^{+\infty} k^{r\theta + \theta\left(\frac{1}{2} + \frac{1}{p} - \frac{1}{s}\right) - 1} a_k^{\theta} \right\}^{\frac{1}{\theta}}.$$

holds. Here $A_{psr\theta} > 0$ depends only on $p$, $s$ and $r$.

**Theorem 2.** Let $r > 0$, $1 < s < \lambda < +\infty$, $1 \le \theta \le +\infty$. Let $f \in B^r_{s\theta}([0,1];H)$. If its Fourier-Haar coefficients are positive and $GM$-monotone, then

$$|f; B^r_{s\theta}([0,1];H)| \ge D_{\lambda sr\theta} \left\{ \sum_{k=1}^{+\infty} k^{r\theta+\theta\left(\frac{1}{2}+\frac{1}{\lambda}-\frac{1}{s}\right)-1} a_k^{\theta} \right\}^{\frac{1}{\theta}}.$$

Here $D_{\lambda sr\theta} > 0$ depends only on $\lambda$, $s$, $r$ and $\theta$.

**Keywords:** Fourier series, Haar system, Fourier-Haar coefficients, $GM$-monotone sequences, Hardy-Littlewood theorem, Besov space

**2010 Mathematics Subject Classification:** 42A16, 46E30

**References:**

[1] Golubov B. I., "Series with respect to the Haar system", *Journal of Soviet Mathematics*, Vol.1, No.6, pp. 704–726, 1973.

[2] Tikhonov C., "Trigonometric series with general monotone coefficients", *Mathematical analysis and applications*, Vol.326, pp. 721-735, 2007.

---

## Integrals with homogeneous - difference kernels on the plane and their applications

Alexander P. SOLDATOV

**Abstract:** Two-dimensional singular integrals and one- dimensional generalized integrals of Cauchy type are considered. The applications to the Riemann -Hilbert problem for a general first order elliptic system on the plane are given. More exactly it is shown how to reduce the problem to a system of Fredholm integral equation on the domain and a singular integral equation on its boundary with the help of homogeneous difference integrals.

The work was supported by International Project (3492/GF4) Ministry of Education and Science of Kazakhstan Republic

---

## Confluent hyper geometric function sand two variables Laguerre polynomials as a solutions of Wilczynski type system

Zhaksylyk TASMAMBETOV [1]

[1] *School of Mathematics and Physics, Aktobe Regional Zhubanov State University, Kazakhstan*
*E-mail: tasmam@rambler.ru*

**Abstract:** In this work it is shown that in the study of systems, solutions of which are Whittaker functions and two variables Laguerre polynomials, the main role is played by Humbert confluent hypergeometric function $\Psi_2(\lambda, \gamma, \gamma'; x, y)$.

It is defined from Appell functions $F_2(\alpha,\beta,\beta',\gamma,\gamma';x,y)$ using the passage to the limit [1, p.121], and it is the solution of the following system

$$(1)\qquad \begin{cases} x\cdot Z_{xx}+(\gamma-x)\cdot Z_x-y\cdot Z_y-\lambda\cdot Z=0,\\ y\cdot Z_{yy}+(\gamma'-y)\cdot Z_y-x\cdot Z_x-\lambda\cdot Z=0, \end{cases}$$

where $\gamma\neq 0,-1,-2,...;\ \gamma'\neq 0,-1,-2,...;\ Z=Z(x,y)$ – general unknown.

The first particular solution of Wilczynski type system (1) is confluent hypergeometric function $Z_1(x,y)=\Psi_2(\lambda;\gamma,\gamma';x,y)$. In [2] it is shown that Humbert confluent hypergeometric function $\Psi_2(\lambda;\gamma,\gamma';x,y)$ is the main tool of investigating the Whittaker type systems. Here, it is introduced that it plays the same role at investigation of Laguerre type systems and simplify its construction of solution in the form of two variables Laguerre polynomials.

With this purpose, from confluent hypergeometric system ($\Psi_2$) at $\gamma=\alpha+1$ ($\alpha>-1$), $\gamma'=\beta+1$ ($\beta>-1$), it is defined the following system

$$(2)\qquad \begin{cases} x\cdot Z_{xx}+(\alpha+1-x)\cdot Z_x-y\cdot Z_y-\lambda\cdot Z=0,\\ y\cdot Z_{yy}+(\beta+1-y)\cdot Z_y-x\cdot Z_x-\lambda\cdot Z=0. \end{cases}$$

Considering the first particular solution $Z_1(x,y)$, it is easy to be assured that the solution in the form of polynomial exists in the case when $\lambda=-n$ ($n$ – integer), is polynomial

$$Z_1(x,y)=\Psi_2(-n;\alpha+1,\beta+1;x,y).$$

**Theorem.** Let $\alpha>-1$, $\beta>-1$, $\alpha\neq 0$, $\beta\neq 0$. The system of second order partial differential equations (2), where $\lambda$ – parametric value, has as a solution the two variables polynomial, not equal to zero identically, if and only if $\lambda$ is in the form of $\lambda=-n$ ($n$ – integer).

Let us call this polynomial the generalized two variables Laguerre polynomial and denote it by

$$L_{n,n}^{(\alpha,\beta)}(x,y)=\Psi_2(-n;\alpha+1,\beta+1;x,y).$$

It is also established that the system of the following type

$$\begin{cases} x^2\cdot U_{xx}-xy\cdot U_y+\left(-\frac{x^2}{4}-\frac{xy}{2}+kx+\frac{1}{4}-\alpha^2\right)\cdot U=0,\\ y^2\cdot U_{yy}-xy\cdot U_x+\left(-\frac{y^2}{4}-\frac{xy}{2}+ky+\frac{1}{4}-\beta^2\right)\cdot U=0, \end{cases}$$

where $k=(\alpha+\beta+2-2\lambda)/2$ is also the Laguerre type system with the solution of the following form

$$U(x,y)=x^{\frac{\alpha+1}{2}}\cdot y^{\frac{\beta+1}{2}}\cdot\exp\left(-\frac{x}{2}-\frac{y}{2}\right)\cdot\sum_{n,m=0}^{\infty}\frac{(\frac{\alpha+\beta}{2}+\lambda)_{m+n}}{(2\alpha+1)_m\cdot(2\beta+1)_n}\cdot\frac{x^m}{m!}\cdot\frac{y^n}{n!}.$$

This aspect is still remaining little-investigated. A variety of theorems is proved.

**References:**

[1] Appell, P., Kampe de Feriet, J. F. "Fonctions hypergeometriques et hyperspheriques: Polynomes d'Hermite. *Gauthier – Villars, Paris*, 1926, p. 434.

[2] Tasmambetov, Zh. N."On irregular singular curves of Whittaker type system *Vestnik Sam. GTV, Seria Fiziko-mathematicheskie nauki*, 2003. Issue 4 (33). p. 25-33.

## On one algorithm to find a solution to a linear two-point boundary value problem

Svetlana TEMESHEVA

Department of Mathematics, Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan
E-mail: nur15@mail.ru

**Abstract:** A two-parameter family of algorithms for finding an approximate solution to a linear two-point boundary value problem for a system of ordinary differential equations is offered. The convergence conditions for the algorithms are obtained. The necessary and sufficient coefficient conditions for the well-posedness of considered problem are established.

Consider a linear two-point boundary value problem

$$\frac{dx}{dt} = A(t)x + f(t), \qquad x \in \mathbb{R}^n, \qquad t \in (0, T), \tag{3}$$

$$Bx(0) + Cx(T) = d, \tag{4}$$

where $A(t)$ and $f(t)$ are continuous on $[0, T]$, $B$ and $C$ are the given $(n \times n)$ matrices, $d$ is a given $n$ vector, $\|x\| = \max\limits_{i=1:n} |x_i|$, and $\|A(t)\| = \max\limits_{i=1:n} \sum\limits_{j=1}^{n} |a_{ij}(t)| \leqslant \alpha$, $\alpha = \text{const}$.

Present article offers an other family of algorithms of parameterizations method [1]. Herein each algorithm's step also consists of two items. The difference of offering algorithms is in item $b$), where the unknown functions $u_r(t)$, $t \in [(r-1)h, rh)$, $r = 1 : N$, are to be computed by the recurrent formula.

**Keywords:** two-point boundary value problem, linear ordinary differential equation, algorithm to find a solution, necessary and sufficient conditions for unique solvability

**2010 Mathematics Subject Classification:** 34B16, 34B40

**References:**

[1] Dzhumabayev D.S., *Criteria for the unique solvability of a linear boundary-value problem for an ordinary differential equation*, USSR Computational Mathematics and Mathematical Physics, 29:1, pp. 34-46, ISSN 0041-5553, Moscow, 1989.

## On the convergence of some iterative algorithms to approximate fixed points of nonexpansive mappings

Balwant Singh THAKUR [1]

[1] *School of Studies in Mathematics, Pt.Ravishankar Shukla University Raipur, India*
*E-mail: balwantst@gmail.com*

**Abstract:** In this paper we present a new iterative algorithm to approximate fixed points of nonexpansive and pseudocontractive mappings. We establish convergence results. We also discuss numerical examples to illustrate theoretical results. In addition, we compare the convergence of these iterations with existing iterations using the numerical computation.

---

## Oscillation of second order half-linear difference equation and discrete Hardy inequality

Ainur TEMIRKHANOVA

*L.N. Gumilyov Eurasian National University, Astana, Kazakhstan*
*E-mail: ainura-t@yandex.kz*

This is joint work with R. Oinarov, A.A. Kalybay and S. Kh. Shalginbayeva

**Abstract:** We consider a second order half-linear difference equation of the following form

$$\Delta(\rho_i|\Delta y_i|^{p-2}\Delta y_i) + v_{i+1}|y_{i+1}|^{p-2}y_{i+1} = 0, \quad i = 0, 1, 2, ..., \tag{1}$$

where $1 < p < \infty$, $\Delta y_i = y_{i+1} - y_i$, $\{\rho_i\}$ - a sequence of positive real numbers and $\{v_i\}$ - a sequence of non-negative real numbers.

In this work for researching oscillation properties of equation (1) we use the variational method, which is reduced to the validity or not validity of the discrete Hardy inequality on the set of finitary sequences. On the basis of the criteria of validity of the discrete Hardy inequality in a local term we establish sufficient, necessary conditions of conjugate, disconjugate on the interval $[m, n]$, $0 \le m < n \le \infty$ of equation (1) and also oscillation, nonoscillation of this equation.

From these results as a corollary we obtained the different interesting conditions of oscillation of the equation (1). In particular, we obtained, that if $\{t_k\} \subset \mathbb{N}$, $\mathbb{N} - \{t_k\} \neq \oslash$, $t_k \to \infty$, as $k \to \infty$, $v_{t_k} \neq 0$, $v_i = 0$, $i \in \mathbb{N} - \{t_k\}$ and the following condition

$$v_{t_k} > \rho_{t_k - 1} + \left( \sum_{i=t_k+1}^{\infty} \rho_i^{1-p'} \right)^{1-p},$$

holds for all $k$, then the equation (1) is oscillatory.

**Keywords:** discrete Hardy inequalities, oscillatory, nonoscillatory, half-linear equations, disconjugate, conjugate, variational principle

**2010 Mathematics Subject Classification:** 39A10, 39A21, 34C10, 26D15.

# Periodic boundary value problem for a system of ordinary differential equations with impulse effects

Agila TLEULESOVA

*L.N. Gumilyov Eurasian national university, Astana, Kazakhstan*
*Institute of Mathematics and Mathematical Modelling, Almaty, Kazakhstan*
*E-mail: agila_72@mail.ru*

**Abstract:** In this work, we investigated a nonlinear periodic boundary value problem with impulse effects. We have found conditions of existence isolated solution of periodic boundary value problem for system of nonlinear differential equations with impulse effects.

We consider the following periodic boundary value problem for a system of nonlinear ordinary differential equations with impulse effects on $[0,T]$:

$$dx/dt = f(t,x), \quad t \in [0,T] \setminus \{\theta_1, \theta_2, \ldots, \theta_m\}, \quad x \in R^n,$$

$$0 = \theta_0 < \theta_1 < ... < \theta_{m+1} = T, \tag{1}$$

$$x(0) = x(T), \tag{2}$$

$$x(\theta_i + 0) - x(\theta_i - 0) = J_i\left(\lim_{t \to \theta_i - 0} x(t)\right), \qquad i = \overline{1,m}, \tag{3}$$

where $f : [0,T] \times R^n \to R^n$, is a piecewise-continuous vector valued function with points of discontinuity of the first kind at $t = \theta_i \, (i = \overline{1,m})$ and $J_i(x)$ $(i = \overline{1,m})$ are continuous vector valued functions of $x$. Many authors discuss solvability of problem (1) - (3) [1–5]. Sufficient conditions for the existence of isolated solutions periodic boundary value problem of impulsive.

**Keywords:** Impules effects, boundray value, differential equation

**2010 Mathematics Subject Classification:** 34B05, 34B37

**References:**

[1] Samoilenko A.M., and Perestyuk N.A., "Impulse Differential Equations" *Vyshcha Shkola*, Kiyv, p. 405, 1987.

[2] Samoilenko A.M., and Ronto N.I., "Numerical-Analytic Methods for the Investigation of the Solutions of Boundary-Value Problems" *Naukova Dumka*, Kiyv, p. 565, 1986.

[3] Liz E., and Nieto J., "Positive solutions of linear impulsive differential equations" *Communications in applied analysis* 2, pp. 565-571, 1998.

[4] Dzhumabaev D.S., and Temesheva S.M., "A Parametrization Method for Solving Nonlinear Two-Point Boundary Value Problems" *Computational Math. and Math. physics* Vol.47, pp. 37-61, 2007.

[5] Dzhumabayev D.S., "Criteria for the unique solvability of a linear boundary-value problem for an ordinary differential equation" *USSR Computational Math. and Math. Physics* Vol.29, pp. 34-46, 1989.

## About sedately growing solutions of one generalized Cauchy-Riemann system

Zhanuzak TOKIBETOV [1], Gulzhan ABDUAKHITOVA [1], Ulbike KUSHERBAYEVA [1]

[1] *Department of Fundamental Mathematics, Al-Farabi Kazakh National University, Almaty, Kazakhstan*

*E-mail: tokibetov@mail.ru, gulzhan_ae@mail.ru, ulbi-70@mail.ru*

**Abstract:** In this paper the existence of only zeroth sedately growing solution of the generalized Cauchy-Riemann system is proved in the case when the coefficients of the generalized Cauchy-Riemann system do not belong to the class $L_{p,2}(E)$, $p > 2$, namely $A(z) = az$, $B(z) = b\bar{z}$, where $a$, $b$ are constants.

Consider the following problem: it is required to find regular solutions of the system

$$\partial w/\partial \bar{z} + azw + b\bar{z}\bar{w} = 0 \tag{1}$$

(where $a$, $b$ are complex numbers), that in the neighborhood of the point of infinity satisfy the following condition

$$|w| < K|z|^N, \tag{2}$$

where $N$ is an arbitrary nonnegative number and $K$ is an arbitrary real number.

The problem, when coefficients of the system (1) from class $L_{p,2}(E)$, $p > 2$, (function $f(z)$ at point $z = x + iy$ belongs to the class $L_{p,2}(E)$, if

$$L_{p,2}(E) = \|f\|_{p,2} = \Big\{ \iint\limits_{|z|\le 1} |f(z)|^p dx\ dy \Big\}^{\frac{1}{p}} + \Big\{ \iint\limits_{|z|\le 1} \Big| z^{-2} f(\frac{1}{z}) \Big|^p dx\ dy \Big\}^{\frac{1}{p}}$$

have been studied by I.N.Vekua [1], and when coefficients are constants - by V.S.Vinogradov [2]. In our case, coefficients do not belong to the class $L_{p,2}(E)$, $p > 2$. We will try to find the solution of equation (1) in the form of representation [3]: $f(z) = F(z)e^{if(z)} + G(z)e^{ig(z)}$, where $F(z)$, $G(z)$ are arbitrary analytic functions, and $f(z)$, $g(z)$ are continuous functions.

**Theorem 2.** *If in a system* (1) *the coefficient a is a real number, then the problem* (1) , (2) *has only trivial solution.*

**Keywords:** Generalized Cauchy-Riemann system, sedately growing solutions

**2010 Mathematics Subject Classification:** 30E25

**References:**

[1] Vekua I., "Obobshhennye analiticheskie funkcij" , *Fizmatgiz*, Moscow, p.600, 1959 (in Russian).

[2] Vinogradov V., "O teoreme Liuvillja dlja obobshennykh analiticheskikh funkcij", *DAN SSSR*, Vol.183, No.3, pp. 503-506, 1968 (in Russian).

[3] Tokibetov Z., " teoreme Liuvillja dlja obobshennoj sistemy Cauchy - Riemann", *Vestnik KazNU. Seriya matematika, mechanika, informatika*, Vol.44, No.1, pp. 91-100, 2005 (in Russian).

## On a class of fractional elliptic problems with an involution perturbation

Batirkhan Kh. TURMETOV [1], Berikbol T. TOREBEK [2]

[1] *Akhmet Yasawi University, Turkistan, Kazakhstan*
*E-mail: batirkhan.turmetov@ayu.edu.kz*
[2] *Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan*
*E-mail: torebek@math.kz*

**Abstract:** To describe the problems, let $\Omega = \{(x,y) \in R^2 : 0 < x < 1, -\pi < y < \pi\}$. The paper is concerned with four boundary value problems concerning the fractional analogue of Helmholtz equation with a perturbation term of involution type in the space variable:

$$D_x^{2\alpha} u(x,y) + u_{yy}(x,y) - \varepsilon u_{yy}(x,-y) - c^2 u(x,y) = 0, (x,y) \in \Omega, \quad (1)$$

where $c, \varepsilon$ are real numbers, $D_x^{2\alpha}$ means $D_x^{2\alpha} = D_x^{\alpha} D_x^{\alpha}$ and the operator $D_x^{\alpha}$ acts by the variable $x$.
Here

$$D_x^{\alpha} u(x,y) = \frac{1}{\Gamma(1-\alpha)} \int_0^x (x-s)^{-\alpha} \frac{\partial u}{\partial s}(s,y) ds$$

is a Caputo differentiation operator of the $\alpha \in (0,1]$ order [1].

Regular solution of Equation (1) is a function $u \in C(\bar{\Omega})$, such that $D_x^{\alpha} u, D_x^{2\alpha} u$, $u_{yy} \in C(\Omega)$. Since for $\alpha = 1 : {}_C D^1 = \frac{d}{dy}$, then $D_x^2 + \frac{\partial^2}{\partial y^2} = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} = \Delta$. Therefore, Equation (1) is a nonlocal generalization of the Helmholtz equation, which at $\varepsilon = 0$ coincides with the Helmholtz equation.

We obtain for them existence and uniqueness results based on the Fourier method.

**Keywords:** Caputo operator, Helmholtz equation, involution, fractional differential equation, Mittag-Leffer function, boundary value problem

**2010 Mathematics Subject Classification:** 34A08, 35R11, 74S25

**References:**

[1] Kilbas A.A., Srivastava H.M., and Trujillo J.J., *Theory and Applications of Fractional Differential Equations*, Elsevier. North-Holland. Mathematics studies, 2006.

## Four-dimensional generalized difference matrix and some double sequence spaces

Orhan TUĞ [1], Feyzi BAŞAR [2]

[1] *Department of Mathematics Education, Ishik University, Erbil, Iraq*
*E-mail: orhan.tug@ishik.edu.iq, tugorhan27@gmail.com*
[2] *Prof.Dr., Turkey*
*E-mail: feyzibasar@gmail.com*

**Abstract:**In this study, we introduce some new double sequence spaces $B(\mathcal{M}_u)$, $B(\mathcal{C}_p)$, $B(\mathcal{C}_{bp})$, $B(\mathcal{C}_r)$ and $B(\mathcal{L}_q)$ as the domain of four-dimensional generalized difference matrix $B(r,s,t,u)$ in the spaces $\mathcal{M}_u$, $\mathcal{C}_p$, $\mathcal{C}_{bp}$, $\mathcal{C}_r$ and $\mathcal{L}_q$, respectively. We show that the double sequence spaces $B(\mathcal{M}_u)$, $B(\mathcal{C}_{bp})$ and $B(\mathcal{C}_r)$ are the Banach spaces under some certain conditions. We give some inclusion relations with some topological properties. Moreover, we determine the $\alpha-$dual of the spaces $B(M_u)$ and $B(\mathcal{C}_{bp})$, $\beta(\vartheta)-$duals of the spaces $B(M_u)$, $B(\mathcal{C}_p)$, $B(\mathcal{C}_{bp})$, $B(\mathcal{C}_r)$ and $B(\mathcal{L}_q)$, where $\vartheta \in \{p, bp, r\}$ and $\gamma-$dual of the spaces $B(\mathcal{M}_u)$, $B(\mathcal{C}_{bp})$ and $B(\mathcal{L}_q)$. Finally, we characterize the classes of four dimensional matrix mappings defined on the spaces $B(\mathcal{M}_u)$, $B(\mathcal{C}_p)$, $B(\mathcal{C}_{bp})$, $B(\mathcal{C}_r)$ and $B(\mathcal{L}_q)$ of double sequences.

**Keywords:** four-dimensional generalized difference matrix; matrix domain; double sequence spaces; alpha-dual; beta-dual; gamma-dual; matrix transformations.

**2010 Mathematics Subject Classification:** 46A45, 40C05.

**References:**

[1] Boos J., "Classical and Modern Methods in Summability", *Oxford university Press, New York*, 2000.

[2] Hardy G.H., "On the convergence of certain multiple series", *Proc. Cambridge Philos. Soc.*, Vol.19 pp. 86-95,(1916-1919).

[3] Móricz F., "Extensions of the spaces $c$ and $c_0$ from single to double sequence", *Acta. Math. Hungar.*, Vol.57, pp. 129-136, 1991.

[4] Başar F., Sever Y., "The space $\mathcal{L}_q$ of double sequences", *Math. J. Okayama Univ.*, Vol.51, pp. 149-157, 2009.

[5] Zeltser M., "On conservative matrix methods for double sequence spaces", *Acta Math. Hung.*, Vol.95(3), pp. 225-242, 2002.

[6] Adams C.R., "On non-factorable transformations of double sequences", *Proc. Natl. Acad. Sci. USA*, Vol.19(5), pp. 564-567, 1933.

[7] Cooke R.C., "Infinite Matrices and Sequence Spaces", *Macmillan and Co. Limited, London,* 1950.

## Four-dimensional generalized difference matrix and almost convergent double sequence spaces

Orhan TUĞ [1], Feyzi BAŞAR [2]

[1] *Department of Mathematics Education, Ishik University, Erbil, Iraq*
*E-mail: orhan.tug@ishik.edu.iq, tugorhan27@gmail.com*
[2] *Prof.Dr., Turkey*
*E-mail:feyzibasar@gmail.com*

**Abstract:** The concept of almost convergence for single sequence was introduced by Lorentz [1] and have studied by several mathematicians (see [2–4]), and for double sequences was introduced by Moricz and Rhoades [5].

In this study, the domains $B(\mathcal{C}_f)$ and $B(\mathcal{C}_{f_0})$ of four-dimensional generalized difference matrix in the spaces of almost convergent and almost null double sequences are introduced and some inclusion relations are given with some topological properties. The $\alpha-, \beta(\vartheta)-$ and $\gamma-$ duals of those spaces are calculated, where $\vartheta = \{p, bp, r\}$. Finally, some classes of four dimensional matrix transformations on those new double sequence spaces are characterized with some significant consequences.

**Keywords:** four-dimensional generalized difference matrix, almost convergence, double sequence spaces, $\alpha$-dual, $\beta$-dual and $\gamma$-dual, matrix transformations.

**2010 Mathematics Subject Classification:** 46A45, 40C05

**References:**

[1] Lorentz G.G., "A contribution to the theory of divergent sequences", *Acta mathematica*, Vol.80(1), pp. 167-190, 1948.

[2] Petersen G.M., "Almost convergence and uniformly distributed sequences", *The Quarterly Journal of Mathematics*, Vol.7(1), pp. 188-191, 1956.

[3] Butkovich D., Kraljevich H., and Sarapa N., "On the almost convergence", *Functional Analysis II. Springer Berlin Heidelberg*, pp. 396-417, 1987.

[4] Başar F., and Kirişçi M., "Almost convergence and generalized difference matrix", *Computers and Mathematics with Applications*, Vol.61(3), pp. 602-611, 2011.

[5] Moricz F., Rhoades B.E., "Almost convergence of double sequences and strong regularity of summability matrices", *Mathematical Proceedings of the Cambridge Philosophical Society, Cambridge University Press*, Vol.104(02), 1988.

## Multiplier Amenability of Banach Algebras

Faruk UYGUL

*Department of Mathematics and Statistics, American University of Sharjah, United Arab Emirates*
*E-mail: fuygul@aus.edu*

**Abstract:** Amenability is a distinctive property of locally compact groups and Banach algebras related to them. The amenability of a locally compact group can be characterized by many properties such as the Følner condition and Leptin's condition. Furthermore. the amenability of locally compact groups can also be studied by the properties of certain Banach algebras related to them (such as the Fourier and the Fourier Stieltjes algebras). In this talk we will discuss and present some new results on the amenability properties of Banach algebras in terms of their multiplier algebras.

**Keywords:** amenable, locally compact, measure algebra, Fourier algebra

**2010 Mathematics Subject Classification:** 43A07; 43A10; 43A20

**References:**

[1] Daws M., Dual Banach Algebras: Representations and Injectivity, Studia Math., Vol.178, pp. 231-275, 2007.
[2] Day M. M., Means on semigroups and groups, Bull. Amer. Math. Soc. 55, pp. 1054-1055, 1949.
[3] Ruan Z.J., The operator amenability of $A(G)$, Amer. J. Math., Vol.17, pp. 1449-1474, 1995.

---

## q-Sumudu transforms of basic generalized hypergeometric functions

Vijay Kumar VYAS [1]

[1] *The ICFAI University Jaipur, India*
*E-mail: vmathsvyas@gmail.com*

**Abstract:** The object of this paper is to evaluate q-Sumudu transforms of the basic analogue of H-function of two variables. Interesting special cases of theorems are also discussed. Further, the results proved in this paper may find certain applications of q-Sumudu transforms to the solutions of the q-integrodifferential equations involving functions.

**Keywords:** q-analogues of Fox's H-function of two variables, q-Sumudu transform, q-analogues of Meijjer's G-function of two variables

## On periodic wave factorization

Vladimir VASILYEV [1]

[1] *Pure Mathematics, Lipetsk State Technical University, Lipetsk, Russia*
*E-mail: vbv57@inbox.ru*

**Abstract:** We consider a special kind of factorization related to certain domains in multidimensional complex space $\mathbb{C}^m$. This approach was introduced by the author [1, 2] for studying solvability of model elliptic pseudo-differential equations in canonical domains. Now we interested in discrete variant of such a theory, and for this purpose we construct a periodic wave factorization to describe solvability properties of some classes of discrete equations.

Let $D \subset \mathbb{R}^m$ be a convex sharp cone, $T(D) \subset \mathbb{C}^m$ is a domain of the following type

$$T(D) = \{z \in \mathbb{C}^m : \ z = x + iy, x \in \mathbb{T}^m, y \in D\}.$$

We say that a function $A(\xi), \xi \in \mathbb{T}^m, A(\xi) \neq 0, \forall \xi \in \mathbb{T}^m$, admits periodic wave factorization with respect to cone $D$ if it can be represented in the form

$$A(\xi) = A_+(\xi)A_-(\xi),$$

where the factors $A_\pm(\xi)$ admit bounded analytical continuation into complex domains $T(\pm D)$ respectively.

If we consider a discrete equation in a cone then an existence of periodic wave factorization with respect to an appropriate cone is a sufficient condition for unique solvability of such equation in Lebesgue spaces. Some preliminary results for special degenerated cones were obtained in [3, 4].

The work was supported by Russian Foundation for Basic Research and government of Lipetsk region of Russia, grant no. 14-41-3595-r-center-a.

**Keywords:** discrete equation, periodic wave factorization, multidimensional Riemann boundary value problem

**2010 Mathematics Subject Classification:** 35J46, 42B37

**References:**

[1] Vasilyev V.B., "Wave factorization of elliptic symbols", *Math. Notes*, Vol.68, No. 5-6, pp. 556-568, 2000.

[2] Vasilyev V.B., "Elliptic equations and boundary value problems in non-smooth domains", *Operator Theory: Advances and Applications*, Vol.213, pp. 105-121, 2011.

[3] Vasilyev A.V., Vasilyev V.B., "Discrete singular operators and equations in a half-space", *Azerb. J. Math.*, Vol.3, No.1, pp. 84-93, 2013.

[4] Vasilyev A.V., Vasilyev V.B., "On some classes of difference equations of infinite order", *Adv. Difference Equ.*, Vol.211, pp. 1-9, 2015.

## On multipliers of Fourier series in the Lorentz space

Aizhan YDYRYS [1], Nazerke TLEUKHANOVA [2]

[1] *Department of Fundamental Mathematics, L.N.Gumilyov Eurasian National University, Astana, Kazakhstan*
*E-mail: aizhanyd@gmail.com*

[2] *Department of Fundamental Mathematics, L.N.Gumilyov Eurasian National University, Astana, Kazakhstan*
*E-mail: tleukhanova@rambler.ru*

**Abstract:** We study the multipliers of Fourier series on the Lorentz spaces, in particular, the sufficient conditions for a sequence of complex numbers $\{\lambda_k\}_{k\in Z}$ in order to make it a multiplier of trigonometric Fourier series of space $L_{p,r}[0;1]$ in the $L_{q,r}[0;1]$. In this work there is a new multipliers theorem which is supplement of the well-known theorems, and given a counterexample.

**Keywords:** Multiplier, Fourier series, Lizorkin theorem, Lorentz space

**2010 Mathematics Subject Classification:** 42A45

**References:**

[1] Reisz M., *Mathematische Zeitschrift*, Vol.27, pp. 218-244, 1928.

[2] Marcinkiewicz J., *Studia Mathematica*, Vol.8, pp. 78-91, 1939.

[3] Mikhlin S., *Vestn. Leningr. Univ., Mat. Mekh. Astron.*, Vol.12, pp. 143-155, 1957.

[4] Nursultanov E., and N.Tleukhanova, "Multipliers of multiple Fourier series", Proceedings of the Steklov Institute of Mathematics, pp. 237–242, 1999.

[5] Nursultanov E., *Mathematical Notes*, Vol.62, pp. 945-949, 1997.

[6] Hörmander L., *Acta Math.*, V0l.104, pp. 93-140, 1960.

[7] Nursultanov E., and Tleukhanova N., *Functional Analysis and its Applications*, Vol.34, pp. 151–153, 2000.

[8] Lizorkin P.I., *Trudy Mat.Inst.Steklov*, Vol.89, pp. 231-248, 1967.

[9] Sarybekova L., Tararykova T., and Tleukhanova N., *Mathematical Inequalities & Applications*, Vol.13, pp. 613–624, 2010.

[10] Persson L.E., Sarybekova L., and Tleukhanova N., *Proceedings of A. Razmadze Mathematical Institute*, Vol.151, pp. 83–101, 2009.

[11] Persson L.E., Sarybekova L., and Tleukhanova N., "A Lizorkin theorem on Fourier series multipliers for strong regular systems", in *Analysis for science, engineering and beyond*, Springer, Heidelberg, vol. 6 of Springer Proc. Math., pp. 305-317, 2012.

## About the defect indices of fourth-order differential operator with rapidly oscillating coefficients

Alibek YESKERMESSULY [1]

[1] *L.N.Gumilyov Eurasian National University, Kazakhstan*
*E-mail: aleke1410@gmail.com*

**Abstract:** We study in the space $L_2(0,\infty)$ of a minimum nonsemibounded differential operator $L_0$ [1], [2], generated by the differential expression

$$ly = y^{(4)} - (h(x)y')' - q(x)y, \quad x \in (0;\infty) \tag{1}$$

$$y^{(4)} - (h(x)y')' - q(x)y = \lambda y, \tag{2}$$

Where $\lambda \in \mathbf{C}$, $Im\,\lambda \neq 0$, the function $q(x)$ satisfies the conditions of regularity Titchmarsh-Levitan [3], and $h(x)$ is rapidly oscillating disturbance, which does not meet the conditions of the Titchmarsh-Levitan.

$$\varphi(\xi,\lambda) = \frac{h(x)}{4q^{\frac{1}{4}}(x)(q(x)+\lambda)^{\frac{1}{4}}}, \qquad \omega(\xi,\lambda) = \frac{q'(x)}{8q^{\frac{1}{4}}(x)(q(x)+\lambda)}, \qquad x = \nu(\xi).$$

$$\left| \int_{\xi}^{\infty} \varphi(s,\lambda)ds \right| < \infty, \tag{3}$$

$$\varphi_1(\xi,\lambda) = \int_{\xi}^{\infty} \varphi(s,\lambda)ds.$$

$$\mu_0\varphi_1(\xi,\lambda) \in L_1(0,\infty), \tag{4}$$

$$\varphi_1(\xi,\lambda)\omega(\xi,\lambda) \in L_1(0,\infty), \tag{5}$$

$$\mu_0\varphi_1^2(\xi,\lambda) \in L_1(0,\infty). \tag{6}$$

**Theorem 3.** *Let the*

**a:** $q(x) \geq c|x|^{\frac{4}{3}+\varepsilon}, \quad c > 0, \quad \varepsilon > 0, |x| \geq R;$
**b:** $q'(x)$, $q''(x)$ *do not change sign for sufficiently large* $R > 0$ *for* $|x| \geq R;$
**c:** $q'(x) = o\left(q^{\zeta}(x)\right)$ *when* $x \in (0;\infty)$, *where* $0 < \zeta < \frac{5}{4}$,

*and the following conditions* (3), (4), (5) *and* (6). *Then the deficiency indices of the operator are equal (3.3).*

**Keywords:** Defect indices, differential operator, rapidly oscillating coefficients.

**2010 Mathematics Subject Classification:** 34K08

**References:**

[1] Naimark M.A. Linear differential operators. – M .: Nauka, 1969 – 526 p.
[2] Kostyuchenko A.G., I.S.Sargsyan. Distribution of eigenvalues (self-adjoint ordinary differential operators). – M .: Nauka, 1979 – 400 p.
[3] Murtazin Kh., Sultanaev Y.T. Formulas for the distribution of the eigenvalues Nonsemibounded Sturm-Liouville // Mathematical notes, Vol. 28:4, pp. 545-553, 1980.

# On a problem for wave equation with data on the whole boundary

Nurgissa YESSIRKEGENOV

*Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan*
*Department of Mathematics, Imperial College London, London, United Kingdom*
*E-mail: n.yessirkegenov15@imperial.ac.uk*

**Abstract:** In this paper we propose a new formulation of boundary value problem for a one-dimensional wave equation in a rectangular domain in which boundary conditions are given on the whole boundary.

Let $\Omega \subset R^2$ be a rectangular domain, bounded by following lines: $AB : 0 \leq x \leq \ell$, $t = 0$, $BC : x = \ell$, $0 \leq t \leq T$, $CD : 0 \leq x \leq \ell, t = T$ and $AD : x = 0$, $0 \leq t \leq T$. Let $E = (T, T), F = (\ell - T, T)$ be points on a boundary $CD$.

We consider a nonhomogeneous wave equation in $\Omega$

$$u_{tt} - u_{xx} = f(x, t). \tag{1}$$

It is well known that the Dirichlet problem for the wave equation (1) in a rectangular domain is ill-posed.

Problem. *Find a solution of Eq.(1), satisfying the boundary condition*

$$u|_{AB \cup BC \cup AD} = 0,$$

*and conditions on the boundary $CD$:*

$$u_t|_{DE} = 0,$$
$$\alpha u_x + \beta u_t|_{EF} = 0,$$
$$u|_{CF} = 0,$$

*where $\alpha$ and $\beta$ are real numbers.*

We prove the well-posedness of boundary value problem in the classical and generalized senses, depending on the coefficients $\alpha$ and $\beta$. The proof of the well-posedness of the formulated problem is reduced to question of the existence and uniqueness of solutions of the corresponding functional equations.

**Keywords:** Wave equation, well-posedness of problems, classical solution, strong solution

**2010 Mathematics Subject Classification:** 35L05, 35L20, 49K40, 35D35

# APPLIED MATHEMATICS

## High order finite-difference method for simulation isotropic turbulence flow

Akmaral ABDIGALIYEVA [1], Zhailaubai ZHUBATOV [2], Dauren ZHAKEBAYEV [3]

[1,2,3] *Al-Farabi Kazakh National University, Almaty, Kazakhstan*
*E-mail: abdigalieva_an@mail.ru*

**Abstract:** The work deals with the modeling of turbulent energy using two different methods: finite-difference and spectral methods. To simulate the turbulent process the filtered three-dimensional non-stationary Navier-Stokes equation is used. The problem is solved by using hybrid methods, where the equation of motion is solved by using finite-difference methods in combination with cyclic penta-diagonal matrix, which allowed to reach high order of accuracy and to simulate turbulence decay at Reynolds number Re=300 for comparing with theoretical solution of Taylor and Green. The spectral method is used for solution of Poisson equation, which is makes it possible to gain the time. For validation of the given algorithm we solved the classical problem of Taylor and Green modeling of isotropic turbulence flow. Where at the work of Taylor and Green [1], the problem is solved analytically and defined all the turbulence characteristics. In the results of our simulation we have compared the turbulence characteristics with analytical solution: the change of turbulent kinetic energy, dissipation energy over the time and longitudinal and transverse one-dimensional spectra are defined. In this paper a three-dimensional non-stationary Navier-Stokes equation is solved for simulation of isotropic turbulence flow. For solving Navier-Stokes equation, we use a splitting scheme by physical parameters that consist of three stages. At the first stage, the Navier–Stokes equation is solved, without taking pressure into account. For approximation of the convective and diffusion terms of the intermediate velocity field finite-difference methods in combination with cyclic penta-diagonal matrix is used [2], which allowed to increase the order of accuracy in time and in space $O(t^3, h^4)$ without changing the amount of points. At the second stage the Poisson equation is solved, which is satisfies the continuity equation with considering the velocity field from the first stage. For solving the Poisson equation we use spectral method in combination with Fourier transform. The obtained pressure field with using fast Fourier transform is translated from the phase space to physical and used at the third stage for the recallculate of the final velocity field [3].

**Keywords:** finite-difference method, spectral method, Poisson equation, cyclic penta-diagonal matrix

**2010 Mathematics Subject Classification:** 94B05, 94B15

**References:**

[1] Taylor, G.I., Green, A.E., "Mechanism of production of small eddies from large ones", *Proceedings of the royal society*, Vol. 158, No. 895, pp.499–521, 1937.

[2] Samarskii, A.A., Nikolaev, E.S., "Methods for solving grid equations", *M.:Nauka*, pp. 73-120, 1978.

[3] Abdibekov, U.S., Zhumagulov, B.T., Zhakebayev D.B., Zhubat K.Z., "Modelling of the decay of isotropic turbulence by the LES ", *Math. Models and Comp. Simulations*, Vol.5(4), pp. 360-370, 2013.

# A new class of orthogonal polynomials (Adeolu polynomials) for derivation of initial value solvers

Emmanuel ADEYEFA [1], R.O. FOLARANMI [2]

[1] *Department of Mathematics, Federal University Oye–Ekiti, Ekiti State, Nigeria*
*E-mail: adeoluman@yahoo.com*
[2] *University of Ilorin, Ilorin, Nigeria*
*E-mail: rotsuccess@yahoo.com*

**Abstract:** This paper presents a set of newly constructed polynomials called Adeolu polynomials, valid in interval [-1, 1] with respect to weight function w(x) = x + 1 and employed as trial function to develop a fast, efficient and reliable block algorithm for the numerical solution of ordinary differential equations with application to third order initial value problems. Collocation and interpolation techniques were adopted for the formulation of self-starting continuous hybrid schemes. Findings from the analysis of the basic properties of the method using appropriate existing theorems show that the developed schemes are consistent, zero-stable and hence convergent. These continuous numerical integration schemes have the ability to yield several output of solutions at the off-grid points without requiring additional interpolation and at no extra cost. On implementation, the schemes compared favourably well with the existing methods owing to the fact that they are accurate and efficient. Further investigation of the properties of these polynomials is ongoing and in future, the scope of application shall be extended to boundary value problems.

**Keywords:** Collocation, Interpolation, Orthogonal polynomials, Block method, hybrid

**2010 Mathematics Subject Classification:** 65L05, 65L06, 94B05, 94B15

**References:**

[1] Lambert J.D.,"$\mathbb{Z}_8$-Computational Methods in Ordinary Differential Equation. John Wiley & Sons Inc.", 1973.

[2] Rosser J.B., "Runge-Kutta for all seasons. $\mathbb{Z}_4$", *SIAM*, Vol.9, pp. 417-452, 1967.

[3] Shampine L.F., and Watts, H.A., "Block Implicit One-Step Methods $\mathbb{Z}_9$", *Journal of Math of Computation*, Vol.23(108), pp. 731-740, 1969.

[4] Tanner L.H., "The spreading of Silicone Oil Drops on Horizontal Surfaces", Vol.12, pp. 1473-1484, 1979.

## Spreading of flame front I by the predictor-corrector scheme

D. ALIBIYEV[1], G. KHAKIMZYANOV[2], A. KAZHIKENOVA[1], E. IBRAEVA[1]

[1] *Karaganda State University, Karaganda, Kazakhstan*
*E-mail: dalibiev@mail.ru*
[2] *Novosibirsk State University, Novosibirsk, Russia*

**Abstract:** In this paper, we present the algorithm for the numerical solution of non-stationary equations describing the process of distribution of a flame in a two-fuel mixture. For the numerical simulations of the process of flame spreading is desired to use moving irregular meshes which are thickened to a region of high gradients of temperature and concentration of the mixture, since the calculations for such grids can have a high precision even with a small amount of nodes [1].

The proposed finite-difference scheme "predictor-corrector" in step "predictor" splitting method is used with the explicit approximation of the convective terms in the first step and an implicit fractional approximation of diffusive members on the second fractional step. This second step involves the fractional member responsible for the burning process. In addition, in step "predictor" of explicit scheme determined by the flow at the edges of cells. We have proposed a method of selecting grid parameters allowing to calculate profiles of numerical solutions without spurious oscillations.

The proposed method of adapting the grid to a moving flame front tested on scalar nonlinear convection-diffusion equation with a nonlinear right-hand side, as well as a model system of two nonlinear equations. For meshing we used equivalence distribution method that focuses on the use of time-dependent problems. We have shown that the use of adaptive grids allows to obtain a numerical solution with the same precision as in a uniform grid, but with the number of nodes on the order of less.

**Keywords:** Predictor-corrector, computational modeling, flame front, flame spreading, flame spreading, algorithm of calculation

**2010 Mathematics Subject Classification:** 97N40, 97M10

**References:**

[1] Varnets Y., Maas U., Dibbl R. Gorenie, "Physical and chemical aspects, modeling, experiments, formation of polluting sub-stances", Phizmatlit, 2003.

## The identification problem for hyperbolic Schrodinger equation

Allaberen ASHYRALYEV[1], Begench GURBANOV [2], Ramil SALIMOV [2]

[1] *Emeritus Professor, Turkey and Turkmenistan*
*E-mail: aaallaberen@gmail.com*
[2] *Faculty of Art and Science, Istanbul University, Istanbul, Turkey*
*E-mail: begagurban@gmail.com; ramilsalimov@yahoo.com*

**Abstract:** We consider the identification problem for hyperbolic "Schrodinger differential equation

$$\begin{cases} u_{tt} + Au(t) = f + p, 0 < t < 1, \\ iu_t + Au(t) = g + p, -1 < t < 0, \\ u(-1) = \varphi, u(1) = \psi \end{cases}$$

In a Hilbert space with self-adjoint positive definite operator A. We are interested in studying the stability of this problem. In applications, the stability estimates for solutions of the two problems for the hyperbolic Schrodinger equations are defined. The stable difference schemes of the approximate solution of this problem are constructed. The convergence estimates for the solutions of these difference schemes are established. In addition, the theoretical statements are supported by the results of numerical examples.

**Key words:** Difference scheme, Schrodinger equation, Stability, Well–posedness

**References:**

[1] Ashyralyev A., Sobolevskii P.E., "New Difference Schemes for Partial Dierential Equations", Operator Theory Advances and Applications, Birkhauser Verlag, Basel, Boston, Berlin, 2004.

[2] Sobolevskii P.E., "Difference Methods for the Approximate Solution of Differential Equations", Izdat. Voronezh. Gosud. Univ., Voronezh, 1975. (Russian).

[3] Ashyralyev A., Amanov D., *Abstract and Applied Analysis* , 2012, Article ID 594802, 14 p., 2012.

[4] Berezin F.A., Shubin M.A., The Schrodinger Equation, Kluwer Academic Publishers Dordrecht, Boston, London, 1991.

**Fractional derivatives and fractional integrals linked to the fractional powers of positive operators**

Allaberen ASHYRALYEV[1], Begench GURBANOV [2], Ramil SALIMOV [2]

[1] *Emeritus Professor, Turkey and Turkmenistan*
*E-mail: aaallaberen@gmail.com*
[2] *Faculty of Art and Science, Istanbul University, Istanbul, Turkey*
*E-mail: begagurban@gmail.com; ramilsalimov@yahoo.com*

**Abstract:** Fractional derivatives and fractional integrals are defined. In addition, definition for positive operators is stated. It is given how fractional derivatives and fractional integrals are connected to fractional positive operators. Finally, the difference formulas for fractional derivative and fractional integral are derived.

**Key words:** Fractional derivative, Fractional integral, Fractional power, Positive operator

**References:**

[1] Podlubny I., *Fractional differential Equations,vol. 198* of Mahematics in Science and Engineering, Academic Press, San Diego, California, USA, 1999.

[2] Kilbas A.A., Sristava H.M., and Trujillo J.J., *Theory and Applications of Fractional Differential Equations,* North-Holland Mathematics Studies, 2006.

[3] Ashyralyev A., and Sobolevskii P.E., *Well-Posedness of Parabolic Difference Equations,* Vol. 69, Birkhäuser Verlag, Basel,Boston, Berlin, 1994.

[4] Ashyralyev A., Emirov N., Cakir Z., *Well-posedness of fractional parabolic differential and difference equations with the Dirichlet-Neumann condition,* Electronic Journal of Differential Equations, Vol.2014, No.97, pp. 1-17, 2014.

[5] Tarasov V.E., *Fractional derivative as fractional power of derivative,* International Journal of Mathematics, Vol.18(3), pp. 281-299, 2007.

[6] Ashyralyev A., *Applied Mathematics Letters* , Vol.24, pp. 1176-1180, 2011.

[7] Ashyralyev A., *Journal of Mathematical Analysis and Applications* , Vol.357(1), pp. 232-236, 2009.

## Source identification problem for an elliptic-hyperbolic equation

Allaberen ASHYRALYEV[1], Fatma Songul OZESENLI TETIKOGLU[2], Tulay KAHRAMAN [3]

[1] *Emeritus Professor, Turkey & Turkmenistan*
*E-mail: aallaberen@gmail.com*
[2,3] *İstanbul, Turkey*
*E-mail: ftetikoglu@fatih.edu.tr*[2],*tulaykahraman@cihangir.k12.tr* [3]

**Abstract:** In the present paper, the boundary value problem

$$(1)\qquad \begin{cases} -\frac{d^2u(t)}{dt^2} + Au(t) = tp + f(t), \ \ (-1 \le t \le 0), \\ \frac{d^2u(t)}{dt^2} + Au(t) = tp + g(t), \ \ (0 \le t \le 1), \\ u(0) = \varphi, u(-1) = \psi, u(1) = \xi, \end{cases}$$

for the differential equation with parameter $p$ in a Hilbert space $H$ with self adjoint definite operator $A$ is investigated. The well-posedness of this problem is established. The stability inequalities for the solution of source identification problem for elliptic-hyperbolic equations are obtained.

**Keywords:** elliptic equation, hyperbolic equation, boundary value problems, stability

**2010 Mathematics Subject Classification:** 35M32, 65M06, 35J25, 35L04

**References:**

[1] Ashyralyev A., and Sobolevskii P. E., "New Difference Schemes for Partial Differential Equations", *Operator Theory: Advances and Applications, Birkhauser, Verlag, Basel, Boston, Berlin,* Vol.148, 2000.

[2] Ashyralyev A., and Judakova G., and Sobolevskii P. E., "A Note on the Difference Schemes fo Hyperbolic-Elliptic Equations", *Abstract and Applied Analysis*, Vol.2006, pp. 1-13, 2006.

[3] Eidelman Y.S., "Boundary Value Problem for Differential Equations with Parameters", *Voronezh State University,* 1984 (in Russian).

[4] Gercek O., and Zusi E, "On the Numerical Solution of a Two Dimensional Elliptic-Parabolic Equation", *AIP Conference Proceedings, Numerical Analysis and Applied Mathematics, ICNAAM 2012,* Vol.2012, pp. 606-610, 2012.

## Stable difference scheme for the solution of an elliptic equation with the involution

Allaberen ASHYRALYEV [1], Baktygul KARABAEVA [2], Abdizhahan SARSENBI [2,3]

[1] *Emeritus Professor, Turkey & Turkmenistan*
*E-mail: aallaberen@gmail.com*

[2] *Department of Mathematical Methods and Modelling, M.Auezov South Kazakhstan State University, Shymkent, Kazakhstan*
[3] *Institute of Mathematics and Mathematical Modelling, Almaty, Kazakhstan*
*E-mail: karabaeva@gmail.com, abzhahan@gmail.com*

**Abstract:** The theory of functional-differential equations with the involution has received less attention than functional-differential equations. Moreover, one of the unstudied areas of differential equations are partial differential and difference equations with the involution (see, e.g., [1]- [3] and the references given therein).

In the paper [4], the boundary value problem of an elliptic equation with the involution and nonlocal boundary conditions in $x$ was investigated. The stability and coercive stability estimates in Hölder norms in $t$ for the solution of this problem were established. In the present paper a stable difference scheme for the approximate solution of an elliptic equation with the involution is constructed. Theorem on stability and almost coercive stability and coercive stability of this difference scheme is established. The theoretical statements for the solution of this difference scheme are supported by the results of the numerical experiment.

**Keywords:** Difference scheme, Elliptic equation with the involution, Stability estimates

**PACS:** 02.30.Jr, 02.30.Ks, 02.60.Lj

**References:**

[1] Wiener J., *Generalized Solutions of Functional-Differential Equations*, World Scientific Pub., River Edge, NJ, 1993.

[2] Burlutskaya M.Sh., *Mixed problem for a first-order partial differential equation with involution and periodic boundary conditions*, Comput. Math. Math. Phys., Vol.54(1), pp. 1-10, 2014.

[3] Sadybekov M.A., Sarsenbi A.M., *Criterion for the basis property of the eigenfunction system of a multiple differentiation operator with an involution*, Differential Equations, Vol.48, pp. 1112-1118, 2012.

[4] Ashyralyev A., Sarsenbi A.M., *Well-posedness of an elliptic equation with involution*, Electronic Journal of Differential Equations, Art. No. Vol.284, 2015.

# A note on the Dirichlet problem for model complex partial differential equations

Allaberen ASHYRALYEV [1], Bahriye KARACA [2]

[1] *Emeritus Professor, Turkey & Turkmenistan*
*E-mail: aallaberen@gmail.com*
[2] *Department of Mathematics, Yeditepe University, İstanbul, Turkey*
*E-mail: bahriye.karaca@yeditepe.edu.tr*

**Abstract:** Complex model partial differential equations of arbitrary order are considered. The uniqueness of the Dirichlet problem is studied. It is proved that the Dirichlet problem for higher order of complex partial differential equations with one complex variable has infinitely many solutions. **Keywords:** Dirichlet problem, homogeneous partial differential equation, inhomogeneous partial differential equation, complex analysis with one variable **2015 Mathematics Subject Classification:** 94B05, 94B15

**References:**

[1] Begehr H., Boundary value problems in complex analysis I, II, Boletin de la Asosiación Matemática Venezolana, XII, pp. 65-85, 217-250, 2005.

[2] Bitsadze A.V., About the uniqueness of the Dirichlet problem for elliptic partial differential equations, Uspekhi Math. Nauk 3, Vol.6(28), pp. 211-212 1948 (Russian).

[3] Vekua I.N., *Generalized Analytic Functions* (Pergamon Press, Oxford, 1962).

[4] Begehr H., and Hile G.N., A hierarchy of integral operators, Rocky Mountain J. Math., Vol.27, pp. 669-706, 1997.

[5] Begehr H., *Complex Analytic Methods For Partial Differential Equations*(World Scientific, Singapore, 1994)

[6] Begehr H., and Vaitekhovich T., Green functions in complex plane domains,Uzbek Math. J., Vol.4, pp. 29-34, 2008.

[7] Begehr H., Integral representations in complex, hypercomplex and Clifford analysis, Integral Transf. Special Funct., Vol.13, pp. 223-253, 2002.

[8] Beloglyadov M.V., and Burskii V.P., On solution uniqueness of the Dirichlet problem for a system of partial differential equations on one function in the unit ball, Journal of Inverse and Ill-Posed Problems Vol.19(1), pp. 13-23 doi: 10.1515/JIIP.2011.021, 2011.

[9] Volovich I.V., Sakbaev V.Z., Universal boundary value problem for equations of mathematical physics, Proceedings of the Steklov Institute of Mathematics, Vol.285(1), pp. 56-80 doi: 10.1134/S0081543814040063, 2014.

[10] Kalmenov T.Sh., Koshanov B.D., Nemchenko M.Y., Green function representation for the Dirichlet problem of the polyharmonic equation in a sphere, Complex Variables and Elliptic Equations, Vol.53(2), pp. 177-183, 2008.

## Crank-Nicolson difference scheme for the stochastic parabolic equation with the dependent operator coefficient

Allaberen ASHYRALYEV [1], Uelker OKUR [2]

[1] *Emeritus Professor, Turkey & Turkmenistan*
*E-mail: aallaberen@gmail.com*
[2] *Department of Mathematics, Faculty University of Stuttgart, Stuttgart, Germany*
*E-mail: uelkero@web.de*

**Abstract:** In the paper the Crank-Nicolson difference scheme for the numerical solution of the stochastic parabolic equation with the dependent operator coefficient is considered. Theorem on convergence estimates for the solution of this difference scheme is established. In applications, convergence estimates for the solution of difference schemes for the numerical solution of three mixed problems for parabolic equations are obtained. The numerical results are given.

**Keywords:** difference scheme, stochastic parabolic equation, convergence estimate

**2010 Mathematics Subject Classification:** 87.10.Ed, 43.35.Ei, 78.60.Mq

## Differential operator equations with interface conditions in modified direct sum spaces

Kadriye AYDEMİR[1], O. Sh. MUKHTAROV [2,3], Hayati OLĞAR[3]

[1] *Faculty of Education, Giresun University, Giresun, Turkey*
*E-mail: kadriyeaydemr@gmail.com*
[2] *Institute of Mathematics and Mechanics, Baku, Azerbaijan*
*E-mail: omukhtarov@yahoo.com*
[3] *Department of Mathematics, Gaziosmanpaşa University, Tokat, Turkey*
*E-mail: hayatiolgar@gmail.com*

**Abstract:** In this study we consider a Sturm-Liouville problem involving an abstract linear operator in equation and determined by eigendependent boundary conditions and two supplementary interface conditions. We give a characterisation of some spectral properties of the considered problem. Particularly, it is established such properties as isomorphism and coerciveness and find asymptotic formulas for eigenvalues.

**Keywords:** Differential operator equations, interface conditions, eigenvalues, isomorphism and coerciveness

**2010 Mathematics Subject Classification:** 34B4, 34L10

**References:**

[1] Aydemir K., and Mukhtarov O.Sh., "Green's Function Method for Self-Adjoint Realization of Boundary-Value Problems with Interior Singularities", *Abstract and Applied Analysis*, Vol.2013, Article ID 503267, 7 p., doi:10.1155/2013/503267, 2013.

[2] Bairamov E., and Uğurlu E., "The determinants of dissipative Sturm-Liouville operators with transmission conditions", *Math. Comput. Modelling*, Vol.53, No.5-6, pp. 805-813, 2011.

[3] Fulton C.T., "Two-point boundary value problems with eigenvalue parameter contained in the boundary conditions", *Proc. Roy. Soc. Edin.*, Vol.77A, pp. 293-308, 1977.

[4] Mukhtarov Oktay Sh., Olğar, Hayati and Aydemir, Kadriye, "Resolvent Operator and Spectrum of New Type Boundary Value Problems", *Filomat*, Vol.29:7, pp. 1671-1680, 2015.

[5] Titchmarsh E.C., "Eigenfunctions Expansion Associated with Second Order Differential Equations I", second edn. *Oxford Univ. Press*, London, 1962.

## Inverse problem for the Verhulst equation of limited population growth with discrete experiment data

Anvar AZIMOV [1], Syrym KASENOV [2], Daniyar NURSEITOV [3], Simon SEROVAJSKY [4]

[1] *Department of Differential Equations and Control Theory, al-Farabi Kazakh National University, Almaty, Kazakhstan*
*E-mail: anvar.aa@mail.ru*

[2] *Department of Methods of Teaching Mathematics, Physics and Computer Science, Abai Kazakh National Pedagogical University, Almaty, Kazakhstan*
*E-mail: syrym.kasenov@mail.ru*

[3] *National Open Research Laboratory of Information and Space Technologies, Kazakh National Research Technical University after Satpayev, Almaty, Kazakhstan*
*E-mail: nurseitovdb@gmail.com*

[4] *Department of Differential Equations and Control Theory, al-Farabi Kazakh National University, Almaty, Kazakhstan*
*E-mail: serovajskys@mail.ru*

**Abstract:** Verhulst limited growth model with unknown parameters of growth is considered. These parameters are defined by discrete experiment data. This inverse problem is solved with using gradient method with interpolation of data and without it [1, 2]. Approximation of the delta-function is used for the latter case. As an example the bacteria population *E.coli* is considered.

We consider the evolution of biological species in limited habitat. The given system is described by Verhulst equation. The coefficients of the equations are unknown. We would like to determine it from an inverse problem by using the results of the experiment. The data are distributed discretely. The inverse problem is transformed to an extremum one. This problem is solved by gradient method. The gradient of the given functional depends on the solution of the adjoint system. The adjoint equation includes delta-functions due to the discreteness of the data. This difficulty can be leave out by the interpolation of the data. This inverse problem is solved by using gradient method with interpolation of data and without it. The delta-function is approximated by Gauss formula [3]. The calculation are realized for the exact values of data and with noised data. The calculation accuracy is high enough for both cases and algorithms. We find also the parameters of growth for bacteria population E.coli by real experiment of Scientific Centre of Anti-infectious Drugs (Almaty).

**Keywords:** Verhulst equation, inverse problem, delta-function

**2010 Mathematics Subject Classification:** 34A34

**References:**

[1] Kabanikhin S. (2011). Inverse and Ill-posed Problems. Theory and Applications. Berlin, Boston: De Gruyter, 2011.

[2] Kabanikhin S.I., Shishlenin M.A., Nurseitov D.B. et al. Comparative analysis of methods for regularizing an initial boundary value problem for the Helmholtz equation // Journal of Applied Mathematics, Vol.2014, p. 7, 2014.

[3] Gel'fand I.M., Shilov G.E. Generalized Functions. Properties and Operations, 400p., 1980.

# A note on the nonlocal boundary value problem for a third order partial differential equation

Kheireddine BELAKROUM [1], Allaberen ASHYRALYEV [2], Assia GUEZANE-LAKOUD [3]

[1] *Department of Mathematics, Frères Mentouri University, Constantine, Algeria*
*E-mail: belakroumkheireddine@yahoo.com* [2] *Emeritus Professor, Turkey & Turkmenistan E-mail: aallaberen@gmail.com*
[3] *Laboratory of Advanced Materials Mathematics Department, Badji Mokhtar Annaba University, Annaba, Algeria*
*E-mail: a_guezane@yahoo.fr*

**Abstract:** It is known that various problem in fluid mechanics (dynamics, electricity) and other areas of physics lead to third order partial differential equations, we derive such equations as models of physical systems and consider methods for solving boundary value problems. This type of equations with constant coefficients can be solved using classical methods like Fourier transform method, and Laplace transform method. In the present study, the nonlocal boundary-value problem for a third order partial differential equation in Hilbert space with a self-adjoint positive definite operator is investigated. Applying operator approach, the theorem on stability for solution of this nonlocal boundary value problem is established. In applications, the stability estimates for the solution of three nonlocal boundary value problems for third order partial differential equations are obtained.

**Keywords:** Third order partial differential equation, Boundary value problems, Self-adjoint positive definite operator, Hilbert Space.

**2010 Mathematics Subject Classification:** 34G10, 47N20

**References:**

[1] Apakov Yu.P., Irgashev B.Yu., "Boundary-Value Problem for a Degenerate High-Odd-Order Equation", Ukrainian Mathematical Journal, Vol.66, Issue 10, pp. 1475-1490, 2015.

[2] Ashyralyev A., Aggez N., Hezenci F., " Boundary value problem for a third order partial differential equation", AIP Conf. Proc. 1470, pp. 130-132, 2012.

[3] Ashyralyev A., Sobolevskii P.E., New Difference Schemes for Partial Differential Equations,Birkh¨auser Verlag, Basel, Boston, Berlin, 2004.

## Solution of the free boundary problem for the parabolic equations with unknown temperature and velocity

Galina BIZHANOVA

*Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan*
*E-mail: galina_math@mail.ru*

**Abstract:** Let $\Omega(t)$ be a domain in $\mathbb{R}^n$, $n \geq 2$, $\gamma(t) := \partial\Omega(t)$, $t \in (0, t)$ and at the initial moment $t = 0$ $\Omega(0) := \Omega$, $\gamma(0) := \Gamma$. $\gamma(t)$ is a free (unknown) boundary.

We consider one–phase multidimensional free boundary problem for the parabolic equations in the unknown domain. Let the substance in $\Omega(t)$ be in a liquid state. The unknown functions determined in $\Omega(t)$ are temperature and velocity of liquid and they satisfy the parabolic equations. Such problem is a mathematical model, for instance, of the transportation of oil, when there is also a paraffined (solid) oil outside of $\Omega(t)$, and the boundary $\gamma(t)$ between the liquid and solid phases is unknown one.

The existence, uniqueness and coercive estimate of the solution of the considered problem locally in time is proved in the Hölder space.

**Keywords:** Hölder space, parabolic equation, free boundary problem, existence, uniqueness, coercive estimates

**2010 Mathematics Subject Classification:** 35R35, 35K20, 35B65

---

## On construction of solutions of linear fractional differential equations with constant coefficients

M.B.BORIKHANOV [1], B.Kh.TURMETOV [2]

[1] *Department of Mathematics, Akhmet Yasawi University, Turkestan, Kazakhstan*
*E-mail: meeir0808@gmail.com*
[2] *Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: turmetovbh@mail.ru*

**Abstract:** One of the effective methods for finding exact solutions of differential equations is the method based on the operator representation of solutions. The essence of this method is to construct a series, whose members are the relevant iteration operators acting to some classes of sufficiently smooth functions. This method is widely used in the papers of Bondarenko [1] for construction of solutions of differential equations of the integer order. In this paper, the operator method is applied to construct solutions of linear differential equations with constant coefficients and generalized Riemann-Liouville fractional derivative of order $\alpha$ and type $\gamma$. Then fundamental solutions are used to obtain the unique solution of the Cauchy problem.

**Keywords:** Linear fractional differential equations with constant coefficients, Riemann-Liouville derivatives, fundamental solutions, Cauchy problem

**2010 Mathematics Subject Classification:** 26A33, 34A08

**References:**

[1] Bondarenko B.A. , "Operator Algorithms in Differential Equations", *Tashkent, "FAN" Publishers*, 1984 (in Russian).

---

## More general common fixed point theorems under a new concept

Hakima BOUHADJERA [1]

[1] *Laboratory of Applied Mathematics, Badji Mokhtar-Annaba University, Annaba, Algeria*
*E-mail: b_hakima2000@yahoo.fr*

**Abstract:** A general common fixed point theorem for two pairs of weakly subsequentially continuous mappings (recently introduced) satisfying a significant estimated implicit function is proved. An extension of this result is thereby obtained. Our results assert the existence and uniqueness of common fixed points in several cases.

**Keywords**: subcompatible mappings, $R$-weakly commuting mappings of type $(\mathcal{A}_f)$ (resp. $(\mathcal{A}_g)$), sequentially continuous mappings of type $(\mathcal{A}_f)$ (resp. $(\mathcal{A}_g)$), weakly reciprocally continuous mappings, weakly subsequentially continuous mappings.

**2010 Mathematics Subject Classification:** 47H10, 54H25, 47H06, 47H09.

**References:**

[1] R.P. Agarwal, R.K. Bisht, N. Shahzad *A comparison of various noncommuting conditions in metric fixed point theory and their applications*, Fixed Point Theory and Applications (2014), 2014:38.

[2] M.A. Al-Thagafi, N. Shahzad *Generalized I-nonexpansive selfmaps and invariant approximations*, Acta Math. Sin. 23 (2007), no. 6, 965-972.

[3] H. Bouhadjera, C. Godet-Thobie *Common fixed point theorems for pairs of subcompatible maps*, arXiv: 0906.3159v1 [math. FA] 17 June 2009.

[4] H. Bouhadjera, C. Godet-Thobie *Common fixed point theorems for pairs of subcompatible maps*, arXiv: 0906.3159v2 [math. FA] 23 May 2011.

---

## Analytical treatment of two-center overlap integral with respect to same screening constants over Slater-type orbitals

Ebru ÇOPUROĞLU [1], Bahtiyar A. MAMEDOV [2]

[1] *Department of Physics, Gaziosmanpasa University, Tokat, Turkey*
*E-mail: ebru.copuroglu@gmail.com*

[2] *Department of Physics, Gaziosmanpasa University, Tokat, Turkey*
*E-mail: bamamedov@yahoo.com*

**Abstract:** By the use of Löwdin- $\alpha$ radial function, an alternative method is introduced for two-center overlap integral over Slater-type orbitals (STOs) at same screening constants (SSC). Notice that the overlap integrals with SSC have an important role in accurate evaluation of multicenter multielectron molecular integrals arising in combined Hartree-Fock-Roothaan (CHFR) theory. This approximation provides us simple and efficient expression for the two-center overlap integrals. To show the effectiveness of established formula we have compared our

results with available literature data [1]. The method ensures high performance with a very short CPU time.

**Keywords:** Löwdin- $\alpha$ radial function, Hartree-Fock-Roothaan (CHFR) theory

**PACS No:** 31.15.-p, 31.10.+z

**References:**

[1] Guseinov I.I., Mamedov B.A., Naturforsch Z., Vol.62a, pp. 467-470, 2007.

---

## Problems of the theory of the consolidation solved in the special functions

Azhibek DASIBEKOV[1], Azimkhan ABZHAPBAROV[1], Peruza DUISEBAYEVA[1], Aigul POLATBEK[1]

[1] *M. Auezov South Kazakhstan State University, Kazakhstan*
*E-mail: dasibekov@mail.ru, azeke_55@mail.ru, peruza_69@mail.ru, padisha66@mail.ru*

**Abstract:** The soil, which deformation modulus is continuously increased with depth is called continuously heterogeneous. In this paper, this heterogeneity is presented in the form of:

$$E = E_m(\alpha + \beta z)^m \quad (\alpha > 0, E_m > 0, \alpha + \beta z > 0),$$

where $E_m, \alpha, \beta, m$ are experimental parameters.

On the basis of this dependence the consolidation problems of elastic and elastically creeping inhomogeneous soils are solved in relation to the restricted area of the consolidation. These solutions make it possible to calculate the values of the pore pressure, the amount of the main stresses and vertical displacements of upper surface points of the condensed inhomogeneous soil mass.

In these solutions for highly compressed water-saturated clay soils is also taken into account that at the initial time the part of loading, instantly enclosed load q to the soil which is equal in value of the structural strength of the compression $p_{str}$, is immediately perceived by a matrix. In addition, Darcy's law is broken, i.e. the initial gradient of pressure is considered. The resulting calculation formulas are presented as a combination of Bessel functions of the first and second kinds. Taking into account that currently one can define any values of the Bessel functions, it is possible to calculate the pressure in the pore fluids and predict the speed of sediments of the compacting mass [1-4].

**Keywords:** soil, consolidation, stress, deformation modulus, pore pressure

**2010 Mathematics Subject Classification:** 74A10, 74L10

**References:**

[1] Klein G.K., *Hydraulic construction*, Vol.2, pp. 7-14, 1948 (in Russian).
[2] Popov G.Y., *Construction and architecture*, Vol.12, pp. 11-19, 1959 (in Russian).
[3] Dasibekov A., Yunusov A.A., Saidullaeva N.S., and Yunusova A.A., *International Journal of Experimental Education*, Vol.8, pp. 67-72, 2012 (in Russian).
[4] Dasibekov A., Yunusov A.A., Yunusova A.A., and Ayashova A., *Scientific journal "Basic Research"*, Vol.8, pp. 323-331, 2013 (in Russian).

## The near optimality of the stabilizing control in a weakly nonlinear system with state-dependent coefficients

Michael DMITRIEV [1], Dmitry MAKAROV [1]

[1] *FRC CSC RAS, Institute for Systems Analysis, Moscow, Russia*
*E-mail: mdmitriev@mail.ru, makarov@isa.ru*

**Abstract:** Currently State-Dependent Riccati Equation (SDRE) technique is quite a popular control approach for the nonlinear feedback control algorithm. However, it needs online solving the SDRE at every sample time that may require high performance computational resources for difficult control tasks. The new SDRE-based approach for constructing nonlinear stabilizing regulator was proposed in [1, 2] to overcome that problem. It deals with weakly nonlinear system with coefficients depending on the state and the formal small parameter. The proposed approach leads to computationally effective control because it uses analytical expressions. This paper is dedicated to near optimality of nonlinear stabilizing control constructed using that approach. First investigation of that problem was made in [3]. Now we use another form of optimal control and gain matrix. The theoretical results analogous to [3] are obtained and appropriate illustrative example of optimal control problem where formal small parameter equals to unit is considered. This work was supported by the Russian Science Foundation (Project No. 14-11-00692).

**Keywords:** weakly nonlinear systems, stabilizing control, small parameter, state-depended Riccati equations, optimal control.

**2010 Mathematics Subject Classification:** 49K40, 49K15, 49L20, 49M30, 93D15, 93D05

**References:**

[1] Dmitriev M.G., Makarov D.A., "Smooth nonlinear controller in a weakly nonlinear control system with state depended coefficients", *Trudy Instituta Sistemnogo Analiza Rossijskoj Akademii Nauk*, Vol.64, No.4, pp. 53-58, 2014 (in Russian).

[2] Emel'yanov S.V., Danik Yu.E., Dmitriev M.G., Makarov D.A, "Stabilization of Nonlinear Discrete-Time Dynamic Control Systems with a Parameter and State-Dependent Coefficients", *Doklady Mathematics*, doi: 10.1134/S1064562416010142, Vol.93, No.1, pp. 121-123, 2016.

[3] Dmitriev M.G., Makarov D.A., "The suboptimality of stabilizing regulator in a quasi-linear system with state-depended coefficients", *2016 International Siberian Conference on Control and Communications (SIBCON), Proceedings. Moscow: National Research University "Higher School of Economics"*. Russia, Moscow, May 12-14, 2016.

# Numerical solution of the hyperbolic-Schrödinger equation with the multipoint nonlocal boundary condition

Yildirim OZDEMIR [1], Sevilay ERDOGAN [2]

[1] *Department of Mathematics, Duzce University, Duzce, Turkey*
*E-mail: yildirimozdemir@duzce.edu.tr*
[2] *Graduate Institute of Sciences and Engineering, Duzce University, Duzce, Turkey*
*E-mail: svlyerdogan@hotmail.com*

**Abstract:** In this work, we suggest a numerical method to solve hyperbolic-Schrödinger partial differential equations with multipoint nonlocal boundary condition. The stability estimates for the solution of the given problem are established. The first and second order of accuracy difference schemes are obtained for the solution of the given problem. These difference schemes are solved by using the method of modified Gauss elimination for one-dimensional hyperbolic-Schrödinger partial differential equations. The results of numerical experiments are given for supporting the method.

**Keywords:** finite difference equation, partial differential equation, stability

**2010 Mathematics Subject Classification:** 65N06, 65N12

**References:**

[1] Ashyralyev A., Ozdemir Y., "Stability of difference schemes for hyperbolic-parabolic equations", *Comput. Math. Appl.*, Vol.50, pp. 1443-1476, 2005.

[2] Ashyralyev A., Ozdemir Y., "On nonlocal boundary value problems for hyperbolic-parabolic equations", *Taiwan. J. Math.*, Vol.11, No.4, pp. 1075-1089, 2007.

[3] Ozdemir Y., Kucukunal M., "On nonlocal boundary value problems for hyperbolic Schrödinger equations", *Abstr. Appl. Anal.*, Vol.2012, pp. 1-12, 2012.

[4] Ringhofet U.C., Koksal M.E., "Parameter extraction of complex pruduction systems via kinetic approach", *Kinet. Relat. Mod.*, Vol.9, No.2, pp. 407-427, 2016.

## Convergence of Modified Homotopy Perturbation Method for Fredholm-Volterra Integro-Differential Equation of order "m'

K. ESHKUVATOV [1], F. S. ZULKARNAIN [2],
Z. MUMINOV [3], N. M. A. NIK LONG, [4]

[1] *Faculty of Science and Technology, Universiti Sains Islam Malaysia (USIM), Ṅegeri Sembilan, Ṁalaysia*
*E-mail: zainidin@usim.edu.my*
[2,4] *Department of Mathematics, Faculty of Science, Universiti Putra Malaysia (UPM), Selangor, Malaysia*
*E-mail: fsamihah88@gmail.com, nmasri@upm.edu.my*
[3] *Faculty of Mathematics-Mechanics, Samarkand State University (SamSU), Samarkand, Uzbekistan*
*E-mail: zimuminov@gmail.com*

**Abstract:**

Homotopy perturbation method (HPM) [1, 2] is the combination of two methods: the homotopy and perturbation method. In recent years, the application of HPM in mathematical problems has been applied by many researchers. This method deforms complicated problem into a simple problem which is easy to solve.

HPM has been used for a wide range of problems; for finding the exact and approximate solutions of linear and nonlinear: ordinary differential equations(ODEs), Partial differential equations, integral equations and integro-differential equations of Volterra-Fredholm type.

In this note Modied homotopy perturbation method (MHPM) has applied to solve the Fredholm-Volterro integrodiffere-tial equation (FVIDE) of the third kind with the derivative of order m. Selective functions and unknown parameters helped us to obtain two step iterations. This proposed method could avoid complex computations. It is found that MHPM is a semianalytical method and easy to apply for FVIDE. Convergence and numerical examples are given to present the efciency and reliability of the method.

**Keywords:** Integral equation, Homotopy perturbation method, Numerical method

**2010 Mathematics Subject Classification:** 65R20, 45D05

**References:**

[1] He J.H., "Homotopy perturbation technique", Comp. Methods Appl. Mech. Engrg., Vol.178, pp. 257-262, 1999.
[2] He J.H., "A coupling method of a homotopy technique and a perturbation technique for non-linear problems", Int. J. Nonlinear Mech, Vol.35, pp. 37-43, 2000.

## Successive approximations to solve higher order fractional differential equations

Assia GUEZANE-LAKOUD[1], Rabah KHALDI[1]
[1] *Department of Mathematics, Badji Mokhtar Annaba University, Algeria*
*E-mail: a_guezane@yahoo.fr, rkhadi@yahoo.fr*

**Abstract:** In this talk, we discuss the existence and uniqueness of solution of an initial value problem for fractional differential equation involving higher order Caputo's fractional derivative. We transform the posed problem to a Volterra integral equation, then under Krasnoselskii-Krein type conditions and by using successive approximations, we discuss the existence and uniqueness questions.

**Keywords:** Fractional differential equation, Existence of solution.

**2010 Mathematics Subject Classification:** 34B15, 34B18, 34A12

**References:**

[1] Agarwal R.P., Meehan M., O'Regan D.," Fixed point theory and applications", Cambridge University, Press, 2001.
[2] Lakshmikantham V., Leela S., "A Krasnoselskii-Krein-type uniqueness result for fractional differential equations", Nonlinear Anal. Vol.71, pp. 3421-3424, 2009.
[3] Yoruk F., Bhaskar T.G., Agarwal R.P., "New uniqueness results for fractional differential equations", Appl. Anal. Vol.92, pp. 259-269, 2013.

## Operational matrices for finding numerical solution of stochastic differential equations arisen in financial mathematics

Farkhondeh HOSSEINI SHEKARABI [1], Hamed HATAMI [2]

[1] *Department of Mathematics, Yadegar-e-Imam Khomeini (Rah) Shahre Rey Branch, Islamic Azad University, Tehran, Iran*
*E-mail: f_hosseini@srttu.edu*

[2] *Department of Mathematics, Computer and Statistics, Faculty of Economics, Allameh Tabataba'i University, Tehran, Iran.*
*E-mail: Hamedhatami90@yahoo.com*

**Abstract:** In this paper, one of the new convenient techniques is used to solve problems formulated by stochastic Volterra integral equations. Cox-Ingersoll-Ross (CIR) model, Vasicek model and Heston model in financial mathematics can be transformed to a stochastic Volterra integral equation. Triangular functions, block pulse functions and their operational matrix and stochastic operational matrix of integration are considered. Properties of the operational matrices are utilized to reduce the computation of this equations to some algebraic equations. This method has several benefits; in addition to validity and good degree of accuracy, arithmetic operations are carried out without the need to derivative or integration, also, computation of presented method is very simple and attractive. The method is applied to a few test examples to illustrate the accuracy and implementation of the method.

**Keywords:** Orthogonal functions; Operational matrix; Stochastic operational matrix; Ito Integral; Brownian motion.

**2010 Mathematics Subject Classification:** 91G30, 91G60, 60H35, 65C20

**References:**

[1] Maleknejad K., Khodabin M., Hosseini Shekarabi F., Modified block pulse functions for numerical solution of stochastic Volterra integral equations, Journal of Applied Mathematics, Vol.2014, Article ID 469308, 10p., 2014.

[2] Khodabin M., Maleknejad K., Hosseini Shekarabi F., Application of Triangular Functions to Numerical Solution of Stochastic Volterra Integral Equations, IAENG International Journal of Applied Mathematics, Vol.43:1, IJAM-43-1-01, 2013.

# Problem with inhomogeneous boundary values for the equations of magnetic elektrogazodinamik

Djamila ISKENDEROVA [1], Aibek TOKTORBAEV [2]

[1] *International Academy of Management, law, finance and business, Bishkek, Kyrgyzstan*

[2] *Osh State University, Osh, Kyrgyzstan*

*E-mail:* [1] *iskenja_2005@mail.ru,* [2] *ain7@list.ru*

**Abstract:** Magnetic system of EGD equations in the Lagrangian mass coordinates is:

$$\frac{\partial v}{\partial t}-\frac{\partial u}{\partial x}=0,\quad v=\frac{1}{\rho},\quad \frac{\partial}{\partial t}vH=\frac{\partial}{\partial x}\Big(\frac{\mu_H}{v}\frac{\partial H}{\partial x}\Big),\qquad \frac{\partial E}{\partial t}=-bE\frac{\partial E}{\partial x},$$

$$\frac{\partial v}{\partial t}=\frac{\partial}{\partial x}\Big(\frac{\mu}{v}\frac{\partial u}{\partial x}\Big)-\frac{\partial p}{\partial x}-\mu_e H\frac{\partial H}{\partial x}+\varepsilon E\frac{\partial E}{\partial x},\quad p=r\frac{\theta}{v},\tag{2}$$

$$\frac{\partial \theta}{\partial t}=\frac{\partial}{\partial x}\Big(\frac{\lambda}{v}\frac{\partial \theta}{\partial x}\Big)-p\frac{\partial u}{\partial x}+\frac{\mu}{v}\Big(\frac{\partial u}{\partial x}\Big)^2+\frac{\mu_e\mu_H}{v}\Big(\frac{\partial H}{\partial x}\Big)^2+b\varepsilon vE^2\frac{\partial E}{\partial x}.$$

The unknown functions satisfy the boundary conditions:

$$\frac{1}{v}\frac{\partial u}{\partial x}-p\Big|_{x=0}=\frac{1}{v}\frac{\partial u}{\partial x}-p\Big|_{x=1}=H\Big|_{x=0}=H\Big|_{x=1}=0, E\Big|_{x=0}=\frac{\partial E}{\partial x}\Big|_{x=0}=0\tag{3}$$

$$\theta\Big|_{x=0}=\chi_0(t),\quad \theta\Big|_{x=1}=\chi_1(t),$$

$$\chi_i(t)\in W_2^1(0,T),\quad \chi_i(t)\geq m_0>0,\quad \chi_i(0)=\theta(i),\quad i=0,1.$$

At the initial time all the characteristics of the medium are known:

$$u\Big|_{t=0}=u_0(x), v\Big|_{t=0}=v_0(x), \theta\Big|_{t=0}=\theta_0(x), E\Big|_{t=0}=E_0(x), H\Big|_{t=0}=H_0(x),\tag{4}$$

**Theorem.** *Let*

$$(v_0,u_0,\theta_0,H_0,E_0)\in W_2^2(\Omega),\quad 0<m_0\leq(v_0(x),\theta_0(x))\leq M_0<\infty,\quad E_0'(x)\geq 0.$$

*Then, in area* $Q=\Omega\times(0,T)$ *with any finite* $T$ *there exists a unique generalized solution of problem (1) - (3), and*

$$(v(t),E(t))\in L_\infty(0,T,W_2^1(\Omega)),\quad (v_t,u_t,\theta_t,H_t,E_t)\in L_2(Q),$$

$$(u(t),\theta(t),H(t))\in L_\infty(0,T,W_2^1(\Omega))\cap L_2(0,T,W_2^2(\Omega)),\quad Q=\Omega\times(0,T),\Omega=(0,1)$$

$v(x,t),\theta(x,t)$ – *strictly positive, limited functions.*

**Keywords:** speed, density, temperature, magnetic field, electric field, generalized solution, apriori estimates, existence

**2010 Mathematics Subject Classification:** 70H03, 70H05

## Principle of Mass Conservation for the Boltzmann's Moment System of Equations in Fourth Approximation

Aizhan ISSAGALI [1], Auzhan SAKABEKOV [2]

[1] *Nazarbayev University, Astana, Kazakhstan*
*E-mail: aizhan.issagali@nu.edu.kz*
[2] *Kazakh-British Technical University, Almaty, Kazakhstan*
*E-mail: a.sakabekov@kbtu.kz*

**Abstract:** We study the one-dimensional non-linear non-stationary Boltzmann's moment system of equations in fourth approximation with the tools developed by Sakabekov in [1-2]. For the third approximation Sakabekov proves the mass conservation law (cf. Theorem in [2]). We consider the initial value problem for the system of hyperbolic partial differential equations in the conservative form. We extend the proof of the principle of mass conservation to the fourth approximation. The principle of mass conservation allows to prove the global existence in time of the solution of the system of nine hyperbolic partial differential equations representing the fourth approximation of the one-dimensional non-linear nonstationary Boltzmann's moment system of equations.

**Keywords:** Boltzmann equation, moment system, mass conservation, initial value problem, hyperbolic partial differential equations

**2010 Mathematics Subject Classification:** 35Q20, 35L65, 35L45

**References:**

[1] Sakabekov A., Initial and boundary value problems for the Boltzmann's moment system of equations. Almaty: Gylym, 2002.
[2] Sakabekov A., Auzhani Y., Analogue of the mass conservation law for third approximation of one-dimensional nonlinear Boltzmann's moment system equations. J. Math. Phys., Vol.54(5), 053512. doi:10.1063/1.4805363, 2013.

## On the well-posedness of the Boltzmann's moment system of equations in fourth approximation

Aizhan ISSAGALI [1], Auzhan SAKABEKOV [2]

[1] *Nazarbayev University, Astana, Kazakhstan*
*E-mail: aizhan.issagali@nu.edu.kz*
[2] *Kazakh-British Technical University, Almaty, Kazakhstan*
*E-mail: a.sakabekov@kbtu.kz*

**Abstract:** We study the one-dimensional non-linear non-stationary Boltzmann's moment system of equations in fourth approximation with the tools developed by Sakabekov in [1–3]. For the third approximation system Sakabekov proves the mass conservation law (cf. Theorem 2.1 in [2]) and discusses the existence and uniqueness of the solution (cf. Theorem in [3]). We extenISSAGALI 1, Auzhan SAKABEKOVd the analysis of the existence and uniqueness of the solution to the fourth approximation system. In particular, for the fourth approximation system we discuss the well-posed initial and boundary value problem and prove the existence and uniqueness of the solution belonging to the space of functions, continuous in time and square summable by spatial variable.

**Keywords:** Boltzmann equation, moment system, initial and boundary value problem, hyperbolic partial differential equations, a-priori estimate

**2010 Mathematics Subject Classification:** 35Q20, 35B45, 35L04

**References:**

[1] Sakabekov A., Initial and boundary value problems for the Boltzmann's moment system of equations. Almaty: Gylym, 2002.

[2] Sakabekov A., Auzhani Y., Analogue of the mass conservation law for third approximation of one-dimensional nonlinear Boltzmann's moment system equations. J. Math. Phys., Vol.54(5), 053512. doi:10.1063/1.4805363, 2013.

[3] Sakabekov A., Auzhani Y., Existence and uniqueness of solution of non-stationary Botzmann's moment system of equations in third approximation. TWMS J. Pure Appl. Math, Vol.5, pp. 36-42, 2014.

## On solvability of some boundary value problems for a biharmonic equation with periodic conditions

V.KARACHIK [1], S.MASANOV [2], B.Kh.TURMETOV [3]

[1] *Department of Mathematics,South Ural State University, Chelyabinsk, Russia Fatih University,*
*E-mail: valerykarachik@mail.ru*

[2] *Department of Mathematics, Akhmet Yasawi University, Turkestan, Kazakhstan*
*E-mail: meeir0808@gmail.com*

[3] *Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: turmetovbh@mail.ru*

**Abstract:** Let $\Omega = \{x \in R^n : |x| < 1\}$ be a unit ball, $n \geq 2, \partial\Omega$ be a unit sphere. For any point we compare its "opposite" point $x^* = (-x_1, ..., -x_n) \in \Omega$, and denote $\partial\Omega_+ = \partial\Omega \cap \{x \in R^n : x_n \geq 0\}$, $\partial\Omega_- = \partial\Omega \cap \{x \in R^n : x_n \leq 0\}$, $I = \partial\Omega \cap \{x \in R^n : x_n = 0\}$.

Consider the following problem in the domain $\Omega$:

$$\Delta^2 u(x) = f(x), x \in \Omega \tag{1}$$

$$D_\nu^m u(x) = g(x), x \in \partial\Omega \tag{2}$$

$$D_\nu^{\ell_1} u(x) - (-1)^k D_\nu^{\ell_1} u(x^*) = g_1(x), x \in \partial\Omega_+ \tag{3}$$

$$D_\nu^{\ell_2} u(x) + (-1)^k D_\nu^{\ell_2} u(x^*) = g_2(x), x \in \partial\Omega_+ \tag{4}$$

where $D_\nu^m = \frac{\partial^m}{\partial\nu^m}, m \geq 1$, $\nu-$ is a normal vector, $k = 1, 2,\ 1 \leq m \leq 3, 1 \leq \ell_1 < \ell_2 \leq 3, \ell_j \neq m, j = 1, 2$.

The exact conditions for solvability of the problems are found. Note, that analogous problems for Poisson equations were studied in [**?**] - [**?**].

**Keywords:** biharmonic equation, periodic boundary value problem, solvability, existence and uniqueness of solution

**2010 Mathematics Subject Classification:** 35J15, 35J25

**References:**

[1] Sadybekov M.A, Turmetov B.Kh, "On an analog of periodic boundary value problems for the Poisson Equation in the Disk", *Differential equations* , Vol.50, pp. 268–273, 2014.

[2] Sadybekov M.A, Turmetov B.Kh, "On analogues of periodic boundary value problems for the Laplace operator in a ball", *Eurasian Math. J.*, Vol.3, No.1, pp. 143-146, 2012.

## A conditional stability estimate of continuation problem for the Helmholtz equation

Syrym KASENOV [1], Altyn NURSEITOVA [2], Daniyar NURSEITOV [3]

[1] *Department of Methods of Teaching Mathematics, Physics and Computer Science, Abai Kazakh National Pedagogical University, Almaty*
*E-mail: syrym.kasenov@mail.ru*

[2] *Department of Methods of Teaching Mathematics, Physics and Computer Science, Abai Kazakh National Pedagogical University, Almaty*
*E-mail: altynna@mail.ru*

[3] *National Open Research Laboratory of Information and Space Technologies, Kazakh National Research Technical University after Satpayev, Almaty*
*E-mail: nurseitovdb@gmail.com*

**Abstract:** In this paper we consider the continuation problem for the Helmholtz equation. The main result is a conditional stability estimate for a solution to the considered problem. The estimate shows that the closer solution to the surface is more stable.

We consider the initial boundary value problem for the Helmholtz equation in the domain $\Omega = (0, l) \times (0, \pi)$ [1]:

$$u_{xx} + u_{yy} + k^2 u = 0, \qquad (x, y) \in \Omega, \tag{1}$$

$$u_x(0, y) = 0, \quad u(0, y) = f(y), \qquad y \in [0, \pi], \tag{2}$$

$$u_y(x, 0) = u_y(x, \pi) = 0, \qquad x \in [0, l], \tag{3}$$

where $k, l$ are given constants. It is required to find a function $u(x, y)$ in $\Omega$ from $f(y)$. The main theoretical result is the following

**Theorem.** Assume that for $f \in L_2(0, \pi)$ there exists a solution $u \in L_2(\Omega)$ to the problem (1) – (3), then the following estimate holds [2]:

$$\|u\|^2(x) \le \left(\|q\|^2 + \left|\|f_y\|^2 - \frac{k^2}{2}\|f\|^2\right|\right)^{\frac{x}{l}} \left(\|f\|^2 + \left|\|f_y\|^2 - \frac{k^2}{2}\|f\|^2\right|\right)^{\frac{l-x}{l}} e^{2x(l-x)} - \left|\|f_y\|^2 - \frac{k^2}{2}\|f\|^2\right|,$$

where $\|u\|^2(x) = \int_0^\pi u^2(x, y)dy$.

**Keywords:** Helmholtz equation, stability estimate, inverse problem

**2010 Mathematics Subject Classification:** 35R30

**References:**

[1] Kabanikhin S., Inverse and Ill-posed Problems. Theory and Applications. Berlin, Boston: De Gruyter, 2011.

[2] Lavrent'ev M.M., Savel'ev L.Y., *Operator Theory and Ill-Posed Problems*, De Gruyter, Germany, 680 p., 2011.

## Recurrence relation for the moments of order statistics from a beta-Pareto distribution

Hossein Jabbari KHAMNEI [1], Roghaye MAKOUYI [2]

[1] *University of Tabriz, Tabriz, Iran*
*E-mail: jabbari@tabrizu.ac.ir*
[2] *Tabriz University of Medical Sciences, Tabriz, Iran*
*E-mail: R.Makuyi@yahoo.com*

**Abstract:** In this paper, a novel cumulative distribution function (*c.d.f.*) for beta-Pareto ($BP$) distribution, through two distinct practical frames, is developed. However, the presented models are obviously more pragmatic than the ones being demonstrated in previous works, in the case of extending the further relations. Then, using the exhibited c.d.f.s, certain recurrence relations for the single and product moments of the order statistics of a random sample of size n arising from beta-Pareto distribution are derived.

**Keywords:** Order statistics, single and product moments, recurrence relations, beta-Pareto

**2010 Mathematics Subject Classification:** 62G30

**References:**

[1] Akinsete A., Famoye F., and Lee C., The beta-Pareto distribution. Statistics, Vol.42(6), pp. 547-563, 2008.

[2] Alshawarbeh E., Lee C., and Famoye F., The beta-Cauchy distribution. Journal of Probability & Statistical Science, Vol.10(1), pp. 41-57, 2012.

[3] Arnold B. C., Balakrishnan N., and Nagaraja H. N., A first course in order statistics, Vol.54, Siam, 1992.

[4] Cordeiro G. M., and Lemonte A. J., The $\beta$-BirnbaumSaunders distribution: an improved distribution for fatigue life modeling. Computational Statistics & Data Analysis, Vol.55(3), pp. 1445-1461, 2011.

[5] David H. A., and Nagaraja H. N., Order statistics. John Wiley & Sons, Inc., 1970.

[6] Eugene N., Lee C., and Famoye F., Beta-normal distribution and its applications. Communications in Statistics-Theory & methods, Vol.31(4), pp. 497-512, 2002.

[7] Famoye F., Lee C., and Olumolade O., The beta-Weibull distribution. Journal of Statistical Theory & Applications, Vol.4(2), pp. 121-136, 2005.

[8] Gradshteyn I. S. and Ryzhik I. M., Table of Integrals, Series, and Products. 7th Edition. Elsevier Inc., 2007.

[9] Jones M. C., Families of distributions arising from distributions of order statistics. Test, Vol.13(1), pp. 1-43, 2004.

[10] Henrick Malik J., Balakrishnan N., and Ahmed S. E., Recurrence relations mid identities for moments of order statistics, i: arbitrary continuous distribution. Communications in Statistics-Theory & Methods, Vol.17(8), pp. 2623-2655, 1988.

[11] Nadarajah S., and Gupta A. K., The beta Fréchet distribution. Far East Journal of Theoretical Statistics, Vol.14(1), pp. 15-24, 2004.

[12] Samuel P., and Thoma P. Y., An improved form of a recurrence relation on the product moments of order statistics. Communications in Statistics-Theory & Methods, Vol.29(7), pp. 1559-1564, 2000.

[13] Thomas P. Y., and Samuel P., A note on recurrence relations for the product moments of order statistics. Statistics & probability letters, Vol.29(3), pp. 245-249, 1996.

## The Motion of Fractional Order Jeffrey Fluid Model

Amir KHAN [1], Gul ZAMAN [2]

[1] *Department of Mathematics, University of Swat, Ṗakistan*
*E-mail: amir.maths@gmail.com*
[2] *Department of Mathematics, University of Malakand, Pakistan*
*E-mail: talash74@yahoo.com*

**Abstract:** This paper presents some new exact solutions corresponding to three unsteady flow problems of a generalized Jeffrey fluid [1-3] produced by a flat plate between two side walls perpendicular to the plate. The fractional calculus approach in the governing equations is used. The exact solutions are established by means of the Fourier sine transform and discrete Laplace transform. The series solution of velocity field and the associated shear stress in terms of Fox H-functions, satisfying all imposed initial and boundary conditions, have been obtained. The similar solutions for ordinary Jeffrey fluid, performing the same motion, appear as limiting case of the solutions are obtained here. Also, the obtained results are analyzed graphically through various pertinent parameter.

**Keywords:** Generalized Jeffrey fluid, Fractional derivatives, Fox H-functions, Discrete Laplace transform.

**2010 Mathematics Subject Classification:** 76A05, 76A10

**References:**

[1] Hayat T., Khan M., Fakhar K., Amin N., Oscillatory rotating flows of a fractional Jeffrey fluid filling a porous space. J Porous Media, Vol.13(1), pp. 29-38, 2010.
[2] Qi H., Xu M., Stokes' first problem for a viscoelastic fluid with the generalized Jeffrey model. Acta Mech Sin, Vol.23, pp. 463-469, 2007.
[3] Khadrawi A.F., Al-Nimr M.A., Othman A., Basic viscoelastic fluid Problems using the Jeffreys model. Chem Eng Sci, Vol.60, pp. 7131-7136, 2005.

## Computational algorithms for analysis of data from thin-film thermoresistors on a radio-electronic printed circuit board

Anna KORNEEVA [1], Vladimir SHAIDUROV [2]

[1] *Siberian Federal University, Krasnoyarsk, Russia*
*E-mail: korneeva_ikit@mail.ru*
[2] *Institute of Computational Modelling, Russian Academy of Sciences, Krasnoyarsk, Russia*
*E-mail: shaidurov04@mail.ru*

**Abstract:** In the paper, a printed circuit board with radio-electronic elements is considered as a geometrical structure with thin-film thermoresistors coated on to it. The local resistance of these thermoresistors depends on local temperature. The measurement of these resistances is carried out not in a local way but totally along horizontal or vertical lines. Therefore, the number of these total data on resistance is significantly less than the number of these local zones. During operating the printed circuit board, overheating of separate elements may occur that leads to a temperature increase of its corresponding parts and a resistance increase of separate parts of a chain. We set up the task of determining these areas of overheating and also their resistance and temperature being guided by some additional physical conditions.

**Keywords:** thin-film thermoresistors, radio-electronic printed circuit boards, local temperature, local resistance, numerical algorithms, underdetermined systems of linear algebraic equations

**2010 Mathematics Subject Classification:** 65F05, 80A23.

**References:**

[1] J.H. Wilkinson. The Algebraic Eigenvalue Problem. *A Clarendon Press Publication. Oxford.* (1988).

# On solvability of boundary value problem of magnetic gas dynamics with cylindric and spherical symmetries

Bakytbek D. KOSHANOV [1], Gulzhazira D. SMATOVA [2]

[1,2] *Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail:* [1] *koshanov@list.ru*

**Abstract:** In this paper we investigate the initial-boundary value problems for the equations motion of a viscous heat-conducting gas, taking into account the magnetic field. The main attention is paid to the case of solutions with cylindrical and spherical symmetry. The paper describes in detail the transition from the basic three-dimensional equations to cylindrical and spherical coordinate system and prove a number of theorems on the existence and uniqueness of the initial-boundary value problems.

$$\frac{\partial \rho}{\partial t}+\rho^2\frac{\partial (x^m u)}{\partial q}=0,\ P=R\rho\theta,$$

$$\frac{\partial u}{\partial t}+\frac{H^2}{x\rho}=x^m\frac{\partial}{\partial q}\left(\rho\frac{\partial (x^m u)}{\partial q}\right)-x^m\frac{\partial P}{\partial q}-x^m H\frac{\partial H}{\partial q},$$

$$\frac{\partial \theta}{\partial t}=æ\frac{\partial}{\partial q}\left(x^{2m}\rho\frac{\partial \theta}{\partial q}\right)+\rho\left[\frac{\partial (x^m u)}{\partial q}\right]^2+$$

$$+\rho x^{2(m-1)}\left[\frac{\partial (xH)}{\partial q}\right]^2-P\frac{\partial (x^m u)}{\partial q}-\frac{3m}{2}\frac{\partial (x^{m-1}u^2)}{\partial q}, \tag{1}$$

$$\frac{\partial H}{\partial t}+x\rho H\frac{\partial (x^{m-1}u)}{\partial q}=x\rho\frac{\partial}{\partial q}\left(\rho x^{2(m-1)}\frac{\partial (xH)}{\partial q}\right),$$

$$(u,\rho,\theta,H)(q,0)=(u_0,\rho_0,\theta_0,H_0)(q), q\in[0,b], \tag{2}$$

$$(u,\frac{\partial \theta}{\partial q},H)(0,t)=(u,\frac{\partial \theta}{\partial q},H)(b,t)=0, t\in[0,T]. \tag{3}$$

This article explores the initial-boundary value problems (1)-(3) for the equations motion of a viscous heat-conducting gas, taking into account the magnetic field with cylindrical and spherical symmetry. The paper describes in detail the transition from the basic three-dimensional equation in cylindrical and spherical coordinates. Under certain conditions the data (2) prove a number of theorems on the existence and uniqueness of the initial-boundary value problems.

**Keywords:** gas dynamics equation, boundary value problems, spherical symmetry

**2010 Mathematics Subject Classification:** 39A10, 35J66, 35J67

## Diffusion equation model for the tumors cell density and immune response

Sanjeev KUMAR [1], Sanjay CHAUDHARY [1]

[1] *Dept. of Mathematics, Dr. B.R. Ambedkar University, Agra, India*
*E-mail: sanjeevibs@yahoo.co.in*

**Abstract:** Over the last 20 years increasingly complex mathematical models of cancerous growths have been developed, especially on solid tumors, in which growth primarily comes from cellular proliferation. Consider a procedure for cancer therapy which consists of interaction between immune response (immune cells) and tumor cells without any specific drug. The cytotoxic T lymphocyte (CTL) and tumor necrosis factor (TNF) cause of the immune response. This process is modeled as a system of tumor cell density (TCD) and tumor necrosis factor (TNF). The purpose of this paper is to establish a rigorous mathematical analysis of the model and to explore the density/concentration of tumor cell and immune response (TNF). The result suggests that although TCD capable to growth of tumor but the immune response is block to direct tumor growth. The model assumes that only two factors need be considered for such predictions: net growth rate and infiltrative ability. The model has already provided illustrations of theoretical glioblastomas, but also shows the distribution of the diffusely infiltrating cell.

---

## Investigation of Laplace transforms for Erlangen distribution of the first passage time into zero level of the semi-Markov random process with positive tendency and negative jumps

Selahattin MADEN[1], Tamilla I. NASIROVA[2], Ulviyya Y. KARIMOVA[2]

[1] *Department of Mathematics, Faculty of Arts and Sciences, Ordu University, 52200, Ordu, Turkey*
*E-mail: maden55@mynet.com*
[2] *Department of Applied Mathematics and Cybernetics, Baku State University, Baku, Azerbaijan*
*E-mails: nasirova.tamilla@mail.ru, kerimova_ulviyye@yahoo.com*

**Abstract:** The investigations of the distributions for the processes of semi-Markov random process have an important value in the random process theory. The purpose of the present article is to find the Laplace transforms for Erlangen distribution of the semi-Markov random processes with positive tendency and negative jumps. Let's the sequence $\{\xi_k, \zeta_k\}_{k\geq 1}$ be given on the probability space $\{\Omega, \Im, P(\cdot)\}$, where the random variables $\xi_k$ and $\zeta_k$, $k \geq 1$ independent and identically distributed. Using these random variables we construct the following semi-markov random process:

$X(t) = z + t - \sum_{i=1}^{k-1} \zeta_i$, if $\sum_{i=1}^{k-1} \xi_i \leq t < \sum_{i=1}^{k} \xi_i$, $k = 1, 2, ...,$ $t, z \geq 0$.

$X(t)$ process is called as the semi-markov random process with positive tendency and negative jumps. Also let us denotethe random variable $\tau_z^0$ the first falling time of the process into zero-level, as $\tau_z^0 = \min\{t : X(t) \leq 0\}$. We need to find the Laplace transform for distribution of random variable $\tau_z^0$. Let's assume that the random variables $\xi_1$ and $\zeta_1$ have an Erlangen distribution of third construction with the parameters $\lambda$ and $\mu$, respectively. According to total

probability formula, we can write the following integral equation for the random variable $\tau_z^0$ :

$$L(\theta \mid z) = \frac{\lambda^3 e^{-\mu z}}{(\lambda+\mu+\theta)^3} + \frac{\lambda^3 \mu e^{-\mu z}}{2} \int_{s=0}^{\infty} s^2 e^{-(\lambda+\mu+\theta)s} \int_{\alpha=0}^{z+s} e^{\mu\alpha} L(\theta \mid \alpha) d\alpha ds.$$

In this case, we get the following fourth order differential equation:

$$\begin{aligned} L^{(4)}(\theta \mid z) &- [3(\lambda+\theta)-\mu]L'''(\theta \mid z) + 3(\lambda+\theta)(\lambda+\theta-\mu)L''(\theta \mid z) \\ &- (\lambda+\theta)^2(\lambda+\theta-3\mu)L'(\theta \mid z) - \mu[(\lambda+\theta)^2 - \lambda^3]L(\theta \mid z) = 0. \end{aligned}$$

Then the general solution of this differential equation will be as follows:

$$L(\theta \mid z) = \frac{\lambda^3}{[\lambda+\theta-k_1(\theta)]^3} e^{k_1(\theta)z}.$$

This expression is the Laplace transform for relative distribution of random variable $\tau_z^0$. Laplace transform for non-relative distribution of $\tau_z^0$ will be

$$L(\theta) = \int_{z=0}^{\infty} L(\theta \mid z)\lambda^3 z^2 e^{-\lambda z} dz =$$

$$\int_{z=0}^{\infty} c_1(\theta) e^{k_1(\theta)z} \lambda^3 z^2 e^{-\lambda z} dz = \frac{\lambda^3}{[\lambda - k_1(\theta)]^3} C_1(\theta).$$

Therefore, we will get the following characteristics for $\lambda m > \mu$ :

$$\begin{aligned} E\tau_z^0 &= -L'(0) = \frac{3(\lambda+\mu)}{\lambda(\lambda-3\mu)}, \\ E(\tau_z^0 \mid z) &= \frac{3(1+z\mu)}{\lambda-3\mu}, \end{aligned}$$

and

$$D(\tau_z^0 \mid z) = \frac{3}{(\lambda-3\mu)^2} + \frac{12(3+z\lambda)\mu}{(\lambda-3\mu)^3}.$$

**Keywords:** Laplace transform, semi-Markov random process, random variable.

**References:**

[1] Borovkov A.A., "On the random walk in the strip with two delaying barriers", *Math. Zametki*, Vol.17, No.5, pp. 647-657, 1975.

[2] Borovkov A.A., Theory of Probabilities, Moscow: Nauka, 2003.

[3] Gikhman I.I., and Skorokhod A.V., Theory of Accidental Processes, Moscow: Nauka, 1973.

[4] Khaniev T.A., "Distribution of a semi-Markov Walk with two delay screens, Some question of the theory of stochastic processes", *Collect sci. Works*, Kiev, pp. 106-113, 1984.

[5] Khaniev T.A., Ünver I., "The study of the level zero crossing time of a semi-Markovian random walk with delaying screen", *Tr. J. of Mathematics*, Vol.21, pp. 257-268, 1997.

[6] Nasirova T.I., and Sadikova R.I., "Laplace Transformation of the Distribution of the Time of System Sojourns within a Band", *Automatic Control and Computer Sciences*, Vol.43, No.4, pp. 190-194.

[7] Nasirova T.I., and Kerimova U.Y., "Definition of Laplace Transformsfor Distribution of the First Passage of Zero Level of the Semi-Markov Random Process with Positive Tendency and Negative Jump", *Appl. Math*, No.2, pp. 908-911, 2011.

## **Statistical inference for the parameter of Burr-type** 2 model using vague information as fuzzy data

Iman MAKHDOOM [1], Parviz NASIRI [1], Abbas PAK [2], Masoud YARMOHAMADI

[1] *Department of Statictics, Payame Noor University (PNU), P.O.BOX, 19395-3697, Tehran, Iran*
*E-mail: makhdoom@pnu.ac.ir*

[2] *Department of Computer Sciences, Faculty of Mathematical Sciences, Shahrekord University, P.O.Box 115,Shahrekord, Iran*
*E-mail: abbas.pak1982@gmail.com*

**Abstract:** Conventional statistical analysis of lifetime models is based on precise data. However, in real world situations, some information about an underlying system might be vague and are represented in the form of fuzzy information. Thus, it is necessary to generalize classical statistical estimation methods for real numbers to fuzzy numbers. Pak et al. ( [1], [2], [**?**], [3]) conducted a series of studies to develop the inferential procedures for the lifetime models on the basis of fuzzy numbers. In this study, different methods of estimation are discussed for the parameter of Burr-type 2 when the available data are in the form of fuzzy numbers. For this, we use the numerical methods such as "Newton-Raphson method" when the available observations are described by means of fuzzy information. Furthermore, we present a Bayesian approach to estimate the parameter of the model. Finally, Comparisons are made between these estimators using Monte-Carlo simulation study.

**Keywords:** Maximum Likelihood Estimate, Markov chain Monte Carlo, EM algorithm, Fuzzy data analysis

**2010 Mathematics Subject Classification:** 94B05, 94B15

**References:**

[1] Pak A., Parham G.H., and Saraj M., "Inference for the Weibull Distribution Based on Fuzzy Data", *Revista Colombiana de Estadistica*, Vol.36(2), pp. 339-358, 2013.

[2] Pak A., Parham G.H., and Saraj M., "Inference for the Rayleigh Distribution Based on Progressive Type-II Fuzzy Censored Data". *Journal of Modern Applied Statistical Methods*, Vol.13(1), 287-304, 2014. sed on Fuzzy Data". *IEEE Transactions on Reliability*, Vol.63(1), pp. 1-10, 2014.

[3] Pak A., Parham G., Saraj M., "Reliability estimation in Rayleigh distribution based on fuzzy lifetime data". *Int. J. Syst. Assur. Eng. Manag.* Vol.5(4), pp. 487-494, 2013.

# Differentiation of the functional in an optimization problem for diffusion and convective transfer coefficients of elliptic imperfect contact interface problems

Aigul MANAPOVA [1] [1]

[1] *Department of Mathematics and IT, Bashkir State University, Ufa, Republic of Bashkortostan, Russia*
*E-mail: aygulrm@mail.ru*

**Abstract:** Convection-diffusion problems are typical for mathematical models of liquid and gas mechanics, since heat and impurities transfer can occur not only due to diffusion, but also to the motion of the medium (see [1]). Currently, the most profound results in the theory of numerical solution of problem for PDEs and optimal control problems are obtained for processes with self-adjoint operators.

In this work we investigate issues of numerical solving of optimal control problems for second order elliptic equations with non-self-adjoint operators - convection-diffusion problems. Control processes are described by semi-linear convection-diffusion equation with discontinuous data and solutions (states) subject to the boundary interface conditions of imperfect type (i.e., problems with a jump of the coefficients and the solution on the interface; the jump of the solution is proportional to the normal component of the flux). Controls are involved in the coefficients of diffusion and convective transfer.

The subject of this paper is related to [2]. We prove differentiability and Lipshitz continuity of the cost functional, depending on a state of the system and a control. The calculation of the gradients uses the numerical solutions of direct problems for the state and adjoint problems.

**Keywords:** optimal control, semi-linear elliptic equation, imperfect contact, convective transfer, cost functional, differentiability

**2010 Mathematics Subject Classification:** 49J20, 35J61, 65N06

**References:**

[1] Samarskii A.A., Vabishchevich P.N., "Computational Heat Transfer", Wiley, New York, 1996; Librokom, Moscow, 2009.

[2] Manapova A.R., Lubyshev F.V., "On frechèt differentiability of cost functional in optimal control of coefficients of elliptic equations", *Ufa Mathematical Journal*, Vol.8, No.1, pp. 84-102, 2016.

[1]This work was supported by a grant of the President of Russian Federation for state support of young Russian scientists" Ph.D. (MK-4147.2015.1).

## Ranking DNA words Using Nonextensive Statistical Mechanics

H. MOGHADDASI [1], K. KHALIFEH [2], A.H.DAROONEH [1]

[1] *Department of Physics, Zanjan University, Zanjan, Iran*

[2] *Department of Biology, Zanjan University, Zanjan, Iran*

*E-mail: h.moghaddasi.ac.ir*

*E-mail: khalifeh.ac.ir*

*E-mail: darooneh@znu.ac.ir*

**Abstract:** Functional DNA subsequences and genome elements are spatially clustered through the genome just as keywords in literary texts. Therefore; some of the methods for ranking words in texts can also be used to compare different DNA subsequences. One of these methods is based on nonextensive statistical mechanics [1]. Here we claim that the distribution of distances between the successive subsequences (words) has q-exponential form which is the Tssalis distribution function in nonextensive statistical mechanics [2].Thus we introduce the q-parameter as a measure for estimating the clustering level of words. In this study, we analyzed the distribution of distances between consecutive occurrences of 16 possible dinucleotides in human chromosomes by comparing their corresponding q-parameters and found that the CGs have the highest clustering level which is a biologically important word concerning its methylation. We extended our study by comparing the genome of some selected organisms and concluded that the clustering level of CG dinucleotides increases in higher evolutionary organisms compared to lower ones

**Keywords:** Statistical Mechanics, DNA sequences, q-exponential function

**References:**

[1] Mehri A. & Darooneh A.H., "Keyword extraction by nonextensivity measure. *Phys. Rev. E.*, Vol.83, pp. 1-6, 2011.

[2] Tsallis C., "Introduction to Nonextensive Statistical Mechanics", *Springer*, ISBN. 978-0-387-85358-1, 2009

## Positiveness of second order differential operators with interior singularity

O. Sh. MUKHTAROV [1,2], Hayati OLĞAR [1], F. S. MUHTAROV [2,3]

[1] *Department of Mathematics, Gaziosmanpaşa University, Tokat, Turkey*
*E-mail: omukhtarov@yahoo.com*
[2] *Institute of Mathematics and Mechanics, Baku, Azerbaijan*
*E-mail: hayatiolgar@gmail.com*
[3] *Mechanics-Mathematics Faculty, Baku State University, Baku, Azerbaijan*
*E-mail: aliriyaziyyat@mail.ru*

**Abstract:** The main goal of this study is to provide an operator-theoretic framework for the investigation of discontinuous Sturm–Liouville problems with eigenparameter appearing in the boundary conditions. We introduce some self-adjoint compact operators in suitable Sobolev spaces such a way that the considered problem can be reduced to an operator-pencil equation. We define a new concept so-called generalized eigenfunctions and prove positiveness of operator-pencil corresponding to the considered problem.

**Keywords:** Sturm-Liouville problem, operator-pencil, interior singularity

**2010 Mathematics Subject Classification:** 34B4, 34L10

**References:**

[1] Aydemir K., and Mukhtarov O., "Variational principles for spectral analysis of one Sturm-Liouville problem with transmission conditions", *Advances in Difference Equations*, 2016:76, 2016.

[2] Gohberg I.C., Krein M.G., "Introduction to The Theory of Linear Non-Selfadjoint Operators", Translation of Mathematical Monographs, vol. 18, *Amer. Math. Soc., Providence*, Rhode Island, 1969.

[3] Keldys M.V., "On completeness of the eigenfunctions for certain classes of non-self-adjoint linear operators", *Upsekhi Mat. Nauk (in Russian)*, 27, 1971, No.4(160), pp. 15-41; *English transl. in Russian Math. Surveys*, 27, 1971.

[4] Ladyzhenskaia O.A., "The Boundary Value Problems of Mathematical Physics", *Springer-Verlag*, New York, 1985.

[5] Mukhtarov O.Sh., Olğar H., and Aydemir K., "Resolvent Operator and Spectrum of New Type Boundary Value Problems", *Filomat*, 29:7 , pp. 1671-1680, 2015.

[6] Mukhtarov O.Sh., and Kandemir M., "Asymptotic behaviour of eigenvalues for the discontinuous boundary- value problem with Functional-Transmissin conditions", *Acta Mathematica Scientia*, Vol.22 B(3), pp. 335-345, 2002.

## Transmission conditions for biharmonic equations with discontinuous coefficients and numerical solution of boundary value problem

Zahir MURADOGLU

*Department of Mathematics, Kocaeli University, Kocaeli, Turkey*
*E-mail: zahir@kocaeli.edu.tr*

**Abstract:** The mathematical model of bending problem for the plate system which is composed of inhomogeneous elastic plates with differing elastic properties is presented by boundary value problems for the biharmonic differential equations of with discontinuous coefficients. Providing continuity at the common borders of the plates, determination of transmission conditions among them and the development of methods for finding suitable numerical aproximate solution for the problem keep up to date for applied mathematics.

The fundamental aim of this study is to obtain the transmission condition of plates system composed of elastic plates with differing elastic properties, their difference equations and then the numerical solution at different boundary conditions. Firstly, we can represent the potantial energy of plates system. Later, finite difference approximations of transmission condition at common boundary of plates is obtained by functional approximation method with the help of numerical approximation of derivative and integral.

In this paper, we obtain numerical solution of biharmonic equation with discontinuous coefficient at different boundary conditions and the analysis of numerical results is given.

**Keywords:** biharmonic equation,transmission condition

**2010 Mathematics Subject Classification:** 65N06, 74G15, 35J40

**References:**

[1] Samarski A.A., Andreev V.B., "Difference methods for elliptic equations", *Nauka*, Moscow, 1976.

## Some properties of eigenvalues and generalized eigenvectors of one boundary-value problem

Hayati OLĞAR [1], O. Sh. MUKHTAROV [1,2], Kadriye AYDEMİR [3]

[1] *Department of Mathematics, Gaziosmanpaşa University, Tokat, Turkey*
*E-mail: hayatiolgar@gmail.com*
[2] *Institute of Mathematics and Mechanics, Baku, Azerbaijan*
*E-mail: omukhtarov@yahoo.com*
[3] *Faculty of Education, Giresun University, Giresun, Turkey*
*E-mail: kadriyeaydemr@gmail.com*

**Abstract:** We investigate one discontinuous boundary value problem which consist of a Sturm-Liouville equation with piecewise continuous potential together with eigenparameter-dependent boundary conditions and supplementary transmission conditions. We establish some spectral properties of the considered problem. In particular it is shown that the generalized eigenfunctions form a Riesz basis of the adequate Hilbert space.

**Keywords:** Discontinuous boundary value problem, boundary conditions, Riesz Basis

**2010 Mathematics Subject Classification:** 34B4, 34L10

**References:**

[1] Aydemir K., "Boundary value problems with eigenvalue depending boundary and transmission conditions", *Boundary value problems*, 2014:131, 2014.

[2] Kerimov N.B., and Aliev Z.S. "On the Basis Property of the System of Eigenfunctions of a Spectral Problem with Spectral Parameter in a Boundary Condition", *Differ. Uravn.*, Vol.43(7), pp. 886-895, 2007.

[3] Ladyzhenskaia O.A., "The Boundary Value Problems of Mathematical Physics", *Springer-Verlag*, New York, 1985.

[4] Mukhtarov O.Sh., Olğar H., and Aydemir K., "Resolvent Operator and Spectrum of New Type Boundary Value Problems", *Filomat*, Vol.29:7 , pp. 1671-1680, 2015.

[5] Mukhtarov O.Sh., and Yakubov S., "Problems for ordinary differential equations with transmission conditions", *Appl. Anal.*, Vol.81, pp. 1033-1064, 2002.

# On a method of finding a solution of semi-periodic boundary value problem for hyperbolic equations

Nurgul ORUMBAYEVA [1,2], Beybitgul SHAYAKMETOVA [3]

[1] *Institute of Applied Mathematics, Karaganda, Kazakhstan*
[2] *Department of Mathematics, Karaganda State University n.a. E.A. Buketov, Karaganda, Kazakhstan*
*E-mail: orumbayevan@mail.ru*
[3] *Department of Mathematics, Karaganda State University n.a. E.A. Buketov, Karaganda, Kazakhstan*
*E-mail: kazahzavod@mail.ru*

**Abstract:** In this paper we investigate the floor semi-periodic boundary value problem for a system of hyperbolic equations with mixed derivative. An algorithm for finding an approximate solution of this problem.

In this paper, a semi-periodic boundary value problem for a system of linear hyperbolic equations is considered:

$$\frac{\partial^2 u}{\partial x \partial t} = A(x,t)\frac{\partial u}{\partial x} + C(x,t)u + f(x,t), \quad (x,t) \in \overline{\Omega} = [0,\omega] \times [0,T],$$

$$u(0,t) = \psi(t), \quad u(x,0) = u(x,T), \quad t \in [0,T], \quad x \in [0,\omega],$$

where $A(x,t), C(x,t)$ matrix of $(n \times n)$ order, $f(x,t)$ $n$-vector-function in continuous $\overline{\Omega}$, $n$-vector-function $\psi(t)$ continuously-differentiable in $[0,T]$. For her study, the interval $[0,\omega]$ is divided into M parts, and parameterization method is applied for each area. The proposed method for finding approximate solutions of semi-periodic boundary value problem for a system hyperbolic equations is illustrated by an example.

**Keywords:** Hyperbolic equations, decomposition, algorithm, the boundary value problem

**2010 Mathematics Subject Classification:** 35K20, 44A10, 45D05, 35L20

**References:**

[1] Assanova A.T., Dzhumabaev D.S., "Correct solvability of nonlocal boundary value problems for systems of hyperbolic equations", *Differential equations*, Vol.41, No.3, pp. 337-346, 2005.

[2] Orumbayeva N.T., "On an algorithm of finding periodical boundary value problem for system of the quasi-linear of hyperbolic equations", *Siberian Electronic Mathematical Reports*, Vol.10, pp. 464-474, 2013.

[3] Dzhumabaev D.S., "Criteria for the unique solvability of a linear boundary-value problem for an ordinary differential equation", *Zhurnal Vychislitel'noi Matematiki i Matematicheskoi Fiziki*, Vol.29, No.1, pp. 50-66, 1989.

## A note on numerical solution of the parabolic-Schrödinger equation

Yildirim OZDEMIR [1], Mustafa ALP

[1] *Department of Mathematics, Duzce University, Duzce, Turkey*
*E-mail: yildirimozdemir@duzce.edu.tr, alp.mustafa81@gmail.com*

**Abstract:** The nonlocal boundary value problem for a parabolic-Schrödinger equation is considered. The stability estimates for the solution of the given problem are established. For aproximately solving this nonlocal boundary problem the first and second order of difference schemes are presented. The theoretical statements for the solution of these difference schemes are supported by the result of numerical examples.

**Keywords:** finite difference equation, partial differential equation, stability

**2010 Mathematics Subject Classification:** 65N06, 65N12

**References:**

[1] Ashyralyev A., Fattorini H. O., "On uniform difference scheme for second order singular perturbation problems in Banach spaces", *SIAM J. Math. Anal.*, Vol.23, No.1, pp. 29-54, 1992.

[2] Ashyralyev A., Ozdemir Y., "Stability of difference schemes for hyperbolic-parabolic equations", *Comput. Math. Appl.*, Vol.50, pp. 1443-1476, 2005.

[3] Ashyralyev A., Ozdemir Y., "On nonlocal boundary value problems for hyperbolic-parabolic equations", *Taiwan. J. Math.*, Vol.11, No.4 pp. 1075-1089, 2007.

[4] Ozdemir Y., Kucukunal M., "On nonlocal boundary value problems for hyperbolic Schrödinger equations", *Abstr. Appl. Anal.*, Vol.2012, pp. 1-12, 2012.

## Well-posedness of high order of accuracy difference scheme

Elif OZTURK BEIGMOHAMMADI

*Department of Economics, Canakkale Onsekiz Mart University, Canakkale, Turkey*

*E-mail: elifozturk16@hotmail.com*

**Abstract:** The simply nonlocal boundary value problem was investigated for the first time by A.V. Bitsadze and A.A. Samarskii in the paper [1]. In recent years, [3]– [5], the Bitsadze–Samarskii type nonlocal boundary value problem and its generalizations for various differential and difference elliptic equations have been studied. In this work, the Bitsadze-Samarskii type nonlocal boundary-value problem with the integral condition

$$\begin{cases} -\frac{d^2u(t)}{dt^2} + Au(t) = f(t),\ 0 < t < 1, \\ u(0) = \varphi,\ u(1) = \int\limits_0^1 \rho(\lambda)u(\lambda)d\lambda + \psi \end{cases} \tag{1}$$

for the differential equation of elliptic type in a Hilbert space $H$ with the self-adjoint positive definite operator $A$ with a closed domain $D(A) \subset H$ is considered. The well-posedness of the fourth order of accuracy difference scheme for approximate solution of nonlocal boundary-value problem (1) in Holder spaces with a weight is proved. The theoretical statements for the solution of this difference scheme are supported by the results of a numerical example.

**Keywords:** elliptic equation, nonlocal boundary value problem, well-posedness

**2010 Mathematics Subject Classification:** 35J40, 34B10, 49K40

**References:**

[1] Bitsadze A.V., Samarskii A.A., "On some simplest generalizations of linear elliptic problems", *Dokl. Akad. Nauk SSSR.*, Vol.185, pp. 139–159, 1969.

[2] Sobolevskii P.E., "On elliptic equations in a Banach space". *Differ. Uravn.* , Vol.4, No.7, pp. 1346-1348, 1969.

[3] Sapagovas M.P., "Difference method of increased order of accuracy for the Poisson equation with nonlocal conditions", *Dif. Equat.*, Vol.44, No.7, pp. 1018-1028, 2008.

[4] Ashyralyev A., Ozturk E., "On Bitsadze-Samarskii type nonlocal boundary value problems for elliptic differential and difference equations: Well-Posedness", *Apply Mathematics and Compupation*, Vol.219, No.3, pp. 1093-1107, 2012.

[5] Ashyralyev A., "A note on the Bitsadze-Samarskii type nonlocal boundary value problem in a Banach space", *Journal of Mathematical Analysis and Applications*, Vol.344, No.1, pp. 557-573, 2008.

## Rank 0 invariant solutions of dynamics of two-phase medium

Alexandr PANOV [1]

[1] *Department of Mathematics, Chelyabinsk State University, Chelyabinsk, Russia*
*E-mail: gjd@bk.ru*

**Abstract:** There is considered a system of partial differential equations, which describes dynamics of two-phase medium [1]

$$\frac{d\rho_1}{dt_1} + \rho_1 \mathrm{div}\ \vec{u}_1 = 0,$$
$$\frac{d\rho_2}{dt_2} + \rho_2 \mathrm{div}\ \vec{u}_2 = 0,$$
$$\rho_1 \frac{d\vec{u}_1}{dt_1} + m_1 \nabla P(\rho_1, \rho_2) = -\frac{\rho_2}{\tau}(\vec{u}_1 - \vec{u}_2),$$
$$\rho_2 \frac{d\vec{u}_2}{dt_2} + m_2 \nabla P(\rho_1, \rho_2) = \frac{\rho_2}{\tau}(\vec{u}_1 - \vec{u}_2),$$

Algebra Lie of symmetry group of this system was found [2]. Optimal systems of subalgebras all dimensionals for this algebra were found in papers [3,4]. Each subalgebra from these optimal systems provides dissimilar invariant, partially invariant, or differential invariant submodel [5]. Solutions of this submodels are called invariant, partially invariant, or differential invariant solutions of original system.

Four-dimensional subalgebras from optimal systems of subalgebras give partially invariant, or invariant solutions. All rank 0 invariant solutions are found. They are received using some 4-dimensional subalgebras. All other 4-dimensional subalgebras will give only partially invariant submodels of this system.

**Keywords:** two-phase medium, rank 0 invariant solutions, symmetry group, partially invariant submodels

**2010 Mathematics Subject Classification:** 35B06, 35Q35

**References:**

[1] RakhmatuliKh., "The basics of dynamics of interpenetrating motion compressible medium", *Prikl. Mat. Mech.*, Vol.20, Isue 2, pp.184-195, 1956.

[2] Panov A., "Invariant submodels of equations system of mixture dynamics of gas and small particles in the case of three spatial variables", *Belgorod state university scientific bulletin. Mathematics and Physics*, Vol.35, No.12 (183), pp. 188-199, 2014.

[3] Ovsyannikov L.V., "On the optimal systems of subalgebras", *Lie Groups and Their Appl.*, Vol.1, No.2, pp. 18-26, 1994.

[4] Khabirov S.V., "Symmetry analisys of uncompressible liquid model with viscosity and thermal conductivity depending from temperature", Gilem, Ufa, 2004.

[5] Ovsyannikov L.V., "The SUBMODELS program. Gas dynamics", *Prikl. Mat. Mech.*, Vol.58, No.4, pp. 30-55, 1994.

## Numerical solution of the elliptic-Schrödinger equation with the multipoint nonlocal boundary condition

Yildirim OZDEMIR [1], Esra PEKONUR [2]

[1] *Department of Mathematics, Duzce University, Duzce, Turkey*
*E-mail: yildirimozdemir@duzce.edu.tr*
[2] *Department of Computer Engineering, Yalova University, Yalova, Turkey*
*E-mail: esrapekonur@yalova.edu.tr*

**Abstract:** The boundary value problem for an elliptic-Schrödinger equation with the multipoint nonlocal boundary condition is considered. The stability estimates for the solution of the given problem are established. For numerically solving this multipoint nonlocal boundary problem the first and second order of difference schemes are presented. The theoretical statements for the solution of these difference schemes are supported by the result of numerical examples.

**Keywords:** finite difference equation, partial differential equation, stability

**2010 Mathematics Subject Classification:** 65N06, 65N12

**References:**

[1] Ashyralyev A., Fattorini H.O., "On uniform difference scheme for second order singular perturbation problems in Banach spaces", *SIAM J. Math. Anal.*, Vol.23, No.1, pp. 29-54, 1992.

[2] Ashyralyev A., Ozdemir Y., "Stability of difference schemes for hyperbolic-parabolic equations", *Comput. Math. Appl.*, Vol.50, pp. 1443-1476, 2005.

[3] Ashyralyev A., Ozdemir Y., "On nonlocal boundary value problems for hyperbolic-parabolic equations", *Taiwan. J. Math.*, Vol.11, No.4 pp. 1075-1089, 2007.

[4] Ozdemir Y., Kucukunal M., "On nonlocal boundary value problems for hyperbolic Schrödinger equations", *Abstr. Appl. Anal.*, Vol.2012, pp. 1-12, 2012.

## On parabolic inverse problems with overdetermination data on spatial manifolds

Sergey PYATKOV

*Yugra State University, Khanty-Mansiisk, Russia*
*E-mail: pyatkov@math.nsc.ru*

**Abstract:** We examine the question on finding a solution to a parabolic system together with a right-hand side and coefficients in parabolic systems of equations. Let $G$ be a domain in $\mathbb{R}^n$ with a boundary $\Gamma$ of class $C^{2m}$ and $Q=(0,T)\times G$. The parabolic system is of the form

$$u_t + A(t,x,D)u = g = \sum_{i=1}^{r} b_i(t,x)q_i(t,x') + f, \quad (t,x)\in Q, \tag{1}$$

where $x'=(x_1,x_2,\dots,x_k)$, $x''=(x_{k+1},x_{k+2},\dots,x_n)$ and $A$ is a matrix elliptic operator of order $2m$ with matrix coefficients of dimension $h\times h$, representable as

$$A(t,x,D)=\sum_{i=r+1}^{sh} q_i(t,x')A_i(t,x,D_x)+A_{hs+1}(t,x,D_x),$$
$$A_i=\sum_{|\alpha|\le 2m} a_{i\alpha}(t,x)D^{\alpha},$$

$$u|_{t=0}=u_0, \quad B_j u|_S=\sum_{|\beta|\le m_j} b_{j\beta}(t,x)D^{\beta}u|_S=g_j(t,x), \tag{2}$$

where $m_j<2m$, $j=1,2,\dots,m$, and $S=(0,T)\times\Gamma$. The unknowns are a solution $u$ and functions $q_i(t,x')$ $(i=1,2,\dots,sh)$ occurring into the right-hand side of (1) and in the operator $A$. The overdetermination conditions are of the form

$$u|_{S_i}=\psi_i(t,x'),\ S_i=(0,T)\times\Gamma_i, \quad i=1,\dots,s \tag{3}$$

where $\{\Gamma_i\}$ $(i=1,2,\dots,s)$ is a set of smooth $n-1$-dimensional surfaces lying in $G$. Under certain conditions on the data, we establish the existence and uniqueness theorems for solutions to these inverse problems.

**Keywords:** inverse problem, parabolic system, existence, uniqueness

**2010 Mathematics Subject Classification:** 35R30, 35K41, 35K35

## Statistical property of earthquakes network: active and passive points

S.REZAEI [1], A.H.DAROONEH [1]

[1] *Department of Physics, Zanjan University, Zanjan, Iran*
*E-mail: s.rezaee@znu.ac.ir*
*E-mail: darooneh@znu.ac.ir*

**Abstract:** Recent results in complex systems have indicated that network theory, which is known as graph theory in mathematics, is a powerful method to handle unsolved complex problems. Earthquakes manifest spatio-temporal complex behavior that can be studied using complex networks. It is so essential for these studies to construct an appropriate network. Then we use the statistical mechanics to study the property of such a network.

We propose a new combination method of Abe - Suzuki method [1] and Telesca - Lovallo method [2] for constructing earthquakes network. We divide a geographical region into small square cells that these cells cover the entire region without any overlapping. If an earthquake with any value of magnitude occurs in a cell, we identify it as a vertex of a network. Then for connecting links between vertexes we use the visibility condition. Indeed in our method two events are connected to each other if visibility condition holds true between them.

For Iran and California earthquakes we divide the longitudinal and latitudinal ranges into cells (cell sizes changes from $4km$-$220km$). We show that the constructed networks are scale free and theirs degree distribution obey the $q$-exponential function which is used in non-extensive statistical mechanics [3]. The diagram of $q$ parameter in terms of the cell size has a peak at $31km$ for Iran and $44km$ for California. Due to dependence of network characteristics on each other for cell sizes less than peak size, only one of the cell sizes is enough to describe the earthquakes network. We found that such model network results in both the Gutenberg-Richter and Omori laws.

Also we find the universal behavior of links and nodes number with time for same areas and same resolution of Iran and California earthquakes. This behavior is power law and similar to Omori law but there are differences between them.

Furthermore, analogy to Darooneh- Lotfi [4] by using the concept of PageRank, we find the passive and active points in the geographical region of Iran.

**Keywords:** Statistical Mechanics, Earthquake Network, PageRank

**References:**

[1] Abe S., Suzuki N., "Scale-free network of earthquakes", *Europhys. Lett.*, Vol.65, pp. 581, 2004.

[2] Telesca L., Lovallo M., "Analysis of seismic sequences by using the method of visibility graph", *Europhys. Lett*, Vol.97, pp. 50002, 2012.

[3] Tsallis C., "Introduction to Nonextensive Statistical Mechanics", *Springer*, ISBN. 978-0-387-85358-1, 2009.

[4] Darooneh A.H., Lotfi N., "Active and passive faults detection by using the PageRank algorithm", *EPL*, Vol.107, pp. 49001, 2014.

## Monge-Ampere equation and pluripotential theory

Azimbay SADULLAEV [1], Bakhrom ABDULLAEV [2]

[1] *National University of Uzbekistan, Tashkent, Uzbekistan*
*E-mail: sadullaev@mail.ru*
[2] *Urgench State University, Urgench, Uzbekistan*
*E-mail: abakhrom1968@mail.ru*

**Abstract:** Pluripotential theory based on the Monge-Ampere operator $(dd^c u)^n$, where $d = \partial + \overline{\partial}$, $d^c = \frac{\partial - \overline{\partial}}{4i}$ are well-known notations, is one of the actual areas of modern mathematics. It is a direct continuation and further development of the classical potential theory, based on the Laplace operator.

In 1959 Bremermann proved that the maximal plurisubharmonic ($psh$) function $\omega(z) = \sup\{u \in psh(D)) : u|_{\partial D} \leq \varphi\}$, where $D \subset \mathbb{C}^n$ is a strictly pseudoconvex domain and $\varphi \in C(\partial D)$, has many properties of harmonic functions, and is connected with the complex Monge-Ampere equation $(dd^c u)^n = 0$. Later Kerzman (1977) proved the opposite, that if $(dd^c u)^n = 0$ then $u$ is maximal. Since maximal $psh$ functions are the basis of the pluripotential theory, then during the 1977-82 years was carried out intensive research aimed at building the pluripotential theory in the complex space $\mathbb{C}^n$: have been introduced and studied the basic objects of the theory, such as extremal Green function $V(z, K)$, $\mathcal{P}-$ measure $\omega(z, K, D)$, pluripolar sets, capacity values $\mathcal{P}(K, D), C(K, D)$, etc.

The most importantly, these studies have been used successfully in solving various problems, that have accumulated in multidimensional complex analysis and in the theory of plurisubharmonic functions. By 1982 the foundation pluripotentil theory was practically built. All the basic fundamental theory of the objects have been identified, the method of their application has been developed.

In this paper we consider a generalized Monge-Ampère equation (equation in Hessian)

$$(dd^c u)^m \wedge \beta^{n-m} = 0, \quad 1 \leq m \leq n, .. \tag{2}$$

where $\beta = dd^c|z|^2$ is the volume form in $\mathbb{C}^n$. Equation (1) for $m = 1$ gives the Poisson equation, and for $m = n-$ the Monge-Ampère equation. Equation (1) and properties of their solutions have been intensively studied in recent years. So, the real equations in the Hessian dedicated by authors such as A. Caffarelli, L.Nirenberg, J.Spruck, N. Trudinger, N. Ivochkina, X. Wang, etc. Complex Hessian equation (1) was first considered by S. Li, Z. Błocki, S. Dinev, S. Kolodziej (2005-2012). The remarkable fact in their research is to determine the class of $m-$subharmonic $(m - sh)$ functions in which it is possible to solve the equation (1). Remember, a function $u \in C^2(D)$ is called $m - sh$, if

$$(dd^c u)^k \wedge \beta^{n-k} \geq 0 \ for \ all \ k = 1, 2, ..., m.$$

The class of $m-$subharmonic functions plays the same role in the solution of equation (1), as the class of plurisubharmonic functions for the Monge-Ampère equation $(dd^c u)^n = 0$. Therefore, it is important to further in-depth study of

the class $m-sh$ functions, in particular its potential-capacity properties. In this way, we formulate a few following fundamental results (see [4-5])

**Theorem 1.** *Locally $m-$polar set is globally $m-$polar in $\mathbb{C}^n$, i.e. if a set $E \subset \mathbb{C}^n$ s locally $m-$polar, then there exists a function $u \in m-sh(\mathbb{C}^n)$ such that $u \not\equiv \infty$, $u|_E = -\infty$.*

Recall that the set $E \subset D \subset \mathbb{C}^n$ is called $m-$ polar, if there exists a function $u \in m-sh(D)$ such that $u \not\equiv \infty$, $u|_E = -\infty$.

**Theorem 2.** *If the domain $D \subset \mathbb{C}^n$ is $m-$ regular, then in it the Dirichlet problem*

$$(dd^c u)^m \wedge \beta^{n-m} = 0, \;\; u|_{\partial D} \equiv \varphi(\xi) \tag{3}$$

*has a unique solution $u \in m-sh(D)) \bigcap C(\overline{D})$ for any given boundary function $\varphi(\xi) \in C(\partial D)$.*

**Theorem 3** (Quasicontinuity, analogue of Luzin's theorem)**.** *Arbitrary $m-$ subharmonic function is continuous almost everywhere with respect to the capacity, i.e. if $u \in m-sh(D)$, then for any $\varepsilon > 0$ there exists open set $U \subset D$ such that its capacity $C(U) < \varepsilon$ and the function $u$ is continuous in $D \setminus U$.*

**Keywords:** Monge-Ampere equation, plurisubharmonic functions, Dirichlet problem

**2010 Mathematics Subject Classification:** 32U05, 32U15

**References:**

[1] Sadullaev A., "Pluripotential theory. Applications", Palmarium Academic Publishing, 2012.

[2] Błocki Z., "Weak solutions to the complex Hessian equation", *Ann.Inst.Fourier, Grenoble, 55*, Vol.5, pp.1735–1756, 2005.

[3] Dinev S., Kolodziej S., "A priori estimates for the complex Hessian equation", *Anal. PDE*, Vol.7, pp. 227–244, 2014.

[4] Sadullaev A., Abdullaev B., "Potential Theory in the Class of m-Subharmonic Functions", *Proceedings of the Steklov Institute of Mathematics*, Vol.279, pp. 155–180, 2012.

[5] Sadullaev A., Abdullaev B., "Capacities and Hessians in the Class of m-Subharmonic Functions", *Doklady Mathematics*, Vol.87, No.1, pp. 88–90, 2013.

## Maxwell-Auzhan boundary conditions for Boltzmann's moment system equations in third approximation

Auzhan SAKABEKOV [1], Yerkanat AUZHANI

[1] *Kazakh-British Technical University, Àlmaty, Kazakhstan*
*E-mail: a.sakabekov@kbtu.kz*

**Abstract:** In this article we prove existence and uniqueness of the solution of the problem with initial and Maxwell-Auzhan boundary conditions [2] for nonstationary nonlinear one-dimensional Boltzmann's six-moment system equations [1] in space of functions continuous in time and summable in square by a spatial variable. In order to obtain a priori estimation of the initial and boundary value problem for the nonstationary nonlinear one-dimensional Boltzmann's six-moment system equations we get the integral equality then use the spherical representation of vector. Then we obtain the initial value problem for Riccati equation. We have managed to obtain a particular solution of this equation in an explicit form [3-4].

**Keywords:** Maxwell-Auzhan Boundary conditions, Boltzmann's moment system equations

**2010 Mathematics Subject Classification:** 35Q20, 35B45, 35L04

**References:**

[1] Sakabekov A., Initial-boundary value problems for the Boltzmann's moment system equations, Gylym, Almaty, 276 p., 2002.

[2] Sakabekov A., Auzhani Y., Boundary conditions for the onedimensional nonlinear nonstationary Boltzmann's moment system equations, *J.Math. Phys.*, 55, 123507, 2014.

[3] Kamke E., Differentialgleichungen: Losungsmethoden und Losungen, I. Gewohuliche Differentialgleichungen, B.G.Teubner, Leipzig, 1977.

[4] Tungatarov A., Akhmed-Zaki D.K., Caushy problem for one class of ordinary differential equations, *Int. Journal of Math. Analysis*, Vol.6, No.4, pp. 31-35, 2012.

# Numerical solution of the relaxational filtration model in relaxationaly-compressed porous environment realized by the linear Darcy law by Monte Carlo and probability difference methods

Kanat SHAKENOV [1], Marzhan SULTANOVA [2]

[1] *Al–Farabi Kazakh National University, Kazakhstan*
*E-mail: shakenov2000@mail.ru*
[2] *Al–Farabi Kazakh National University, Kazakhstan*
*E-mail: marzhan.ss@mail.ru*

**Abstract:** The model of linear relaxational filtration is considered. Initial and boundary conditions are set for them (Dirichlet, Neumann and mixed). Obtained problem solved by Monte Carlo methods – "random walk on spheres" , "random walk on balls" and "random walk on lattices" of Monte Carlo methods and by probability difference methods. Filtration model in relaxationaly-compressed porous environment realized by the linear Darcy law.

$$\chi\partial_x^2\Big(p(t,x)+\lambda_m\partial_t p(t,x)\Big)=\partial_t\Big(p(t,x)+\lambda'_m\partial_t p(t,x)\Big), \tag{4}$$

$$\mathbf{W}(t,x)=-\frac{\kappa}{\mu}\mathrm{grad}_x p(t,x). \tag{5}$$

Here $p(t,x)$ is pressure, $\mathbf{W}(t,x)$ is velocity of filtration, $\chi$, $\lambda_m$, $\lambda'_m=\lambda_m\frac{\beta_*}{\beta}$, $\beta_*$, $\beta$ is parameters, and all parameters are nonnegative given numbers. The model in bounded region of filtration $\Omega\in\mathbb{R}^3$ with boundary $\partial\Omega$, and for $t\in[0,T]$, and for pressure $p(t,x)$ we consider equation (4). Then we set initial conditions for them. We consider three boundary conditions: Dirichlet, Neumann and mixed for equation (4). Obtained problems solved by Monte Carlo methods – "random walk on spheres" , "random walk on balls" and "random walk on lattices" of Monte Carlo methods and by probability difference methods. [1].

**Keywords:** Numerical modeling, model, relaxation, filtration, random walk, Monte Carlo, probability, difference, method, approximation

**2010 Mathematics Subject Classification:** 65C05

**References:**

[1] Kanat Shakenov, "Numerical Modeling of the one Model of Filtration Process by Monte Carlo Methods", Series *"Applied and Numerical Harmonic Analysis". Book "Methods of Fourier Analysis and Approximation Theory"*. Springer. Berkhäuser, pp. 237–258, 2016.

## New Economy Form (NEF) as a Plan-Market Economy

S. SHARIPOV [1], Kub.S. SHARIPOV [2], Kad.S. SHARIPOV [3]

[1,2] *IGU after Tynystanov, Kyrgyzstan*
*E-mail: 7847526@mail.ru*
[3] *KUPS, Kazakhstan*
*E-mail: 7847526@mail.ru*

**Abstract:** We propose to research a theory of growing income with a Plan-Market Economy in the New Economy Form. Humanity tends to get an increasing income to make a prosperous life. The method that leads us to the rising income has been a dream of economists. However, they fail to find it. The mentioned problem belongs to the class of economic growth and development. We started the research with the model of economic growth and development of income equal to the sum of costs.

**Keywords:** New Economy Form (NEF), improved Harrod-Domar mode, old style, abstract model

**2010 Mathematics Subject Classification:** 330.564:64.066.2

Consider the abstract model of Harrod-Domar model as follows

$$y'(t) = \lambda u(t), t \in [t_0, +\infty) \tag{1}$$

$$y(t) = u(t) + c(t), t \in [t_0, +\infty) \tag{2}$$

Initial income $y(t_0) = y_0$ (3)

**New Economy Form (NEF) as the new style of economic work.** We split the time interval $[t_0, +\infty)$ into $n$ parts as follows. Take $n+1$ points $t_0, a_1, a_2, ..., a_n \in [t_0, +\infty)$. Then we get a sequence of time intervals

$$[t_0, a_1] \cup [a_1, a_2] \cup ... \cup [a_{n-1}, a_n] \subset [t_0, +\infty)$$

Then we look at the improved Harrod-Domar model from (1) - (3) on this sequence of time intervals.

Obviously, this is the only way to fulfill the plan in a plan-market economy:

$$y' = \lambda y(t) + \lambda c(t), t \in [t_0, a_1] \cup ... \cup [a_{n-1}, a_n] \subset [t_0, +\infty)$$

with the following conditions $y(t_0) = y_0, y(a_1) = y_1, ..., y(a_n) = y_n$.

With the help of the given model we could find the economic unknown quantities.

**References:**

[1] Sharipov S., Sharipov K.S., Sharipov K.S., *Control solutions of differential and integral equations*, Vestnik IGU after K. Tynystanov, No.12, Karakol, 2004.

## Initial–Boundary–Problems for Vlasov–Poisson Equations with Homogeneous Magnetic Field

Alexander L. SKUBACHEVSKII [1]

[1] *Department of Applied Mathematics, RUDN University, Moscow, Russia*
*E-mail: skub@lector.ru*

**Abstract:** We consider the Vlasov-Poisson equations in infinite cylinder with the Dirichlet boundary condition for electric potential [1]. The Vlasov-Poisson system describes evolution of electric potential and distribution functions for densities of charged particles in high-temperature rarefied plasma. The cylindrical shape of domain corresponds to thermonuclear reactor that is called "mirror trap". We study classical solutions with supports of distribution functions strictly inside domain. Such solutions are modelling plasma confinement. We obtain explicite relations providing existence and uniqueness of such solutions in Hölder spaces.

This work was supported by Ministry for Education and Science of Russia, project No.02.a03.21.0008.

**Keywords:** Vlasov–Poisson equations, classical solutions, plasma confinement

**References:**

[1] Skubachevskii A.L., "Vlasov–Poisson equations for a two–components plasma in a homogeneous magnetic field", *Russain Math. Surveys*, Vol.69, No. 2, pp. 291–330, 2014.

---

## Numerical solution of the multiphase one dimensional model of gas lift process

Nurlan TEMIRBEKOV [1], Amankeldy TURAROV [1]

[1] *Department of Higher Mathematics, East Kazakhstan State Technical University, Ust-Kamenogorsk, Kazakhstan*
*E-mail: temirbekov@rambler.ru, t010183@gmail.com*

**Abstract:** Study of the processes of field development is of great practical importance in the oil and gas industry due to the increasing need to improve production methods. A very large number of works is dedicated to the mathematical modeling of these processes [1–3]. We consider and conduct a numerical simulation of a one-dimensional model of gas-lift process, where the movement in gas-lift well is described by partial differential equations of hyperbolic type. Difference schemes for the gas-lift model of the process on a non-uniform grid condensing near the boundaries of subdomains with gas, liquid and a gas-liquid mixture are developed.

In this paper, the mathematical model of gas lift wells is developed. A one-dimensional model of gas-lift wells is examined in which it is assumed that the flow is two-phase and isothermic in the annular portion and in the well. The system, which describes the process under study, consists of the equations of motion and continuity equations of thermodynamic state, concentration, and hydraulic resistance. At the interface, matching conditions for pressure, velocity and concentration are set that allows to obtain a formula for determining the density of the liquid phase in an explicit form. Finite-difference scheme on adaptive non-uniform grid condensing on the boundaries of the gas, liquid and liquid-gas phase, is developed. When building a grid, cubic spline function is used. The results of the proposed algorithm is illustrated by the example of a real well.

**Keywords:** gas lift process, multiphase flow, finite difference method, numerical solution

**2010 Mathematics Subject Classification:** 47.85.Dh

**References:**

[1] Aliev F.A., and Ilyasov M.H., and Dzhamalbekov M.A., "Simulation of gas lift wells", *Izvesitiya AN SSSR*, Vol.4, pp. 107-116, 2008.

[2] Shurov V.I., "Technology and engineering oil", *Moscow, Nedra*, 510 p., 1983.

[3] Barashkin R.L., "Development of models and algorithms for operation of gas-lift well as the object of an operational management system", *I.M. Gubkin Russian State University of Oil and Gas*, pp. 152, 2011.

# Numerical implementation of the method of fictitious domains for elliptic equations

Almas TEMIRBEKOV [1]

[1] *Al-Farabi, Kazakh National University, Almaty, Kazakhstan*
*E-mail: almas_tem@mail.ru*

**Abstract:** In the paper we study the elliptical type equation with strongly changing coefficients. We are interested in studying such equation because the given type equations are yielded when we use the fictitious domain method. In this paper we suggest a special method for numerical solution of the elliptic equation with strongly changing coefficients. We have proved the theorem for the assessment of developed iteration process convergence rate. We have developed computational algorithm and numerical calculations have been done to illustrate the suggested method effectiveness.

**Keywords:** elliptical equation, Dirichlet problem, the equation with rapidly changing coefficients, computational algorithm, iterative process, fictitious domain method, boundary conditions

**2010 Mathematics Subject Classification:** 97N40

**References:**

[1] Samarskiy A.A., "About One Economic Algorithm of Numerical Computation of Differential and Algebraic Equation System (in Russian), *Journal of Computation Mathematics and Mathematical Physics*, V.4, No. 3, pp.580–585, 1964.

[2] Kucherov A.B., Nikolayev E.S., "Alternately-Triangular Method of Mesh Elliptical Equiations Solution in Rectangle, *Journal of Computation Mathematics and Mathematical Physics,*, V.16, No. 5, pp. 1164–1174, 1976.

[3] Lebedev V.N., "Method of Composition, *OBM. AS USSR, Moscow*,1986.

[4] Volkov E.A., "About Methods Solving Difference Equations for Sectionally-Uniform Media and With the Right Part of Given along Curve, *DAN USSR*, 283, No. 2, pp.274–277, 1985.

[5] Bugrov A.N., Konovalov A.N., Scherbak V.A.,"Fictitious Domain Method in Plane Elastic Static Problems, *Numerical Methods of Mechanics Of Continua. Novosibirsk*, V.5, No.1, pp. 20–30, 1974.

[6] Konovalov A.N., "Fictitious Domain Method in Torsion Problems, *Numerical Methods of Mechanics Of Continua. Novosibirsk*, V.4, No.2, pp. 109–115, 1973.

[7] Smagulov Sh.S., "Fictitious Domain Method for Boundary Problem of Navie-Stocks Equation, *Novosibirsk: Ed. V.Ts. SO AN USSR, Predprint*, No.68, pp. 68–73, 1979.

[8] Orunkhanov M.K., Smagulov Sh.S., "Fictitious Domain Method for Navie-Stocks Equation in Terms of Flow Function and Vorticity with Inhomogeneous Boundary Conditions *Calculation Technologies. Novosibirsk: SO RAN*, V.5, No.3, pp. 46–53, 2000.

[9] Kuttykozhaeva Sh.N., "Fictitious Domain Method for Navie-Stocks Equation *Vestnik KazGU. Section Math., Mech., Inf.*, No.13, pp. 54–59, 1998.

## The gyroscope movement with variable moments of inertia

A.N. TYUREKHOJAYEV [1], G.U. MAMATOVA [1]

[1] *K.I. Satpayev Kazakh National Technical University, Almaty, Kazakhstan*
*E-mail: tyurekhodja@mail.ru, mamatovag@mail.ru*

**Abstract:** It took more than two hundred years since the publication of the equations of rigid body dynamics with a fixed point, but research is still ongoing. Great interest in this problem is caused by the fact that in the motion of rigid body with a fixed point are observed gyroscopic effects, widely used in modern technology. The topicality of considering the problem about motion of rigid body with a fixed point is also due to the need to allow for perturbing gravitational, electrical, magnetic and other forces, variability of inertia moment of the object and a wide application in practice.

This kind of problems can be reduced to the study of systems of nonlinear and with variable coefficients differential equations, generating analytical solutions of which offers great mathematical difficulty, and it is possible in a relatively small number of cases. Therefore, the construction of analytical solutions for a wide class of such problems is topical.

This paper discusses motion equations of an axisymmetric rigid body with a fixed point with variable moment of inertia, which is described by a system of nonlinear dynamic Euler equations

$$\begin{cases} \dfrac{d[A(t)]p}{dt} + [C(t) - B(t)]qr = M_x, \\ \dfrac{d[B(t)]q}{dt} + [A(t) - C(t)]rp = M_y, \\ \dfrac{d[C(t)]r}{dt} + [B(t) - A(t)]pq = M_z, \end{cases} \tag{1}$$

where $A(t)$, $B(t)$, $C(t)$ are body's inertia moments relative to $x$, $y$, $z$ axes connected with the body; $p$, $q$, $r$ are projections of the body's angular velocity vector on these axes; $M_x$, $M_y$, $M_z$ are moments of external resistance forces.

The system of differential equations (1) is considered jointly with initial conditions

$$\begin{gathered} t = 0 : p(0) = p_0, \;\; q(0) = q_0, \;\; r(0) = r_0, \\ \dot{p}(0) = 0, \;\; \dot{q}(0) = 0, \;\; \dot{r}(0) = 0. \end{gathered} \tag{2}$$

The solution of this problem is obtained by the technique of partial discretization of non-linear differential equations, built by A.N. Tyurekhodzhayev on the base of the generalized functions theory.

**References:**

[1] Courant, R. and Hilbert, D., *Methods of Mathematical Physics*, vol. 1, Wiley-VCH Verlag GmbH, Weinheim (2004).

## On the stability of the telegraph equation with time delay

Allaberen ASHYRALYEV [1], Deniz AĞIRSEVEN [2], Koray TÜRK [2]

[1] *Emeritus Professor, Turkey & Turkmenistan*
*E-mail: aallaberen@gmail.com*
[2] *Department of Mathematics Trakya University, Edirne, Turkey*
*E-mail: denizagirseven@trakya.edu.tr, korayturk@trakya.edu.tr*

**Abstract:** Telegraph equation is mostly interested in physical systems. Many physicists and engineers use telegraph equation without time delay [1]-[**?**]. However, for several reasons, in problems encountered in real life, time delay should be considered in modeling. In this study, the initial value problem for telegraph equations with delay

$$\begin{cases} \frac{d^2v}{dt^2} + \alpha\frac{dv}{dt} + Av(t) = aAv([t]), t > 0, \\ \\ v(0) = \varphi, v'(0) = \psi \end{cases} \tag{1}$$

in a Hilbert space $H$ with a self-adjoint positive definite operator $A$, is considered. Here $A \geq \delta I$, $\delta \geq \frac{\alpha^2}{4}$ and $0 \leq a < 1$. Theorem on stability estimates for the solution of this problem is established. As a test problem one-dimensional delay telegraph equation with Dirichlet boundary conditions is considered. Numerical solutions of this problem are obtained by first and second order of accuracy difference schemes.

**Keywords:** initial and boundary value problems, difference schemes, delay telegraph equations.

**2010 Mathematics Subject Classification:** 35L20, 65N06

**References:**

[1] Lamb H., Hydrodynamics, Cambridge Mathematical Library, Cambridge University Press, Cambridge, UK, 6th edition, 1993.

[2] Lighthill J., Waves in Fluids, Cambridge University Press, Cambridge, UK, 1978.

[3] Taflove A., Computational Electrodynamics: The Finite-Difference Time-Domain Method, Artech House, Boston, Mass, USA, 1995.

[4] Sieniutycz S., and Berry R.S., "Variational theory for thermodynamics of thermal waves" Physical Review E, Vol.65, No.4, Article ID 046132, 11 p., 2002.

# Determination of a control parameter of the $r$-modified Crank-Nicholson difference scheme for the Schrödinger equation

Allaberen ASHYRALYEV [1], Mesut URUN [2]

[1] *Emeritus Professor, Turkey & Turkmenistan*
*E-mail: aallaberen@gmail.com*
[2] *Faculty of Mechanical Engineering, İstanbul Technical University, İstanbul, Turkey*
*E-mail: aashyr@fatih.edu.tr mesuturun@gmail.com*

**Abstract:** The differential equations with parameters have been studied extensively by many researchers (see, [1]- [9] and the references therein). However, such problems were not well-investigated in general. In the present paper, the second order of accuracy $r$-modified Crank-Nicholson difference schemes are presented for the numerical solution of the boundary value problem for the Schrödinger differential equation with parameter $p$

$$\begin{cases} i\frac{du(t)}{dt} + Au(t) + iu(t) = f(t) + p, \ 0 < t < T, \\ u(0) = \varphi, \ u(T) = \psi \end{cases}$$

in a Hilbert space $H$ with self-adjoint positive definite operator $A$. The well-posedness of this difference schemes are established. The stability inequalities for the solution of difference schemes for three determination of a control parameter problems for the Schrödinger equation are obtained.

**Keywords:** determination of a control parameter problem, difference scheme, Schrödinger equation, stability, well-posedness

**References:**

[1] Eidelman Y.S., *Boundary Value Problems for Differential Equations with Parameters*, PhD Thesis, Voronezh State University, Voronezh, 1984 (in Russian).

[2] Di Blasio G., and Lorenzi A., "Identification problems for parabolic delay differential equations with measurement on the boundary", Journal of Inverse and Ill-Posed Problems, Vol.15, No.7, pp. 709-734, 2007.

[3] Orlovsky D., and Piskarev S., "On approximation of inverse problems for abstract elliptic problems", Journal of Inverse and Ill-Posed Problems, Vol.17, No.8, pp. 765-782, 2009.

[4] Ashyralyev A., "On a problem of determining the parameter of a parabolic equation" , Ukranian Mathematical Journal, Vol.62, No.9, pp. 1200-1210, 2010.

[5] Serov V., and Päivärinta L., "Inverse scattering problem for two-dimensional Schrödinger operator", Journal of Inverse and Ill-Posed Problems, Vol.14, No.3, pp. 295-305, 2006.

[6] Ashyralyev A., Sobolevskii P.E., "New Difference Schemes for Partial Differential Equations", Operator Theory Advances and Applications, Birkhauser Verlag, Basel, Boston, Berlin, 2004.

[7] Sobolevskii P.E., "Difference Methods for the Approximate Solution of Differential Equations", Izdat. Voronezh. Gosud. Univ., Voronezh, 1975 (in Russian).

[8] Ashyralyev A., Urun M., "Determination of a control parameter for the difference Schrö dinger equation", Abstract and Applied Analysis, Article ID 548201, 8, 2013.
[9] Ashyralyev A., Urun M., "A second order of accuracy difference scheme for Schrödinger equations with an unknown parameter", Filomat, Article Vol.28:5, pp. 981-993, 2014.

---

## On the analytical solution methods for sine-Gordon equation

Ozgur YILDIRIM [1], Meltem UZUN [2]

[1] *Department of Mathematics, Yildiz Technical University, Istanbul, Turkey*
*E-mail: ozgury@yildiz.edu.tr*
[2] *Department of Mathematics, Yildiz Technical University, Istanbul, Turkey*
*E-mail: meltemu@yildiz.edu.tr*

**Abstract:** In this work some solution methods are considered to obtain the analytical solution of nonlinear sine-Gordon equation. Theoretical aspects and some basic concepts on solution of this equation are given. Several applications of sine-Gordon equation in biology and biophysics are introduced.

**Keywords:** soliton solutions, nonlinear waves, traveling wave solutions

**2010 Mathematics Subject Classification:** 35C08, 74J30, 35C07

**References:**

[1] Kudryashov N.A., "Simplest equation method to look for exact solutions of nonlinear differential equations", *Chaos, Solitons and Fractals*, Vol.24, No.5, pp. 1217-1231, 2005.
[2] Zhao Y.M., "Exact solutions of coupled sine-Gordon equations using the simplest equation method", *Journal of Applied Mathematics*, Vol.2014, pp. 5 p., 2014.
[3] Ablowitz M.A., and Clarkson P.A., "Solitons, Nonlinear Evolution Equations and Inverse Scattering", *Cambridge University Press*, 1991.
[4] Yomosa S., "Solitary excitations in deoxyribonucleic acid (DNA) double helices", *Physical Review A*, Vol.30, pp. 474-480, 1984.

## On the numerical solution of sine-Gordon equation

Ozgur YILDIRIM [1], Meltem UZUN [2]

[1] *Department of Mathematics, Yildiz Technical University, Istanbul, Turkey*
*E-mail: ozgury@yildiz.edu.tr*

[2] *Department of Mathematics, Yildiz Technical University, Istanbul, Turkey*
*E-mail: meltemu@yildiz.edu.tr*

**Abstract:** In this work the approximate solution of well-known nonlinear equation is studied. Finite difference method and the fixed point iteration are used to obtain numerical solution of the nonlinear equation. Error analysis is presented using MATLAB implementation.

**Keywords:** nonlinear boundary value problems, difference equations, numerical analysis

**2010 Mathematics Subject Classification:** 34B15, 39A10, 97N40

**References:**

[1] Yildirim O., and Uzun M., "On the numerical solutions of high order stable difference schemes for the hyperbolic multipoint nonlocal boundary value problems", *Applied Mathematics and Computation*, Vol.254, pp. 210-218, 2015.

[2] Yildirim O., and Uzun M., "On third order stable difference scheme for hyperbolic multipoint nonlocal boundary value problem", *Discrete Dynamics in Nature and Society*, Vol.2015, 16 p., 2015.

[3] Ablowitz M.J., Herbst B.M., and Schober C.M., "On the numerical solution of the sine-Gordon equation. 2. Performance of numerical schemes", *Journal of Computational Physics*, Vol.131, pp. 354-367, 1997.

# Dynamic analogues of Green and Gauss's formulas for unsteady dynamics of anisotropic elastic medium at antiplane deformation

Gulmira K. ZAKIR'YANOVA [1]

[1] *Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: gulmzak@mail.ru*

**Abstract:** The paper is devoted to the solution of non-stationary boundary value problems (BVP) for an anisotropic elastic medium at antiplane deformation. Wave propagation in continuous medium is described, as a rule, by hyperbolic equations and systems [1, 2]. In this case we have hyperbolic equation of second order with constant coefficients

$$(a_{ij}\partial_i\partial_j - \rho\partial_t^2)u(x,t) = G(x,t), \;\; i,j = \overline{1,N}$$

where $a_{ij} \geq \lambda\|\xi\|$, $\lambda = const > 0$, $\rho = const$, $G$ is regular function. For the solution of the non-stationary BVP for this equation the Method of Generalized Functions (MGF) [3] is used. This method allows the original BVP to lead to the solution of the equations in the space of generalized functions with a certain right-hand side of the class of singular generalized functions such as simple and double layers and use the properties of fundamental solutions of equations for the construction of their generalized solutions.

Here MGF is used for investigation of unsteady dynamics of anisotropic elastic medium at antiplane deformation. The statement of two initial boundary problems are given. Dynamic analogues of Green and Gauss's formulas constructed [4] and their integral representations obtained.

**Keywords:** hyperbolic equation, boundary value problem, generalized solutions, elastic medium

**2010 Mathematics Subject Classification:** 35L15, 74J05

**References:**

[1] L.A. Alexeyeva, Sh.A. Dildabayev, G.K. Zakiryanova, A.B. Zhanbyrbaev "Boundary Integral Equations Method in two-and three-dimensional problems of elastodynamics *Computational mechanic*. Vol.18:2, pp.147-157, 1996.

[2] Alexeyeva L.A., Zakiryanova G.K., "Generalized Solutions of InitialBoundary Value Problems for Second Order Hyperbolic Systems *Computational Mathematics and Mathematical Physics*. V0l.51:7. 11941207, 2011.

[3] Alexeyeva L.A., "The generalized functions method in non-stationary boundary value problems for wave equation *Mathematical Journal*. Vol.6:1. pp. 16–32, 2006.

[4] Zakiryanova G.K., "Dynamic analogies of Green's and Gauss formula for hyperbolic equations", *Mathematical Journal*, Vol.10, No.3, pp. 67-74, 2010.

## Optimization of turbulent flow in a radial reactor with fixed bed

Uzak ZHAPBASBAYEV, Gaukhar RAMAZANOVA, Madina ZHAKENOVA

*Kazakh-British Technical University, Almaty, Kazakhstan*
*E-mail: uzak.zh@mail.ru*

**Abstract:** The data of the computation of turbulent flow in the CF-$\pi$ and CP-$\pi$ configurations of the radial reactor with a fixed bed are presented.

Turbulent flow in the annular duct and in the central pipe is described by the Reynolds equations [1]. The macroscopic $k$-$\epsilon$ turbulence model [2, 3] in the variables averaged over the local volume of the catalyst bed and is used for determining eddy viscosity in a porous media.

The macroscopic variables determined over a local volume of the porous medium are used in a fixed bed for flow description [4]. The motion equations with the Ergan drag law in macroscopic variables are used for flow description in the catalyst bed.

The computational data are obtained for the averaged and turbulent characteristics, and it is shown that the flow in the fixed bed causes the generation of the turbulence kinetic energy and its dissipation rate; the flow in the CF-$\pi$ configuration is distributed more uniformly as compared to the CP-$\pi$ configuration of the radial reactor. Computed data are compared with the experimental ones.

**Keywords:** radial reactor, fixed bed, $k$-$\epsilon$ turbulence model.

**2010 Mathematics Subject Classification:** 02.60.Cb, 44.15.+a

**References:**

[1] Launder B.E., and Sharma B.I., "Application of the energy dissipation model of turbulence to the calculation of flow near a spinning disk", *Lett. Heat and Mass Transfer*, Vol.3, pp. 269-289, 1974.

[2] Nakayama A., and Kuwahara F., "Macroscopic Turbulence Model for Flow in a Porous Medium", *J. Fluids Engng*, Vol.121, pp. 427-433, 1999.

[3] Nakayama A., and Kuwahara F., "A general macroscopic turbulence model for flows in packed beds, channels, pipes and rod bundles", *J. Fluids Engng*, Vol.130, pp. 1-7, 2008.

[4] Slattaery J.C., "Advanced Transport Phenomena", Cambridge University Press, Cambridge, 1999.

## Instability of control system in the neighborhood of program manifold

Sailaubay ZHUMATOV

*Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: sailau.math@mail.ru*

**Abstract:** We will consider the problem of construction of the steady systems of automatic control on the program manifold. Given program $\omega(t, x) = 0$ is performed only subject to condition, if initial values of the state vector of the system satisfies to equality $\omega(t_0, x_0) = 0$. But these equalities can be not always performed. Therefore at the construction of the systems of differential equations it is necessary to have some requirements on quality properties of the program manifold as stability, convergence, dissipatedness relatively to some function. To the construction of the systems of differential equations on the program manifold, possessing properties of stability, to the optimality and establishment to estimations for quality of transient's indexes in the neighborhood of program manifold, were devoted great number of works. The review of these works is conducted in [1 - 3].

In this work the control system with respect to vector-function $\omega$ is investigated with nonlinearity $\varphi(\sigma)$, satisfying to the local quadratic conditions. For this system was built Lyapunov function of the form

$$V(\omega,\ \xi) = \omega^T L\omega + \int_0^\sigma \varphi^T \beta d\sigma,$$

where $L = L^T,\ \beta = diag(\beta_1, \ldots, \beta_r)$ are matrices not necessarily sign definiteness. The sufficient conditions of instability are obtained with respect to vector-function $\omega$ by means of this function.

**Keywords:** program manifold, automatic control system, absolute stability, Lyapunov's function

**2010 Mathematics Subject Classification:** 34K20, 93C15, 34K29

**References:**

[1] Galiullin A.S., Mukhametzyanov I.A., Mukharlyamov R.G., "Review of researches on the analytical construction of the systems programmatic motions", *Vestnik RUDN M.*, No.1, pp. 5–21, 1994.

[2] Zhumatov S.S., Krementulo B.D., Maigarin B.G., "Lyapunov second method in the problems of stability and control by motion", *Almaty*, 228 p., 1999.

[3] Mukharlyamov R.G., "Differential-algebraic equations of programmed motions of Lagrangian dynamical systems", *Mechanics of Solids*, Vol.46, No.4, pp. 534–543, 2011.

## Role of non-commutative geometry in gravitational collapse scenarios

Amir Hadi ZIAIE [1]

[1] *Department of Physics, Shahid Bahonar University, Kerman, Iran*
*E-mail: ah.ziaie@gmail.com*

**Abstract:** The collapse process of a homogeneous scalar field has been studied by many authors and it has been shown that such a process would end in a curvature singularity which can be either hidden behind a horizon or be visible to external observers. In the present work, we revisit the collapse process of a spherically symmetric homogeneous scalar field (in FRW background) minimally coupled to gravity, when the phase-space deformations are taken into account. Such a deformation is mathematically introduced as a particular type of noncommutativity between the canonical momenta of the scale factor and of the scalar field. In the absence of such deformation, the collapse culminates in a spacetime singularity. However, when the phase-space is deformed, we find that the singularity is removed by a non-singular bounce, beyond which the collapsing cloud re-expands to infinity. More precisely, for negative values of the deformation parameter, we identify the emergence of a negative pressure term, which slows down the collapse to finally avoid the singularity formation. Depending on the model parameters, one can find a minimum value for the boundary of the collapsing cloud or correspondingly a threshold value for the mass content below which no horizon would form. Such a setting predicts a threshold mass for black hole formation in stellar collapse and manifests the role of non-commutative geometry in physics and especially in stellar collapse and supernova explosion. [1–3].

**Keywords:** gravitational collapse, non-commutative geometry, non-linear differential equations, numerical method

**References:**

[1] Pankaj S. Joshi, Gravitational collapse and spacetime singularities," Cambridge University Press, 2007.

[2] Rasouli S.M.M., Ziaie A.H., Marto J., Moniz P.V., Phys. Rev. D 89, 044028 (2014).

[3] Douglas M.R., and Nekrasov N.A., Rev. Mod. Phys. 73 977 (2001);
R. J. Szabo, Phys. Rep. 378, 207, 2003.

# MATHEMATICS EDUCATION

## Geometry study using the interactive whiteboard

Bayramsoltan ANNADURDYYEVA [1], Nursoltan NURMYRADOVA [2]
[1] *30 th school, Gokdepe, Ahal, Turkmenistan*
*ogulmengli70@gmail.com*
[2] *84 th school, Ashgabat, Turkmenistan*
*dowletnurmyradow@gmail.com*

**Abstract:** Information technology can be used in different learning modes, and most importantly - in the mode of the graphic illustration of the studied material.

The effect of information technology on the secondary mathematics classroom has been studied by several authors ( [1–4] and references therein).

The computer as a tracer has a number of advantages compared with compass and straightedge. So much for the image of bodies of rotation required to build the image of the circle is an ellipse. However, ruler and compass can construct an approximate image of the ellipse, it has not always been good quality. Using SMART Board interactive whiteboard and SMART Notebook software and other computer softwares, it can be created a wide variety of models of geometric figures, which is difficult in the case of material models, both technically and in material terms.

In geometry lessons students work a lot with graphic representation of spatial geometrical figures, which are not always clearly reflect their properties. Therefore, of particular interest are image editors allow teacher and students to create and modify computer models of geometric objects.

The potential of information technology in conducting computer simulation in order to independently obtain new knowledge about the geometric objects based on the study of computer models makes these technologies in learning an instrument of knowledge.

In this paper, methodocial aspects of using information technologies in the teaching geometry. The prospects of development of the learning environment in geometry study by using interactive whiteboard are revealed.

**Keywords:** information technologies, secondary school, teacher professional development, interactive whiteboard

**2010 Mathematics Subject Classification:** 97U30, 97U70

**References:**

[1] Lavy I., Lebron U., "The emergence of mathematical collaboration in an interactive computer environment", *International Journal of Computers for Mathematical Learning*, Vol.9(1), pp. 1-23, 2004.

[2] Krymova L.N., "Interactive board in math class", *Mathematics in school,* Volume 2008, No.1, pp. 31-39, 2008.

[3] Sohoreva,T.A., "Using the interactive whiteboard in math class", *In Proceedings of the V All-Russian Scientific-Practical Conference: Actual problems mat. Education in schools and ped. high school*, April 8-10, 2009, edited by Payson, B.D., Barnaul, pp. 245-249, 2009.

[4] Ashyralyyeva M.A., Arslanova B.M., The methodical aspects of the mathematics study using the interactive whiteboard, The abstract book of the conference AMS 2015, Antalya, Turkey, p. 54, 2015.

## Usage of recognizing algorithm handwriting in mathematical education

Maral Allaberenovna ASHYRALYYEVA [1,1], Abdyresul Nurgeldiyevich CHARYYEV [1,2]

[1] *Department of Applied of Mathematics and Informatics, Magtymguly Turkmen State University, Ashgabat, Turkmenistan*
*E-mails:*[1] *ashyrmaral2010@mail.ru,* [2] *abdyresul@gmail.com*

**Abstract:** Improving quality of modern mathematical education can be achieved by using modern information technologies ( [1–4]).

It is well known that recognizing algorithms handwriting can be divided to various types by objective of application.

In this work, we use Neural networks method. We propose software which based on recognizing algorithm. Proposed programm permit us to organize lessons, save education materials and view of created early lessons.

The applied programm is an addition tool for teachers and allows them to show students in class all handwritten, painted and tagged items.

The program permit us to save the classified lessons, drafts and help fast viewing.

**Keywords:** mathematics education, Neural networks method, recognating algorithm

**2010 Mathematics Subject Classification:** 97D40, 97U30

**References:**

[1] Popel M., The methodical aspests of the algebra and the mathematical analysis study using the Sagemath Cloud, *Journal of Information Technologies in Education*, Vol. 19, pp. 93-100, 2014.

[2] Thomas M.O.J., and Hong Y.Y., Teacher integration of technology into mathematics learning, *International Journal for Technology in Mathematics Education*, Vol.20(2), pp. 69-84, 2013.

[3] Ashyralyyeva M.A., Arslanova B.M., The methodical aspects of the mathematics study using the interactive whiteboard, The abstract book of the conference AMS 2015, Antalya, Turkey, p. 54, 2015.

[4] Gurbanmyradov G., Meredaliyeva O., Increasing mathematics maturity, The abstract book of the conference AMS 2015, Antalya, Turkey, p. 113, 2015.

## Fundamentalization of mathematical knowledge while teaching students inverse and incorrect problem

Yessen BIDAIBEKOV [1], Victor KORNILOV [2], Guldina KAMALOVA [3], Nagima AKIMZHAN [4]

[1,3,4] *Department of Computer Science and Informatization of Education, KazNPU named after Abai, Almaty, Kazakhstan*
*E-mail:* [1] *esen_bidaibekov@mail.ru*
[2] *Department of Informatization of Education, Moscow City Training University, Moscow, Russia*
*E-mail:*[2] *vskornilov@mail.ru*

**Abstract:** Fundamentalization of mathematical knowledge of students of physical and mathematical areas of training in higher educational institutions is considered in the article [1, 2].

Training inverse and incorrect problems, the solution of which requires the development of a number of mathematical theories has a significant opportunity to improve the fundamental mathematical knowledge. To confirm this, the article presents several productions of inverse problems and methods of their study, included in the content of training inverse and incorrect problems of the future specialist of physical and mathematical profile [3–5].

**Keywords:** training of inverse and incorrect problems, mathematical education, competence in the field of applied mathematics, student.

**2010 Mathematics Subject Classification:** 65L09, 65M32

**References:**

[1] Bidaibekov, Y.Y., Kornilov, V.S., Kamalova, G.B."Teaching of future mathematics and computer science teachers of inverse problems for differential equations", *Vestnik of Moscow city training university. Series of Informatics and informatization of education"*, No.3(29), pp. 57-69 , 2014.

[2] Bidaibekov, Y.Y., Kornilov, V.S., Kamalova, G.B., Akimzhan, N.Sh. "Fundamentalization of knowledge system on applied mathematics in teaching students of inverse problems for differential equations", *AIP Conference Proceeding*, 1676, 020044-1020044-5, AIP Publishing, Antalya, doi: 10.1063/1.4930470 2015, p.6.

[3] Bidaibekov, Y.Y., Kornilov, V.S., Kamalova, G.B., Akimzhan N.Sh., "Using computer technologies in teaching university students of inverse problems for ordinary differential equations", *Vestnik of Peoples' friendship university of Russia. Series of "Informatization of education"*, No.2, pp. 57-72, 2015.

[4] Kabanikhin, S.I., "Inverse and incorrect problems ", Siberian scientific edition, Novosibirsk p. 458, 2009.

[5] Romanov, V.G.,"Inverse problems of mathematical physics: monograph", Nauka, Moscow p. 264, 1984.

## Geometric heritage of al-Farabi in education

Yessen BIDAIBEKOV [1], Guldina KAMALOVA [1],
Bektas BOSTANOV [1], Kairat UMBEBAEV [1]

[1] *Abai Kazakh National Pedagogical University, Kazakhstan, Almaty, Kazakhstan*
*E-mail: bbgu@mail.ru*

**Abstract:** On the base of studying geometrical treatise of al-Farabi and A.Kubesov's works devoted to their research the expediency of introduction geometric heritage of al-Farabi to modern mathematics education due to its uniqueness, which is reflected in use of algorithmic approach in solving geometrical problems and its application-oriented direction is proved in the article.

**Keywords:** mathematical heritage, geometric construction, construction with help of compass and ruler, algorithm, algorithmic approach, application-oriented direction, mathematics education.

**2010 Mathematics Subject Classification:** 373+51.7+004

**References:**

[1] Bidaibekov Y.Y., Bostanov B.G., & Kamalova G.B. (2013). The mathematical heritage of Al-Farabi by A.Kubesov in modern conditions of education. In IX International ISAAC Congress, Poland, Krakow, August 5-9, 2013. pp.33-34

[2] Carry J. Tee (University of Aucland), Kubesov A.K. The Mathematical Heritage of al-Farabi (in Russian). Journal for the history of Arabic science. - No.1 (1978), pp. 150-153.

[3] Rescher N. (1962). Al-Farabi: An Annotated Bibliography. Pittsburgh: University of Pittsburgh Press.

[4] Kubesov A. (1974). Mathematical heritage of al-Farabi. Alma-Ata: Nauka (in Russian).

[5] Bidaibekov Y.Y., Bostanov B.G., Kamalova G.B., & Dzhanaberdieva S.A. (2015). Mathematical heritage of al-Farabi in modern education. Vestnik of Kazakh national university named after al-Farabi. Series of politology, philosophy, culturology, 2(51), 50-57 (in Kazakh).

[6] Al-Farabi (1972). Mathematical treatises. Alma-Ata: Nauka (in Russian).

[7] Courant R., & Robbins H. (2001). What is Mathematics? (An elementary approach to ideas and methods). Moscow: MCFCME (in Russian).

[8] Government program "Madeni mura" - Cultural heritage of Kazakhstan:- http://www.madenimura.kz/ru/government-program-madenimura/programs-madenimura/.

## Features and advantages of training teachers to use means of informatization to profile teaching mathematics

Yessen BIDAIBEKOV [1], Vadim GRINSHKUN [2], Guldina KAMALOVA [1]

[1] *Abai Kazakh National Pedagogical University, Almaty, Kazakhstan*
*E-mail: esen_bidaibekov@mail.ru, g_kamalova@mail.ru*
[2] *Moscow City Teacher Training University, Russia*
*E-mail: vadim@grinshkun.ru*

**Abstract:** The usage advantages of modern means of informatization of mathematics education are discussed in the article. Aspects of training teachers to effective use of such means within the framework of profile teaching mathematics pupils are considered.

**Keywords:** profile education, mathematics, informatization of education, training teachers

**2010 Mathematics Subject Classification:** 373+51.7+004

**References:**

[1] Organization rules of educational process on distance learning technologies / Appr. by Resolution of the Government of the RK 112 from January 19, 2012. - (https://www.do.ektu.kz/laws/MONRK/p2.pdf).

[2] Grigoryev S.G., Grinshkun V.V. Informatization of education. Fundamental basis: textbook for students of pedagogical HEIs and listeners of the improvement system of teachers' qualification. - Tomsk: "TML-Press", 2008. - 286 p.

[3] Bidaibekov Y.Y., Kamalova G.B., Kisselyova Y.A. Theoretical and methodological basis of training teachers to use information and communication technologies in the context of e-learning. - Almaty: KazNPU named after Abai, 2012. - 143 p.

## Value of new technologies in improving the branch of learning

Abdyresul Nurgeldiyevich CHARYYEV [1,1], Maral Allaberenovna ASHYRALYYEVA [1,2],

[1] *Department of Applied of Mathematics and Informatics, Magtymguly Turkmen State University, Ashgabat, Turkmenistan*

*E-mails:* [1] *abdyresul@gmail.com,* [2] *ashyrmaral2010@mail.ru*

**Abstract:** The creation of electronic mathematical textbooks with multimedia devices and realization of school lessons using a computer is studied. A practical program that allows us to automate the use of new technologies in practical departments is being created. Practical program enables upgrading of schoolwork. The PHP web-programming language and MySQL information management system of practical program has been prepared.

The practical program offers new additional tools for school technique using modern technologies.

The practical program is an additional tool for teachers, it makes possible to show all the things shown, drawn and marked by teacher directly on students' computers. Besides, the program gives students the opportunity to carry out home works through this program. And it allows the teacher to identify academic student performance on a particular lesson or a specific topic. Because the teacher is able to not only see the result of the performed work, but also to monitor the performance of tasks from the beginning of the process to the end. Practical program gives some information about the user by using the handwriting recognition algorithms. Interpretation of lessons and works into the electronic form is made in the form of vector graphics. The difference between vector graphics and video inveterate method is that it takes 20-30 times less space and preserves the quality of the recording regardless of the demonstration screen. Therefore, it can increase the number of participants in electronic form during the lesson and spreading of these lessons in the program with ease.

**Keywords:** mathematics education, PHP, MySQL

**2010 Mathematics Subject Classification:** 97U30

**References:**

[1] Thomas M.O.J., and Hong Y.Y., Teacher integration of technology into mathematics learning, *International Journal for Technology in Mathematics Education*, Vol.20(2), pp. 69-84, 2013.

## Formation of project and research skills of students in calculation of limits

Burkhan KALIMBETOV [1], Indira OMAROVA [1]

[1] *Department of Mathematics, H.A. Yasawi International Kazakh-Turkish University, Turkestan, Kazakhstan*
*E-mail: bkalimbetov@mail.ru, omarovai10@mail.ru*

**Abstract:** Today different goals and objectives are qualitatively assigned to the educational system, which are defined by the dictated social order, by a new model of expert and professional. Consequently, it is necessary to form a personality, which is able to organize his professional career in the ever-changing socio-cultural conditions. Therefore, current priorities of higher education are aimed at development of specialist personality, the leading component of which is the possession of an independent creative and research activities [1].

According to the heads of project and research works, the most of students are not able to independently propose and substantiate a hypothesis, plan an activity, formulate a goal, search and analyze necessary information, represent results of research, realize reflection, competently write a report. This is due to the fact that not only the students but also their teachers are not trained to project and research activities [2].

The method of rationalization [3], applied to rationalize various irrational and trigonometric expressions, and hence to solve irrational and trigonometric equations and inequalities, to calculate limits, etc., is a generalization of the well-known idea about the use of rationalize integration substitutions [4], where these substitutions are widely used to rationalize the various classes of differential expressions by I. Newton, L. Euler, P. Chebyshew and other mathematicians.

**Keywords:** project-research activity, radical expression, rationalization, indeterminacy, rational

**References:**

[1] Pavlova E.B., "Bulletin of BSU, Ulan-Ude, Vol.15, pp. 59-63, 2011.
[2] Alnikova T.Y., "Autoabstract. ... cand. ped. sciences, Tomsk, 2007.
[3] Fikhtengol'ts G.M., "Course of differential and integral calculus, M.: Nauka, 1969.
[4] Demidovich B.P., "QCollection problems and exercises in mathematical analysis, M.: Nauka, 1969.

## Methods of professionally-oriented training of complex function theory

Bahytgul KASKATAEVA [1], Yssen BIDAIBEKOV [2], Nurbol ABISHEV [2]

[1] *Kazakh State Women's Pedagogical University, Almaty, Kazakhstan*
*E-mail: kaskataeva@yandex.ru*
[2] *Kazakh National Pedagogical University, Almaty, Kazakhstan*
*E-mail: esen_bidaibekov@mail.ru, n.abishev@mail.ru*

**Abstract:** This article describes the interactive methods of professionally-oriented education of theory analytic functions for the purpose of formation of professional competence of students; The paper defined the concept of "professional competence" and "activity approach to professional-focused training to the mathematician" and the classification of interactive forms and methods of teaching.

**Keywords:**

**2010 Mathematics Subject Classification:** 97I80, 97I20, 97I30

**References:**

[1] B.R. Kaskatayeva, Formation of methodical competence of the future mathematics teachers. Monograph. Almaty: KazgoszhenPU. (2009) 37-40.
[2] T.G. Brazhe, Development of the general culture of the teacher in system of a continuous pedagogical education. The relationship of general and professional culture of the teacher. Materials and conferences. - St. Petersburg: the Institute of Adult Education of RAO. (1992) 86-87.
[3] Anan'ev, Selected psychological works.V.2. (1980).
[4] A.K.Markova, Preparation of the future teachers to the active professional activity in the course of course studying pedagogy: Abstract ... cand. ped. Sciences. (1994) 12–16.
[5] L.S. Vygotski, Collected works: V. 6 - N. 2. – (1982) 47–50.
[6] A.N. Leont'ev, Selected psychological works:. In 2 vols II. M: Pedagogy (1983) 127-135.
[7] S.L. Rubenshtein, Fundamentals of general psychology. - SPb .: Peter. (1999) 127-133
[8] V.V. Davydov, The theory of developmental education. - (1996) 145-150.
[9] B.D. El'konin, The concept of competence from the standpoint of developing training // Modern approaches to competence-oriented education. – Krasnoyarsk. (2002) 25–27.
[10] L.S. Vygotskii, Questions of theory and history of psychology // Collected works: In 6 vols 1. (1982)
[11] N.N.Tul'kibayeva, A.V. Usova, Methods of teaching students how to solve problems. Chelyabinsk. (1981).

## Student-centered model for teaching numerical methods course

Margarita KLUNNIKOVA [1], Nikolai PAK [2],
Tatyana PUSHKARYEVA [1], Tatian STUPINA [1]

[1] *Siberian Federal University, Krasnoyarsk, Russia*
*E-mail: mklunnikova@gmail.com, a_tatianka@mail.ru, stupina_tv@mail.ru*
[2] *Krasnoyarsk State Pedagogical University, Krasnoyarsk, Russia*
*E-mail: nik@kspu.ru*

**Abstract:** The quality of modern education is strongly influenced by students' personal motivation. Long scientific and educational experience confirms that the success of students depends on various factors mainly related to their personal ambitions and preferences. The purpose of this work is to prove applicability of student-centered model for teaching "Numerical methods" course that weakens students' personal and psychological barriers to conscious material learning and contributes to their computational thinking development.

**Keywords:** student-centered model, computational thinking, "Numerical methods" course

**2010 Mathematics Subject Classification:** 97Q30, 97N99

**References:**

[1] Chenglie Hu, Computational thinking: what it might mean and what we might do about it, Proceedings of the 16th annual joint conference on Innovation and technology in computer science education, Darmstadt, Germany, June 27-29, 2011.

[2] *Exploring Computational Thinking (ECT)* Available at: https://www.google.ru/edu/resources /programs/exploring-computational-thinking/. (Accessed 19/04/16).

[3] Henner E.K., Computational thinking // Education and science, 2. pp. 18-33, doi: 10.17853/1994-5639-2016-1-18-33. 2016 (In Russian).

[4] Podlinjaev O.L., Molokova O.A., Person-centered approach as a basis for students adaptation in learning at the university / Theory and practice of social development, Vol.11, 2013 (In Russian).

[5] Rogers K., The freedom to learn / K. Rogers, D. Freiberg. Moscow: Meaning, 527 p. 2002 (In Russian).

[6] Andreeva N.M., Innovative experience of training students of economic specialties in "Computer science" course // Bulletin of Astafev Krasnoyarsk state teachers university. Vol.2(20), pp. 18-21, 2012(In Russian).

[7] Pak N.I., Projective approach in training as an information process: monograph. Krasnoyarsk, 112 p., 2008(In Russian).

[8] Andreeva N.M., Pak N.I., The role of roadmaps for students e-learning computer science in classical universities // Open education, Vol.3, 2015(In Russian).

[9] Kuzyk J., What is a roadmap. Available at: http://www.strf.ru/material.aspx?CatalogId=223&d_no=20108#.VSsllPCdeEQ (Accessed 25/04/16).

[10] *The concept for developing roadmaps / Agency for strategic initiatives / National business initiative for improving investment climate in the Russian Federation.* Moscow: July, 2012 (In Russian).

[11] Lee S., Park Y., Customization of technology roadmaps according to roadmapping purposes: overall process and detailed modules, Technology Forecasting & Social Change, Vol.72, pp. 267-583, 2005.

## Didactic bases of system formation of learners' methodological knowledge

Dosymkhan RAKHYMBEK [1], Marzhan ABDUALIYEVA [1], Nurlibai MADIYAROV [1], Shadyar ALTYNBEKOV [1]

[1] *M. Auezov South Kazakhstan State University, Kazakhstan*
*E-mail: rakhymbek_d@mail.ru, Madnur@rambler.ru, abdualieva82@mail.ru, altynbekov_shadiar@mail.ru*

**Abstract:** The article deals with the role and significance of methodological knowledge in education. Studied the basic sources, factors of educational content and the expected results of the inclusion of methodological knowledge in school education. Didactic conditions of formation and learning of methodological knowledge in teaching the basics of sciences of a school course. Analyzes the principles and criteria for acquiring methodological knowledge of learners.There are some demands representation of methodological knowledge in educational subject. The selection procedure of the complex methodological knowledge and its didactic and methodological foundation. [1]. On the basis of analysis of the main sources, shows that methodological knowledge are reflected in all elements of the content of education, at all levels in the form of the whole complex of knowledge. Methodological knowledge included in the content of education in an implicit form of knowledge: about the knowledge of scientific methods (universal, general scientific and private scientific or specific); logic of scientific knowledge (formal and dialectical); regularities of the process of scientific cognition; the logic of scientific activity. Methodological knowledge present in the content of education and in dissected forms. In studying academic subject is recommended to show the nature of scientific cognition and its effectiveness, specific techniques of every field of knowledge, and use those and other methods that have the general scientific and educational character. [2].

**Keywords:** methodological knowledge, school, teacher training, academic subject

**2010 Mathematics Subject Classification:** 00A35, 97C70

**References:**

[1] Taubayeva Sh.T., Pedagogics methodology. Almaty: "Karasay", 2013.
[2] Rakhymbek D., Developing metodological knowledge of pupils. Almaty: RBC, 1998.

## Analysis of theoretical and methodological bases of teaching object-oriented programming languages in higher school

Beybitgul SHAYAKHMETOVA [1], Nurgul ORUMBAYEVA [1,2], Sholpan OMAROVA [3]
[1] *Buketov Karaganda Stated University, Karaganda, Kazakhstan E-mail:* kazahzavod@mail.ru
[2] *Institute of Applied Mathematics, Karaganda, Kazakhstan, E-mail:*orumbayevan@mail.ru
[3] *Karaganda Economic University, Karaganda, Kazakhstan E-mail:* sheo_1953@mail.ru

**Abstract.** In this paper, we affect one of the most urgent calls of our time - problem of educating at higher school.

One of the most pressing challenges of our time is the problem of teaching in higher school. Training of young people in the sphere of the latest computer technologies is the teachers' task. When teaching teachers should change the methodology of training specialists in higher school in accordance with the adequately changing development pace. Software changes due to the appearance on the market of new information technologies. The subject area of computer science changes in an extremely dynamic way. Teachers of higher school that teach subjects associated with the technology of software creation are constantly faced with the problems of changing the content of curricula, working programs, methodical support (lectures, labs, tests, etc.), the development of new teaching and learning aids.

In connection with the application in the educational process of the various versions of educational software, teaching guides on the use of this software have to be accordingly changed. A certain bias towards the creation of software for complex systems has been recently observed; the tasks under consideration become much more complicated, so the old techniques and methods previously used to create simple or formalized programs are not enough. At the modern stage of social development, the needs of educational, industrial, commercial entities have significantly increased, and simple programs no longer meet them. Education of students of information specialties has not been provided yet with the necessary theoretical basis for the creation of complex software systems, the necessary subjects are often presented in the curricula fragmentary, in the list of elective courses.

**Keywords:** analysis, development, complex systems, information, object-oriented approach, model, education, methods, pedagogics, algorithmization

**2010 Mathematics Subject Classification: 68N15**

**References:**

[1] The State Compulsory Standard of Education of the Republic of Kazakhstan. Higher professional education. Baccalaureate. SES RK 23.08.1080-2012. Astana: Ministry of Education and Science of the Republic of Kazakhstan, 2012.

[2] Shayakhmetova B.K., Technology of creating programs for complex systems. Astana: Turan-Pro, 170p., 2010.

[3] National Report of the Russian Federation at the II International UNESCO Congress "Education and Computer Science", "Education policy and new information technologies". M., 34p., 1996.

[4] Zhuzhzhalov V.Ye., Integration methods of learning programming in the higher school course of computer science. Bulletin of Moscow State Pedagogical University. A series of computer science and informatization of education. M., 2003. No.1.
[5] Virt N. Algorithms + program structure = program. – M.: Mir, 406p., 1985.

## History of mathematics and its teaching in higher school

Simon SEROVAJSKY

*Department of Differential Equations and Control Theory, al-Farabi Kazakh National University, Almaty, Kazakhstan*
*E-mail: serovajskys@mail.ru*

**Abstract:** Currently teaching hours for basic mathematics are reduced because of the education reform. This is typical for the specialty "*Mathematics*", and for the natural sciences and technical fields where mathematics is not critical discipline. This adversely affects the quality of trained specialists. We cannot change this situation generally. However, we would like somehow influence it.

Modern educational programs usually provide elective courses. Then we could offer a special non-standard mathematical course for the students. It will be able to interest them, is aimed at improving the general mathematical culture of pupils and, thus, to fill the numerous gaps in their mathematical education. It will be able to interest them, will aimed at improving the general mathematical culture of pupils and, thus, to fill the numerous gaps in their mathematical education. This is "*History of Mathematics*". We are teaching this course for many years at the Mechanics and Mathematics Faculty of al-Farabi Kazakh National University as an elective course. We recommend [1] as an appropriate textbook for it.

**Keywords:** history of mathematics, mathematical education

**2010 Mathematics Subject Classification:** 01A05

**References:**

[1] Serovajsky S., "Thinking about Mathematics and its History, *Almaty, Instant printing*, 856 p., 2015.

## Methodological questions of filling the virtual university program modules

Aigerim ZHANBOLOVA [1], Bogdat ZHANBUSINOVA [2], Gaukhar ISKAKOVA [3], Shaukenova KELBET [4], Beybitgul SHAYAKHMETOVA [5]
[1,2,3,4,5]Karaganda State University, Karaganda, Kazakhstan
E-mail: [1]zhanbolova.aigerim@mail.ru, [5]kazahzavod@mail.ru

**Abstract:** Methodological questions of filling the virtual university program modules. Consider the questions filling Virtual University program modules with teaching tools. Here are the main teaching tools used in the content and also the types of computer training programs:

Work module tutorial (unit). Units are designed by each module, the average during one academic period. Each student receives 25 -27 unit and methodical recommendations for work with workers modular textbook.

Training didactic or methodical tools logically complement each other and enhance didactic functions of the discipline. General view of the virtual university materials must be more than 2000 items of educational products. Among them - the simulated movies, music videos, video lectures (review and modular), satellite TV lectures, author's lecture courses, slide lectures. Computer training programs . The university (virtual) should have more than 300 computer-based training programs for all academic disciplines. Work with such programs aimed at enhancing the student's mental activities, the formation and consolidation of skills.

Training programs represent one of the forms of independent work of students on the stage when a certain knowledge is already formed.

Collective training is a classroom group session aimed at updating and processing of professional skills and allows students unleash your creativity in an interactive way. Student Appraisal System. Continuous monitoring and assessment of assimilation quality of each module in accordance with standard criteria is also the original difference of this technology. Certification system is divided into a learning module, current and complete[1].

We note the characteristic features of virtual education: flexibility, modularity, parallelism, large audience, economical, workability, social equality, internationalization, the new role of the teacher, positive impact on student (listener), quality.

Creating information systems focuses on the complex means of interaction with the end user who performances a "smart interface" role on modern level and enables interactive solution of information problems on a computer. Note that for this purpose capabilities of the Web environment are widely used; this capabilities ensure the provision of a user interface for working with one or more databases. Thus, it can be seen that the known three-tier client-server architecture of the database is displayed naturally on the Web environment, where the Web - browser plays a role of a "thin" client, a Web-server - the role of the application server.

**Keywords:** supertyutor, proftyutor, training, certification, module, unit, integration, telecommunications, infrastructure, configuration, portal, environment.

**References:**

[1] Morozov V.A., "Creativity high school teacher, *ZHigher Education Today (in Russian)*, No.3, p. 64-72, 2004.

# OTHER TOPICS

# Furstenberg type theorem for a multiplicative product of Markov random matrices

Nursadyk AKANBAY [1]

[1] *Department of Mechanics and Mathematics, Al-Farabi Kazakh National University, Almaty, Kazakhstan*
*E-mail: noureke@mail.ru*

**Abstract:** In the work, considering the certain conditions of "general situation", Furstenberg type theorem is proved for multiplicative product of a special type of Markov random matrices.

The classical theorem of Furstenberg of strict positivity of the largest Lyapunov exponent for the product of a large number of independent matrices and its generalization to the case of the dependent (for example, forming a Markov chain) random matrices play a key role in the whole range of issues related to the spectral properties of disordered structures.

Currently, there are several different approaches to proof theorems of Furstenberg [1], [2], [3]. In this paper we prove the the Furstenberg type theorem for multiplicative product (in terms of [4] multiplicative integral) Markov random matrices of a special form.

Note that, the necessity of studying of such multiplicative products of random matrices arise in the problems of the evolution of the magnetic field in so-called Markov linear incompressible flows [5].

**Keywords:** random matrix, a multiplicative product, transition function, transition density, the formula of Kac-Feynman, the Lyapunov exponent

**2010 Mathematics Subject Classification:** 15B52, 37H15

**References:**

[1] Furstenberg H., "Non-commuting random products", *Trans. Amer. Math. Soc.*, Vol.108,3, pp. 377-428, 1963.

[2] Tutubalin V.N., "A central limit theorem for product of random matrix and some of its applivations", *-Symposin Hath., Bologna,XXI, "*, pp. 101-116, 1977.

[3] Wircer A.D., "On matrix and operator of casual works", *Teor.probab. and its approx. m. 24, No. 2"*, pp. 361-370, 1979.

[4] Gantmakher F.R., "Matrix theory", *M.*, p. 576, 1967.

[5] Akanbay N., "Some questions of the theory of matrix fields in random media,", *Abstract. cand. diss.(Manuscript) Moscow, Moscow State University,"*, 1987.

# On asymptotic normality of solutions of the Cauchy problem of a parabolic equation with random right side

Nursadyk AKANBAY [1], Zoiya SULEYMENOVA [1], Samal TAPEEVA [1]

[1] *Department of Mechanics and Mathematics, Al-Farabi Kazakh National University, Almaty, Kazakhstan*
*E-mail: suleymenova.zoiya@mail.ru*

**Abstract:** Consider the Cauchy problem for the heat equation

$$\frac{\partial u(t,x)}{\partial t} = \frac{1}{2}\frac{\partial^2 u(t,x)}{\partial x^2} + C(t,x), \quad u(0,x) = f(x), \qquad (1)$$

where $t \geq 0,\ x \in \mathbb{R}, f(x),\ C(t,x)$are continuous bounded (generally speaking, random) functions. Then to solve the problem (1) we have

$$u(t,x) = M_x\left[f(W_t) + \int_0^t C(t-s, W_s)ds\right],$$

where $W_s$ is Wiener process and sign $M_x$ means taking the conditional expectation on all facing at the initial time t = 0 of point $x$, trajectories of $W_t$ process. Next we consider the special case of the equation (1), namely, consider the equation

$$\frac{\partial u(t,x)}{\partial t} = \frac{1}{2}\frac{\partial^2 u(t,x)}{\partial x^2} + g(x)\dot{W}_t, u(0,x) = f(x), \qquad (2)$$

where $t \geq 0,\ x \in \mathbb{R}, f(x), g(x)$ are limited continuous functions, $\dot{W}_t$ is "white noise" [1, 2].

**Theorem 1.** Let $f(x), g(x)$ be continuous and bounded functions. Then, if the function $g(x)$ at $|x| \to \infty$ satisfies relation $|g(x) - \sigma(x)| \to 0$, where $\sigma(x) = \sigma_1\ , x > 0;;\ \sigma(x) = \sigma_2\ ,\ x < 0$; then distribution of the random function $\frac{u(t,x)}{\sqrt{t}}$, where $u(t,x)$ is a solution of Cauchy problem given by equation (2), which at $t \to \infty$ converges to distribution of the normal random variable with parameters $\left(0, \frac{(\sigma_1+\sigma_2)^2}{4}\right)$.

**Keywords:** parabolic equation, Wiener process, white noise, asymptotic normality, infinitesimal operator

**2010 Mathematics Subject Classification:** 35R60, 60H15

**References:**

[1] Venttsel A.D., "The course of the theory of random processes", *M .: Nauka*, 1975.
[2] Akanbay N., "Fundamentals of probability theory, mathematical statistics and the theory of random processes", *Almaty: ed. "Kazakh University"*, 2007.

## Modeling of thorium transport by the water flow in the river Kechi-Kemin

Valentina ALEKHINA [1], Ilya STARODUMOV [2], Oleg STARODUMOV [2]

[1] *Institute of Physical-Technological Problems and Materials Science, Bishkek, Kyrgyz Republic*
*E-mail: alekhina@jet.kg*

[2] *Laboratory of Multi-Scale Mathematical Modeling, Ural Federal University, Ekaterinburg, Russia*
*E-mail: ilya.starodumov@urfu.ru*

**Abstract:** Radioactivity of the Kechi-Kemin river is mainly due to transported thorium. The problem of the spread of radioactive pollution of the river Kechi-Kemin and their impact on the life is an important task that requires permanent monitoring and research. An important part of such work is mathematical modeling.
The current talk presents the modeling of thorium transfer in the section of the river Kechi-Kemin. To solve this problem, we used the numerical method [1] for the Saint-Venant equation to describe behaviour of water flow. This method has used together with thorium advection-diffusion transfer model along the river watercourse. By the use of river regime information and thorium source data we have simulated the possible behavior of thorium propagation in water flow. This report contains a description of the methods and the main results of the calculations.

**Keywords:** Kechi-Kemin, radioactive, river modeling

**2010 Mathematics Subject Classification:** 35Q35, 35Q86

**References:**

[1] Starodumov I., "Numerical Scheme for Solution of an Approximation of Saint-Venant Equation , *Contemporary Engineering Sciences*, Vol. 7, No. 35, 1861 - 1866, 2014.

# One-types in ordered theories

Bektur BAIZHANOV

*Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: baizhanov@math.kz*

**Abstract:** Let $A$ be a subset of a saturated enough model $N$ of $T$. The analysis of one-types is based on using the properties of $A$-definable family of 1-formulas (**Points A, B**) [1], [2], [3], of 2-formulas (**Point C**) and properties of interactions of one-types (**Point D**) [4]. Taking into consideration the points **A, B, C** it is possible to show the existence of 18 kinds of one-types in ordered theories. The properties of of 1-formulas permits to give an example complete ordered theory with two non-orthogonal 1-types one is definable, second is non-definable.

Let $p, q \in S_1(A)$. It said to be $p$ is *weakly orthogonal to* $q$ ($p \perp^w q$) if if $p(x) \cup q(y)$ is complete two-type over $A$ [5].

**Definition 1.** *Type $p$ is definable if and only if $\forall\varphi(x,\bar{y})\ \exists\psi_\varphi(\bar{y},\bar{a})$, such that*

$$A \models \psi_\varphi(\bar{b},\bar{a}) \Leftrightarrow \varphi(x,\bar{b}) \in p$$

**Theorem 2.** *There exist linear ordered structurel $M = \langle M; =, <, E^2, S^2\rangle$, infinite set $A \subset M$, two one-types over $A$, $p, q \in S_1(A)$ such that $p \not\perp^w q$, $p$ is definable and $q$ is non-definable.*

**Keywords:** ordered structures, 1-formula, 1-type.

**2010 Mathematics Subject Classification:** 03C64.

**References:**

[1] Baizhanov B.S., "Expansion of an o-minimal model by unary convex predicates", in: T. Nurmagambetov (Eds), *Researches in theory of algebraic systems*: Collection of papers, University of Karaganda, pp. 3–23, 1995.

[2] Baizhanov B.S., "Expansion of a model of a weakly o-minimal theory by family of unary convex predicates", *Journal of Symbolic Logic*, Vol. 66, No. 3, pp. 1382–1414, 2001.

[3] Baizhanov B.S. and Verbovskyi V.V., "O-stable theories", *Algebra and Logic*, Vol. 50, No. 3, pp. 211–225, 2011.

[4] Baizhanov B.S., "Orthogonality of one-types in weakly o-minimal theories", in A.G. Pinus and K.N. Ponomaryov, *Algebra and Model Theory II*, Novosibirsk State Technical University, pp.3–28, 1999.

[5] Shelah S. "Classification Theory and the number of non-isomorphic models", *North-Holland, (Studies in Logic and the Foundations of Mathematics)*, Amsterdam–New York–Oxford–Tokyo, Vol. 92, 544 p., 1978.

## On the Tietze extension theorem in soft topological spaces

Cigdem GUNDUZ ARAS [1], Sadi BAYRAMOV [2]

[1] *Department of Mathematics, Kocaeli University, Kocaeli, 41380, Turkey*
*E-mail: caras@kocaeli.edu.tr*
[2] *Department of Algebra and Geometry, Baku State University, Baku, 1148, Azerbaijan*
*E-mail: baysadi@gmail.com*

**Abstract:** The theory of soft set was introduced by Molodtsov in 1999 as a new mathematical tool for deal with uncertainties. The idea of soft topological space was first given by Shabir and Naz. The purpose of this paper is to prove Tietze Extension Theorem using Uryshon's Lemma in the so-called "soft topological space". After giving the necessary separation axioms based on the soft point definition in [3] such as soft completely regular space and soft normal space, we prove important theorem for soft separation axioms. Later we give Uryshon's Lemma. Finally we prove Tietze Extension Theorem using Uryshon's Lemma.

**Keywords:** Soft set, soft completely space, soft normal space, Urysohn Lemma and Tietze Extension Theorem.

**2010 Mathematics Subject Classification:** 54D15, 54D35

**References:**

[1] Molodtsov D., "Soft set-theory-first results", *Comput. Math. Appl.*, Vol. 37, pp. 19-31, 1999.
[2] Shabir M., Naz M., "On soft topological spaces", *Comput. Math. Appl.*, Vol.61, pp. 1786-1799, 2011.
[3] Bayramov S., Gunduz (Aras) C., "Soft locally compact and soft paracompact spaces", *J. Math. and System Sci.*, Vol.3, , pp. 122-130, 2013.

## A study on intuitionistic fuzzy supra soft topological spaces

Cigdem GUNDUZ ARAS

*D*epartment of Mathematics, Kocaeli University, Kocaeli, Turkey
E-mail: caras@kocaeli.edu.tr

**Abstract:** The aim of this paper is to introduce some necessary definitions on the so called "intuitionistic fuzzy supra soft topological spaces". Furthermore, we give definitions of fuzzy strongly supra soft connected and fuzzy supra soft continuous mapping. We present characterization theorems of their.Finally, we define intuitionistic fuzzy supra soft compactness and invesitigate some results of intuitionistic fuzzy soft compactness.

**Keywords:** intuitionistic fuzzy supra topology, fuzzy strongly supra soft connected

**2010 Mathematics Subject Classification:** 03E75, 54A40

**References:**

[1] Molodtsov D., "Soft set-theory-first results", *Comput. Math. Appl.*, Vol.37, pp. 19-31, 1999.

[2] TuranlıN., "An Overview of Intuitionistic Fuzzy Supratopological Spaces", *Hacettepe Journal of Math. Statistics*, Vol.32, pp. 17-26, 2003.

[3] Bayramov S., Gunduz Aras C., "On Intuitionistic Fuzzy Soft Topological Spaces", *TWMS J. Pure Appl. Math.*, Vol.5, No.1, pp. 66-79, 2014.

## The probability distribution function for the sum of squares of independent random variables

Yu. FATEEV [1], D. DMITRIEV [1], V. TYAPKIN [1], N. KREMEZ [1], V. SHAIDUROV [2]

[1] *Siberian Federal University, Krasnoyarsk, Russia*
*E-mail: fateev_yury@inbox.ru*
[2] *Institute of Computational Modelling, Russian Academy of Sciences, Krasnoyarsk, Russia*
*E-mail: shaidurov04@mail.ru*

**Abstract:** In the paper, the probability distribution function is derived for the sum of squares of random variables for nonzero expectations. This distribution function enables one to develop an efficient one-step algorithm for phase ambiguity resolution when determining the spatial orientation from signals of satellite radio-navigation systems. Threshold values for rejecting false solutions and statistical properties of the algorithm are obtained.

**Keywords:** probability distribution functions normal distribution, variance, squared standard deviation

**2010 Mathematics Subject Classification:** 60A05, 60C05

**References:**

[1] Yu. Fateev, D. Dmitriev, V. Tyapkin, E. Garin, and V. Shaidurov. *AIP Conference Proceedings*. **1611**, 12–14 (2014).
[2] G. Giorgi, P.J.G. Teunissen, S. Verhagen, P.J. Buist. Improving the GNSS Attitude Ambiguity Success Rate with the Multivariate Constrained LAMBDA Method, *in Geodesy for Planet Earth*. International Association of Geodesy Symposia. **136**, 941–948 (2012).
[3] P. Buist. "The baseline constrained LAMBDA method for single epoch, single frequency attitude determination applications", *in Proceedings of the 20th International Technical Meeting of the Satellite Division*, Institute of Navigation, **3**, 2962–2973 (2007).
[4] Yu. Fateev, A. Kurnosov. Ionosphere parameters definition. *Transactions of International Siberian Conference on Control and Communications, SIBCON 2013*. Proceedings. doi: 10.1109/SIBCON.2013.6693622.
[5] A. Borka, G. Even-Tzur. *Survey Review*. **46**, 335, 122–131 (2014).
[6] V.I. Tikhonov. *Statistical radiotechnics*, Moscow: Soviet Radio, 1966 (in Russian).
[7] B.R. Levin. *Theoretical Foundations of Statistical Radio Engineering*, Moscow: Soviet Radio, 1974 (in Russian).
[8] Yu. Fateev, D. Dmitriev, V. Tyapkin, I. Ishchuk, and E. Kabulova. *ARPN Journal of Engineering and Applied Sciences*. **10**, 8264–8270 (2015).

## Hyperimmunity and $A$-computable universal numberings

Assylbek ISSAKHOV

*Department of Mechanics and Mathematics, Al-Farabi Kazakh National University, Almaty, Kazakhstan*
*E-mail: asylissakhov@mail.ru*

**Abstract:** Whether there exists a computable universal numbering for a computable family is the key question in theory of numberings. In a very general setting, this problem was explored in [1]. For sets $A$ that are Turing jumps of the empty set, the problem was treated in [2] and other papers.

It is well known that, in the classical case, every finite family of c.e. sets has a universal (principal) computable numbering.

S.A. Badaev and S.S. Goncharov have showed that for every set $A$ such that $\emptyset' \leq_T A$, a finite family $S$ of $A$-c.e. sets has an $A$-computable universal numbering if and only if $S$ contains the least set under inclusion. But also they have showed that for an infinite family of $A$-c.e. sets if $\emptyset' \leq_T A$ then the presence of the least set under inclusion is neither necessary nor sufficient to have an $A$-computable universal numbering, [3].

We considered a family $F$ of total functions, computable relative to oracle $A$. Taking into the account an infinite set $A$ is hyperimmune iff no recursive function majorizes $A$, [4], we have showed that if $A$ is a hyperimmune set and if $A$-computable (finite or infinite) family $F$ of total functions contains at least two elements, then $F$ has no universal $A$-computable numbering. However, if Turing degree of a set $A$ is hyperimmune-free then every $A$-computable finite family of total functions has $A$-computable universal numbering.

**Keywords:** universal numbering, hyperimmune set, hyperimmune-free degree

**2010 Mathematics Subject Classification:** 03D45, 68Q15

**References:**

[1] Ershov Yu.L., "Theory of numberings", *Handbook of Computability Theory. - North-Holland; Amsterdam: Stud. Log. Found. Math.*, Vol.140, pp. 473-503, 1999.

[2] Badaev S.A., Goncharov S.S., Sorbi A., "Completeness and universality of arithmetical numberings", *Computability and Models, S.B. Cooper and S.S. Goncharov (eds.). - New York: Kluwer Academic/Plenum Publishers*, pp. 11-44, 2003.

[3] Badaev S.A., Goncharov S.S., "Generalized computable universal numberings", *Algebra and logic*, Vol.53, No.5, pp. 355-364, 2014.

[4] Miller W., Martin D.A., "The degree of hyperimmune sets", *Z. Math. Logik Grundlag. Math.*, Vol.14, pp. 159-166, 1968.

## A system dynamics model to forecasting of the United States gross savings

Seyit KERIMKHULLE [1]

[1] *L. N. Gumilyov Eurasian National University, Astana, Kazakhstan*
*E-mail: kerimkhulle@gmail.com*

**Abstract:** This work presents a macroeconomic model for long-term forecasts describing compounding process of the gross saving on based the feedback loops of the theory of system dynamics and their empirical evidence on the statistical data of the U.S. economy [1] with the help of econometric analysis under different assumptions on the function of adjusting time. The system dynamics model for the forecasting indicators of economy, business and finance are widely used and they was studied detail in [2–9].

Indeed, to build the model compounding gross saving used of variables: $y_t > 0$, $t \in Z$ - a initial state, $y_{t+T} > 0$, $T = 1, 2, \ldots$ - a stock; $m \gg 1$ - a $S$-shaped limit, $r_t \geq 0$ - a growth rate; $y_{t+j}^{4/5} > 0$, $j = 0, 1, \ldots, T-1, \ldots$ - a reinforcing feedback, $1 - (y_{t+j}/m)^{1/5} > 0$ - a balancing feedback, $\Delta(t+j)$ - a function of adjusting time, $\Delta y_{t+j} = y_{t+j+1} - y_{t+j}$ - a endogenous flow, $r_t y_{t+j}^{4/5}(1 - (y_{t+j}/m)^{1/5})\Delta(t+j)$ - a exogenous flow and for stock-flow relation chosen the Von Bertalanffy logistic growth equation [10], then the forecasting problems of the gross savings was reduced to following econometric model:

$$\log_a(1-(y_{t+t}/m)^{1/5}) = \log_a(1-(y_t/m)^{1/5}) - r_t \sum_{j=0}^{T-1} \Delta(t+j)/(5m^{1/5}\ln a) + \varepsilon_{t+T-1},$$

$$E(\varepsilon_{t+T-1} \mid \mathrm{X}) = 0,$$
$$Var(\varepsilon_{t+T-1} \mid \mathrm{X}) = \sigma^2 \mathrm{I},$$
$$Cov(\varepsilon_{t+T-1}, \varepsilon_{t+S-1} \mid \mathrm{X}) = 0,$$
$$\varepsilon_{t+T-1} \mid \mathrm{X} \approx N(0, \sigma^2 \mathrm{I}),$$

where $\varepsilon_{t+T-1}$ - random errors, X - a matrix of the observations, $E(\cdot)$ - a expectation, $Cov(\cdot)$ - a covariance, $Var(\cdot)$ - a variance, I - a identity matrix, $N(\cdot, \cdot)$ - a Normal distribution and $a > 0$, $a \neq 1$ - parameter for adjusting time and this parameter determined by Otelbaev average conditions [11]: $T_t = \inf_{T=1,2,\ldots} \left\{ T \mid r_t \sum_{j=0}^{T-1} \Delta(t+j)/(5m^{1/5}\ln a) \geq 1 \right\}$.

**Keywords:** model of system dynamics, von Bertalanffy growth model, econometric measures

**2010 Mathematics Subject Classification:** 91B62, 91B82, 93A30

**References:**

[1] Federal Reserve Economic Data. Federal Reserve Bank of St. Louis, "Gross Saving", *U.S. Bureau of Economic Analysis*, https://fred.stlouisfed.org/series/GSAVE, 2016.

[2] Forrester, J., Industrial Dynamics. Cambridge: Pegasus Communications, 1961.

[3] Sterman, J. D., Business Dynamics: Systems Thinking and Modeling for a Complex World. Irwin / McGraw-Hill, 2000.

[4] Yamaguchi, K., Money and Macroeconomic Dynamics. Accounting System Dynamics Approach, *Japan Futures Research Center*, Osaka, 2015.

[5] Clements, M. P., Galvao A. B., Forecasting US output growth using leading indicators: An appraisal using MIDAS models. *J. of Appl. Econometrics*, Vol. 24, No. 7, pp. 1187-1206, 2009.

[6] Elliott, G., Timmermann A., Economic Forecasting, *J. of Econ. Literature*, Vol.46, pp. 3-56, 2008.

[7] Lyneis, J. M. System Dynamics for Market Forecasting and Structural Analysis, *System Dynamics Review*, Vol. 16, pp. 3-25, 2000.

[8] Modis, T., "Long-term GDP forecasts and the prospects for growth", *Technol. Forecast. Soc. Chang*, Vol. 80, pp. 1557-1562, 2013.

[9] Yamashita, T., "Macroeconomic Model of the Regional Economy in a Society with a Declining Population: A Simulation of Shizuoka Prefecture's Economy", *J. of Econ. Policy Studies*, Vol. 4, No. 2, pp. 67-70, 2007.

[10] Tsoularis, A., "Analysis of Logistic Growth Models", *Res. Lett. Inf. Math. Sci*, Vol. 2, pp. 23-46, 2001.

[11] Otelbaev, M., "On summary ability with a weight of a solution of the Sturm-Lowville equation", *Mathematical Notes*, Vol. 16, No. 6, pp. 1172-1179, 1974.

---

## Selections, games and Karamata theory

Ljubiša D.R. KOČINAC

*University of Niš, Faculty of Sciences and Mathematics, 18000 Niš, Serbia*
*E-mail: lkocinac@gmail.com*

**Abstract:** We present various results which show how ideas from selection principles theory (developed in 1920s and 1930s by K. Menger, W. Hurewicz, F. Rothberger, W. Sierpiński) and corresponding game theory can be applied to the classical Karamata theory (1930s) for sequences of real numbers. These results are related to regularly and rapidly varying sequences and a number of their variations, as well as to rates of divergence of such sequences. Statistical versions of the mentioned notions will be also discussed.

**Keywords:** Selection principles, game theory, regular variation, rapid variation, rate of divergence

**2010 Mathematics Subject Classification:** 26A12, 40A05, 91A05

**References:**

[1] Bingham, N.H., Goldie, C.M. and Teugels, J.L., "Regular Variation", Cambridge University Press, Cambridge, 1987.

[2] Djurčić, D., Kočinac, Lj.D.R. and Žižović, M.R., "Some properties of rapidly varying sequences, *Journal of Mathematical Analysis and Application*, Vol. 327, No. 2, pp. 1297–1306, 2007.

[3] Djurčić, D., Kočinac, Lj.D.R. and Žižović, M.R., "A few remarks on divergent sequences: rates of divergence II, *Journal of Mathematical Analysis and Applications*, Vol. 367, No. 2, pp. 705–709, 2010.

[4] Haan, L. de, "On Regular Variations and its Applications to the Weak Convergence of Sample Extremes", Mathematical Centre Tracts, Vol. 32, Amsterdam, 1970.

[5] Karamata, J., "Über die Hardy-Littlewoodschen Umkehrungen des Abelschen Stätigkeitssatzes", *Mathematische Zeitschrift*, Vol. 32, pp. 319-320, 1930.

[6] Karamata, J., "Sur un mode de croissance réguliére des fonctions", Mathematica (Cluj), Vol. 4, pp. 38-53, 1930.

## Application of the theory of games for selection of effective defense

Damir KULBAEV, A.Z. ABDENOV

*Faculty of Informational Systems, Eurasian National University, Astana, Kazakhstan*
*E-mail: damirkaz@mail.ru*

**Abstract:** This article aims to prove that the use of games theory in decision making can aid in choosing the best methods of defense against attack weighed against the possible costs of such methods. Decision making in games theory assumes interaction between the seller or buyer with the environment as the environment fluctuates and the outcome is the expected payoff for such decisions. The decision maker makes decisions based on the possibility of risk and the expected payoff and whether or not the risk will be advantageous. The Wald Test is justified in situations where the possibility of environmental health is unknown, the solution occurs only once, and there is no risk taken. The Hurwitz pessimism-optimism theory is used when environmental information is missing or unreliable, recalculations must be made for each state of the environment, the number of implemented solutions is small, and some risk is allowed. Integrating these methodologies into the possible attacks and protections, we can evaluate the methods of protection in theory and in practice. Given the possible methods of defense being none, firewall, intrusion detection tools, and link redundancy, we can evaluate each method individually and in combination. Although the most effective method in theory would be to use firewall, intrusion detection tools, and link redundancy together, in practice the cost is prohibitive. If cost is factored into the equation, then the most effective method in theory is a firewall, however this fails to take into account the possible loss if an attack is successful. Taking into account this factor as well, a firewall works in a limited number of attacks, link redundancy for higher, and the most effective is to use all three methods in combination (firewall, intrusion detection tools, and link redundancy).

**Keywords:** game theory, Wald test, Hurwitz pessimism-optimism theory, security threat, firewall, link redundancy, intrusion detection tools, effective defense

**References:**

[1] Owen G. Games theory: World, 1971.

[2] Gordon L.A., Loeb M.P, Lucyshyn W., Richardson R. 2006 CSI/FBI COMPUTER CRIME AND SECURITY SURVEY. - San Francisco: Computer Security Institute, 29 p., 2006.

[3] Shun-Chieh Lin and Shian-Shyong Tseng. Constructing detection knowledge for DDoS intrusion tolerance. Expert Systems with Applications. - Atlanta: Expert Systems with Appli-cations, - Vol.27, Issue 3, pp. 379-390, 2004.

# Binarity for almost $\omega$-categorical quite o-minimal theories

Beibut KULPESHOV

*Department of Mathematical and Computer Modeling, International Information Technology University, Almaty, Kazakhstan*
*E-mail: b.kulpeshov@iitu.kz*

**Abstract:** The present lecture deals with the notion of *weak o-minimality*, which initially deeply studied in [1]. A subset $A$ of a linearly ordered structure $M$ is *convex* if for any $a, b \in A$ and $c \in M$ whenever $a < c < b$ we have $c \in A$. A *weakly o-minimal structure* is a linearly ordered structure $M = \langle M, =, <, \ldots \rangle$ such that any definable (with parameters) subset of the structure $M$ is a finite union of convex sets in $M$.

In the following definitions (introduced in [2] and [3] respectively) we assume that $M$ is a weakly o-minimal structure, $A \subseteq M$, $M$ is $|A|^+$-saturated, and $p, q \in S_1(A)$ are non-algebraic types. We say that $p$ is not *weakly orthogonal* to $q$ ($p \not\perp^w q$) if there are an $A$-definable formula $H(x, y)$, $\alpha \in p(M)$, and $\beta_1, \beta_2 \in q(M)$ such that $\beta_1 \in H(M, \alpha)$ and $\beta_2 \notin H(M, \alpha)$. We say that $p$ is not *quite orthogonal* to $q$ ($p \not\perp^q q$) if there is an $A$-definable bijection $f : p(M) \to q(M)$. We say that a weakly o-minimal theory is *quite o-minimal* if the relations of weak and quite orthogonality for 1-types coincide.

Almost $\omega$-categoricity has been introduced in [4] and studied in [5].

**Theorem.** Any almost $\omega$-categorical quite o-minimal theory is binary.

**Keywords:** almost $\omega$-categoricity, weak o-minimality, quite o-minimality, binary theory

**2010 Mathematics Subject Classification:** 03C64

**References:**

[1] Macpherson H.D., Marker D., and Steinhorn C., "Weakly o-minimal structures and real closed fields", *Transactions of The American Mathematical Society*, Vol.352, pp. 5435-5483, 2000.

[2] Baizhanov B.S., "Expansion of a model of a weakly o-minimal theory by a family of unary predicates", *The Journal of Symbolic Logic*, Vol.66, pp. 1382-1414, 2001.

[3] Kulpeshov B.Sh., "Convexity rank and orthogonality in weakly o-minimal theories", *News of the National Academy of Sciences of the Republic of Kazakhstan, physical and mathematical series*, Vol. 227, pp. 26–31, 2003.

[4] Ikeda K., Pillay A., Tsuboi A., "On theories having three countable models", *Mathematical Logic Quarterly*, Vol.44, pp. 161-166, 1998.

[5] Sudoplatov S.V., "Classification of countable models of complete theories", Novosibirsk State Technical University, Novosibirsk, parts 1 and 2, 2014.

## Prospective study on the potential of big data in railways

Ainur KUSSAINOVA [1], Tamara ZHUKABAYEVA [1], Abai JUSSUPOV [1], Kazizat ISKAKOV [1], Zhanar ORALBEKOVA [1]

[1] *Department of Information Technology, L.N. Gumilyov Eurasian National University, Astana, Kazakhstan*
*E-mail: ainurkussainova89@gmail.com*

**Abstract:** Now new information technologies are actively implemented and used for development of different areas of industry, transportation, including the railway industry. In our work solutions to the problems of railway industry based on Big Data are presented. Key terms, basic characteristics and stages of big data are presented. Global achievements in railway industry are analysed, provided examples of foreign authors, functional block diagram of big volumes of data is developed.

**Keywords:** big data, data analysis, railway engineering.

**2010 Mathematics Subject Classification:** 68U35, 68P01

**References:**

[1] Lady Mitchell Hall, The Vocabulary of Big Data, Demystifying Data Science. Cambridge, 26th January 2015.

[2] Nii Attoh-Okine, The Future Marriage of Big Data and Railroad Engineering. // 2014 IEEE International Conference on Big Data. 2014

[3] Divakar Mysore , Shrikant Khupat, Shweta Jain. Big data architecture and patterns, Part 1: Introduction to big data classification and architecture. // IBM Corporation. -2013.

---

## An Efficient Method for the Determination of Fourth Virial Coefficient with Lennard-Jones (12-6) Potential and Its Application

Bahtiyar Akber MAMEDOV [1], Elif SOMUNCU [2], Iskender M. ASKEROV [3]

[1] *Department of Physics, Gaziosmanpasa University, Tokat, Turkey*
*E-mail: bamamedov@yahoo.com*
[2] *Department of Physics, Giresun University, Giresun, Turkey*
*E-mail: elf_smnc@hotmail.com*
[3] *Department of Physics, Giresun University, Giresun, Turkey*
*E-mail: iskender.askeroglu@giresun.edu.tr*

**Abstract:** In this work, a new theoretical approach is proposed for calculating fourth virial coefficient with Leonard-Jones potential(12-6). The established algorithm can be used to evaluate the thermodynamics properties and the intermolecular interaction potentials of liquids and gases with an improved accuracy. Note that the evaluation of the high-order virial coefficients is very valuable for accurate calculation of thermodynamic parameters. By using the suggested method, the fourth virial coefficient of $CH_4, Ar, C_2H_6$ and $SF_6$ molecules are evaluated. The calculation results are useful for accurate interpretation of the experimental data and of the determination of related physical properties [1].

**Keywords:** Virial equation of state, Fourth virial coefficient, Lennard-Jones (12-6) potential

**PACS No:** 51.30.+i, 05.10.-a

**References:**

[1] Dyer M.K., Perkyns J.S., Pettitt B.M., "A reexamination of virial coefficients of the Lennard-Jones fluid Theoretical Chemistry Accounts", Theor. Chem. Acc., Vol.105, pp. 244-251, 2011.

---

## Analyzing High-Dimensional Survival Data Using Random Forests

K. MOTARJEM[1], M. MOHAMMADZADEH [1], A. ABYAR [1]

[1] *Department of Statistics, Tarbiat Modares University, Tehran, Iran.*
*E-mail: k.motarjem@modares.ac.ir*

**Abstract:** Recently developments in genetics create cases that contain numerous covariates and few observations. These kinds of data are called High-dimensional data. In these cases, use of usual statistical methods is not appropriate. Especially in survival analysis, modeling of high-dimensional survival data is of utmost importance. One of the usual methods for analyzing survival data is the use of logistic regression. In this paper will be shown that the performance of logistic regression is not acceptable for analyzing high-dimensional survival data [1]. Generalized Breiman's Random Forests method for analyzing high-dimensional survival data, namely Random Survival Forests (RSF), has a strong efficiency for modeling survival data [2-3]. The main advantage of this method is that there is no distributional assumption on the regression model. Then the efficiency of RSF and usual methods for analyzing survival data such as logistic regression model are compared for modeling a real high-dimensional survival dataset and results are evaluated.

Finally using random forests and logistic regression models, a real dataset has been modeled and the accuracy of results are compared by Akaike Information Criterion [4]. It is shown that the random forests method and logistic regression model may presented different significant covariates. So, based on the data conditions the researchers should precisely choose the models.

**Keywords:** Random Survival Forests, Logistic regression, High-dimensional data, Survival analysis.

**2010 Mathematics Subject Classification:** 62F01, 62F02, 62F99

**References:**

[1] Mittal, S. and Madigan, D., "High-dimensional, massive sample-size Cox proportional hazards regression for survival analysis", *Biostatistics*, Vol. 15, pp. 1–15, 2014.

[2] Braiman, L., "Random forests, Machine Learn", Vol. 45, pp. 5-32, 2001.

[3] Genuer, R., "Random Forest: some methodological insights", 2008.

[4] Akaike, H., "Information theory and the extension of the maximum likelihood principle", *In: Petrov, V., Csaki, F., Editors, Proceedings of the Second International Symposium on Information Theory*, Budapest: Akailseoniaikiudo, pp. 267-281, 1973.

# Features in simulation of crystal growth using the hyperbolic PFC equation and the dependence of the numerical solution on the parameters of the computational grid

Ilya STARODUMOV [1], Nikolai KROPOTIN [2]

[1] *Laboratory of Multi-Scale Mathematical Modeling, Ural Federal University, Ekaterinburg, Russia*
*E-mail: ilya.starodumov@urfu.ru*
[2] *AO NPO MKM, Izhevsk, Russia*
*E-mail: nikokrop@yandex.ru*

**Abstract:** We investigate the three-dimensional mathematical model of crystal growth called PFC (Phase Field Crystal) in a hyperbolic modification. This model is also called the modified model PFC (originally PFC model is formulated in parabolic form) and allows to describe both slow and rapid crystallization processes on atomic length scales and on diffusive time scales [1, 2]. The modified phase field crystal model describes a continuous field of atomic density $\phi(x,t)$ and it can be expressed by the following equation:

$$\tau\frac{\partial^2\phi}{\partial t^2}+\frac{\partial\phi}{\partial t}=\nabla^2\mu, \tag{1}$$

where $t$ is the time, $\tau$ is the relaxation time of the atomic flux to its stationary state, and $\mu$ represents the chemical potential, obtained from the free-energy functional

$$\mathcal{F}[\phi,\nabla\phi,\nabla^2\phi]=\int_\Omega\left[f(\phi)-|\nabla\phi|^2+\frac{1}{2}(\nabla^2\phi)^2\right]\mathrm{d}\Omega, \tag{2}$$

associated to the domain $\Omega$. The chemical potential can be obtained as the variational derivative of the free-energy functional $\mathcal{F}$, namely

$$\mu(\phi)=\frac{\delta\mathcal{F}}{\delta\phi}=f'(\phi)+2\nabla^2\phi+\nabla^4\phi. \tag{3}$$

The function $f$ represents the homogeneous part of the free energy density. It takes on the form

$$f(\phi)=\frac{1-\epsilon}{2}\phi^2+\frac{\alpha}{3}\phi^3+\frac{1}{4}\phi^4. \tag{4}$$

Here, $\epsilon=(T_c-T)/T_c$ is the undercooling, where $T$ and $T_c$ are the temperature and critical temperature of transition, respectively. $\alpha$ is a coefficient which means a measure of metastability.

The solution of this equation is possible only by numerical methods. Previously, authors created the software package for the solution of the Phase Field Crystal problem, based on the method of isogeometric analysis (IGA) [4] and PetIGA program library [5].

Using the modified PFC model for the simulation allows description of fast and slow dynamics of structure rearrangement from homogeneous unstable phase to the emerged periodic structure [3]. We consider that if the element size is larger than lattice parameter of the emerged structure on the interface, then the type of finite elements and its shape functions predetermine the behavior of the interface front movement, because of approximating of the atomic density inside the element using inappropriate functions. In this report, we show solutions for

both stationary and non-stationary problems which confirm our assumption of a necessary condition for the computational cell size: to be at least less then the lattice parameter.

This work was supported by the Russian Science Foundation [grant number 16-11-10095]. We also thank Immers Ltd. (Michurinskiy Ave, 19, bld. 3, 119192, Moscow, Russia, www.immers.ru) for technical support and ability to run numerous calculations on the newest computing cluster of their production.

**Keywords:** crystal growth, phase field crystal, isogeometric analysis, crystallization

**2010 Mathematics Subject Classification:** 74E15, 74S05

**References:**

[1] P. K. Galenko, D. A. Danilov, and V. G. Lebedev, Physical Review E 79, 051110 (2009).

[2] P. K. Galenko, and K. R. Elder, Physical Review B 83, 064113 (2011).

[3] P. Galenko, H. Gomez, N. Kropotin, and K. Elder, Phys. Rev. E 88, 013310 (2013).

[4] J. Cottrell, T. Hughes, and Y. Bazilevs, Isogeometric Analysis: Toward Integration of CAD and FEA, Wiley, 2009.

[5] J. Bueno, I. Starodumov, H. Gomez, P. Galenko, and D. Alexandrov, Computational Materials Science 111, 310 312 (2016).

## The Moutard transformation of two-dimensional Dirac operators and the Mobius geometry

Iskander A. TAIMANOV

*Sobolev Institute of Mathematics, Novosibirsk, Russia*
*E-mail: taimanov@math.nsc.ru*

**Abstract:** We discuss the Moutard type transformation for two-dimensional Dirac operators of the form

$$\mathcal{D} = \begin{pmatrix} 0 & \partial \\ -\bar{\partial} & 0 \end{pmatrix} + \begin{pmatrix} U & 0 \\ 0 & U \end{pmatrix}$$

with $U$ a real-valued potential. In particular, we show that such a transformation has a very simple geometrical meaning. Given a solution of the equation $\mathcal{D}\psi = 0$ an a point $x_0 \in \mathbb{R}^3$, it maps the Dirac operator $\mathcal{D}$ into an operator $\tilde{\mathcal{D}}$ of the same form with another potential $\tilde{U}$ and gives an explicit procedure how to describe all solutions of the equation $\tilde{\mathcal{D}}\psi = 0$ in terms of solutions of the equation $\mathcal{D}\psi = 0$. However such a function $\psi$ and a point $x_0$ determine a surface $\Sigma$ via the Weierstrass representation such that $U$ is the potential of the surface. The inversion of $\mathbb{R}^3$ maps $\Sigma$ into a surface $\tilde{\Sigma}$ such that $\tilde{U}$ is the potential of the Weierstrass representation of $\tilde{\Sigma}$.

We use this transform for constructing blow-up solutions of the modified Novikov–Veselov equation which is a two-dimensional generalization of the modified Korteweg–de Vries equation.

**Keywords:** two-dimensional Dirac operator, Moutard transformation, inversion, modified Novikov–Veselov equation, blow-up.

**2010 Mathematics Subject Classification:** 35A30, 35Q41, 35Q53, 37K10, 53A30

**References:**

[1] Taimanov, I. A., "The Moutard transformation of two-dimensional Dirac operators and Mobius geometry". *Math. Notes*, Vol. 97, No. 1–2, pp. 124–135, 2015.

[2] Taimanov, I.A., "Blowing up the solution of the modified Novikov-Veselov equation and minimal surfaces". *Theoret. and Math. Phys.*, Vol. 182, No. 2, pp. 173–181, 2015.

## On stochastic stability of the integral manifold under permanently acting random perturbations

Marat TLEUBERGENOV [1], Gulmira VASSILINA [2]

[1] *Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: marat207@mail.ru*
[2] *Al-Farabi Kazakh National University, Kazakhstan*
*E-mail: v_gulmira@mail.ru*

**Abstract:** In this paper the stability in probability under an action of permanently acting random perturbations of the analytically given integral manifold is investigated. On the one hand these results generalize the theorems of stability under an action of permanently acting perturbations of the invariant set [1] in the class of ordinary differential equations. On the other hand they extend the theorems of stability under permanently acting perturbations of the unperturbed motion [2] to the case of invariant sets. To proof these theorems on stochastic stability under an action of permanently acting random perturbations of the analytically given invariant set we used the theorems in [3].

**Keywords:** integral manifold, stochastic stability

**2010 Mathematics Subject Classification:** 34H05,60H10

**References:**

[1] Tleubergenov M.I., *On the stability of program motion in the first approximation*, Proceedings of the Academy of Sciences of KazSSR, Series Physics and Mathematics, No.5, pp. 58-61, 1984.

[2] Khas'minskii R.Z., *Stability of Differential Systems equations with random perturbations of their parameters*, Moscow, 368 p., 1969.

[3] Tleubergenov M.I., Vassilina G.K., *On solving of the problem of stochastic stability of the integral manifold by Lyapunov's second method*, Ukrainian Mathematics Journal, Vol.68, No.1, pp. 14-27, 2016.

# Automorphisms of differential polynomial algebras

Ualbai UMIRBAEV

*Eurasian National University, Kazakhstan,*
*and Wayne State University, USA*
*E-mail: umirbaev@gmail.com*

**Abstract:** It is well known that all automorphisms of polynomial algebras and free associative algebras in two variables are tame. It is also known that polynomial algebras and free associative algebras in three variables over fields of characteristic zero have wild automorphisms.

The basic concepts of differential algebras can be found in [1]. Let $\Delta = \{\delta_1, \ldots, \delta_m\}$ be a basic set of derivations and let $A = \mathbf{k}\{x_1, x_2\}$ be the differential polynomial algebra in two free differential variables $x_1, x_2$ over an arbitrary field $\mathbf{k}$ of characteristic zero. We assume that $m \geq 2$, i.e., the number of basic derivations is at least two. Let $\theta$ be the automorphism of $A$ defined by

$$\theta(x_1) = x_1 + \delta_2(w), \quad \theta(x_2) = x_2 + \delta_1(w), \quad w = \delta_1(x_1) - \delta_2(x_2).$$

This automorphism represents an analogue of the Anick automorphism for polynomial and free associative algebras in four variables. The Anick automorphism of free associative algebras in three variables is wild [2].

We prove that $\theta$ is a wild automorphism of $A$. The question on the tameness of automorphisms of ordinary (with only one basic derivation) differential polynomial algebras in two variables remains open.

**Keywords:** Automorphisms, polynomial algebras, differential polynomial algebras, derivations.

**2010 Mathematics Subject Classification:** 12H05, 14H37, 13N15

**References:**

[1] Kolchin E.R., "Differential Algebra and Algebraic Groups", *Pure and Applied Mathematics*, Vol. 54, Academic Press, New York-London, 1973.

[2] Umirbaev U.U., "The Anick automorphism of free associative algebras", *J. Reine Angew. Math.*, Vol. 605, pp. 165–178, 2007.

## Combining goodness of fit tests for multivariate normality

Vassilly VOINOV [1], Natalya PYA [2], Rashid MAKAROV [1],
Yevgeniy VOINOV [1], Roza RAKHIMOVA[1]

[1] *KIMEP University and Institute for Mathematics and Mathematical Modeling, Almaty, Kazakhstan*
*E-mail: voinovvv@kimep.kz*
[2] *Nazarbayev University, Department of Mathematics, Astana, Kazakhstan*
*E-mail: natalya.pya@nu.edu.kz*

**Abstract:** Multivariate statistical analyses often require the data to come from multivariate normal distribution. Vector-valued tests for multivariate normality were introduced in [1]. The authors proposed two-dimensional procedures by combining several scalar tests such as non-parametric, Wald's type chi-squared tests [2] and Henze-Zirkler test. Two types of rejection regions, intersection and union, were considered there. Their preliminary simulation study to compare powers of the tests was limited to the multivariate $t$ distribution with 10 degrees of freedom as an alternative and rejection regions with equal weights for the marginal significance levels. In this work we investigate the power of the vector-valued tests as a function of the weight $w$ for the rejection regions of intersection type. Scalar tests of different origins are combined to form two-dimensional vector-valued tests. Moreover, vector-valued tests based on Neyman-Pearson intervals are developed and their performance is investigated through simulation study. This study assesses power of the proposed testing procedures implemented in a contributed package `mvnTest` [3] for the statistic software R [4].

**Keywords:** vector-valued goodness of fit tests, multivariate normality, modified chi-square type tests

**2010 Mathematics Subject Classification:** 62H15, 62H10

**References:**

[1] Voinov V., Pya N., Makarov R., Voinov Y., Rakhimova R., "Vector-valued tests for multivariate normality", *Materials of the International Conference on Statistics and its applications.*, October 16-17, Tashkent, Uzbekistan, pp. 41-48, 2015.

[2] Voinov V., Pya N., Makarov R., and Voinov Y., "New invariant and consistent chi-squared type goodness-of-fit tests for multivariate normality and a related comparative simulation study", *Communications in Statistics- Theory and Methods*, doi: 10.1080/03610926.2014.901370, 2015.

[3] Pya N., Voinov V., Makarov R., Voinov Y., "mvnTest: Goodness of fit tests for multivariate normality", *R package*, Version 1.1-0, 2016.

[4] R Development Core Team, *R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing*, Vienna, Austria, ISBN 3-900051-07-0; url: http://www.R-project.org., 2016.

# The stability of forcing companion for center of Jonsson set's fragment

Aibat YESHKEYEV, Olga ULBRIKHT

*E.A.Buketov Karaganda State University, Karaganda, Kazakhstan*
*E-mail: modth1705@mail.ru, ulbrikht@mail.ru*

**Abstract:** Let $T$ is Jonsson theory [1] complete for the existential sentences in the language $L$ and $C$ is its semantic model. Let $X$ is Jonsson subset [**?**] of $C$ and $M$ is existentially closed submodel of semantic model $C$ of considered Jonsson theory $T$, where $dcl(X) = M$. $Th_{\forall\exists}(M) = T_M$, where $T_M$ is the Jonsson fragment of Jonsson set $X$.

Let $A \subseteq C$, $\sigma_\Gamma(A) = \sigma \cup \{c_a | a \in A\} \cup \Gamma$, where $\Gamma = \{P\} \cup \{c\}$. Consider the following theory $T_\Gamma(A) = Th_{\forall\exists}(C, a)_{a\in A} \cup \{P(c)\} \cup \{"P \subseteq"\}$, where $\{"P \subseteq"\}$ is an infinite set of sentences, which says that a interpretation of symbol $P$ is positively existentially closed submodel in the signature $\sigma$. This theory is not necessarily complete. Through $S_\Gamma$ denoted the set of all $\Sigma$-completions of theory $T_\Gamma(A)$. Theory $T$ is said to be $P - \lambda$-stable, if $|S_\Gamma| \leq \lambda$ for any $A$, such that $|A| \leq \lambda$.

Consider all completions of the center $T^*$ of the theory $T$ in a new signature $\sigma_\Gamma$, where $\Gamma = \{c\}$. By virtue that $T^*$ is Jonsson theory, there exist its center and we denote it so $T^C$. If restrictions $T^C$ to the signature $\sigma$, that the theory $T^C$ becomes a complete type. This type we call a central type of theory $T$.

**Theorem 1.** Let $\lambda$ be an arbitrary infinite cardinal, $T_M$ is perfect Jonsson theory, complete for positive existential sentences. Then the following conditions are equivalent:

(1) $T^*_M$ is $P - \lambda$-stable;
(2) $\left(T^*_M\right)^F$ is $\lambda$-stable in the classical sense, where $\left(T^*_M\right)^F$ is forcing companion of theory $T^*_M$ in enriched signature;
(3) $T^C_M$ is $\lambda$-stable in the classical sense.

**Keywords:** Jonsson theory, Jonsson set, forcing companion, stable theory

**2010 Mathematics Subject Classification:** 03C05, 03C25, 03C45

**References:**

[1] Yeshkeyev A.R., "Jonsson Theories", *Publisher of the Karaganda state university*, 249 p., 2009.
[2] Yeshkeyev A.R., "On Jonsson sets and some their properties / 2014 European summer meeting of the association for Symbolic Logic Colloquium '14 Vienna, Austria, July 14-19, 2014", *The Bulletin of Symbolic Logic*, Vol. 21, No.1, pp. 99-100, 2015.

## Categorical Jonsson fragments in existentially prime convex Jonsson theory

Aibat YESHKEYEV, Nazgul SHAMATAYEVA

*E.A.Buketov Karaganda State University, Karaganda, Kazakhstan*
*E-mail: modth1705@mail.ru, naz.kz85@mail.ru*

**Abstract:** The theory Ò is called convex if for any its model $\mathfrak{A}$ and for any family $\{\mathfrak{B}_i \mid i \in \mathfrak{I}\}$ of substructures of $\mathfrak{A}$, which are models of the theory T, the intersection $\bigcap_{i\in I} \mathfrak{B}_i$ is a model of T. If it is assumed that this intersection is not empty. If this intersection is never empty, then the theory is called strongly convex.

The inductive theory Ò is called the existentially prime if

1. It has a prime algebraic (AP) model, (the class of its AP denote by $T_{AP}$)

2. Class ($E_T$) non trivial intersects with class AP, i.e. $T_{AP} \bigcap E_T \neq 0$.

Let Jonsson theory $T$ [1] is complete for the existential sentences in the language $L$ and $C$ is its semantic model.

Let X is Jonsson subset [**?**] in $T$, and $M$ is existentially closed submodel of semantic model C of considered Jonsson theory $T$, where $dcl(X) = M$ .

$Th_{\forall\exists}(M) = T_M$, $T_M$ is the Jonsson fragment of Jonsson set X [**?**].

**Theorem 1.** Let theory T will be existential prime strongly convex Jonsson existentially complete theory. Then the following conditions are equivalent:

(1) $T_M^*$ is $\omega$- categorical;

(2) $T_M$ is $\omega$ - categorical.

**Theorem 2.** Let T be an existential prime strongly convex Jonsson existentially complete theory. Then the following conditions are equivalent:

(1) $T_M^*$ is $\omega_1$ - categorical;

(2) any countable model $E_{T_M}$ has a prime algebraic extension in $E_{T_M}$.

**Keywords:** Jonsson theory, Jonsson set, Jonsson fragment

**2010 Mathematics Subject Classification:** 03C05, 03C35

**References:**

[1] Yeshkeyev A.R., "Jonsson Theories", *Publisher of the Karaganda state university*, 249 p., 2009.

[2] Yeshkeóev A.R., "On Jonsson sets and some their properties / 2014 European summer meeting of the association for Symbolic Logic Colloquium '14 Vienna, Austria, July 14-19, 2014", *The Bulletin of Symbolic Logic*, Vol.21, No.1, pp. 99-100, 2015.

# Strongly minimal fragments in existentially prime convex Jonsson theory

Aibat YESHKEYEV, Maira KASSYMETOVA

*E.A.Buketov Karaganda State University, Karaganda, Kazakhstan*
*e-mail: modth1705@mail.ru, mairushaasd@mail.ru*

**Abstract:** Let Jonsson theory $T$ [1] is complete for the existential sentences in the language $L$ and $C$ is its semantic model.

Let X is Jonsson subset [**?**] of $T$, and $M$ is existentially closed submodel of semantic model C of considered Jonsson theory $T$, where $dcl(X) = M$ .

$Th_{\forall\exists}(M) = T_M$, $T_M$ is the Jonsson fragment of Jonsson set X [**?**] .

Let $I(E_T, \aleph_0)$ denotes the number of countable existentially closed models of Jonsson theory T. On strongly minimality for Jonsson sets in the frame of the Jonsson theories one can find in [**?**].

**Theorem 1.** If theory T is the strongly minimal existentially prime convex Jonsson theory complete for existential sentences, then $T_M$ is categorical for $k \geq \aleph_1$ and $I(E_T, \aleph_0) \leq \aleph_0$ .

**Theorem 2.** If theory T is Jonsson theory complete for the existential sentences is uncountable categorical and it has Jonsson strong minimal $\mathfrak{L}$-formula, then either $T_M$ is $\aleph_0$- categorically or $I(E_{T_M}, \aleph_0) = \aleph_0$ .

**Theorem 3.** If theory T Jonsson existentially prime convex theory complete for the existential sentence is uncountable categorical, but not $\aleph_0$-categorical, then $I(E_{T_M}, \aleph_0) \geq \aleph_0$.

**Keywords:** Jonsson theory, Jonsson set, forcing companion, stable theory

**2010 Mathematics Subject Classification:** 03C05, 03C25, 03C45

**References:**

[1] Yeshkeyev A.R., "Jonsson Theories", *Publisher of the Karaganda state university*, 249 p., 2009.
[2] Yeshkeyev A.R., "On Jonsson sets and some their properties / 2014 European summer meeting of the association for Symbolic Logic Colloquium '14 Vienna, Austria, July 14-19, 2014", *The Bulletin of Symbolic Logic*, Vol.21, No.1, pp. 99-100, 2015.

# MS1

# MATHEMATICAL MODELLING

## Numerical solution to inverse source problems for hyperbolic equation

Kamil AIDA-ZADE [1,3], Anar RAHIMOV [1,2]

[1] *The Institute of Control Systems, Azerbaijan National Academy of Sciences, Baku, Azerbaijan*
*E-mail: kamil_aydazade@rambler.ru*
[2] *Aix Marseille Univ, CNRS, Centrale Marseille, Institut Fresnel, Marseille, France*
*E-mail: anar.rahimov@fresnel.fr, anar_r@yahoo.com*
[3] *Baku State University, Baku, Azerbaijan*

**Abstract:** In this work, we consider two inverse source problems with respect to processes described by boundary value problems of hyperbolic type. In particular, these classes of problems arise in the study of boundary value problems with nonlocal (integral) boundary conditions. The specific character of the considered classes of inverse problems is that the identifiable coefficients belong to the right-hand side, and they depend only on the time or only on the space. This specific character allows reducing the problem to specially built Cauchy problems with respect to a system of ordinary differential equations using the method of lines [1–3]. Thus, we do not use any iterative procedures in the approach proposed in the work. Some of the results of the carried out numerical experiments are given. The obtained results show the efficiency of the proposed approach.

**Keywords:** inverse problem, method of lines, hyperbolic equation, space and time dependent

**2010 Mathematics Subject Classification:** 35R30, 35K20, 65N40, 65N21

**References:**

[1] Aida-zade K.R., Rahimov A.B., "An approach to numerical solution of some inverse problems for parabolic equations", *Inv. Prob. in Sci. and Eng.*, Vol.22, No.1, pp. 96-111, 2014.

[2] Aida-zade K.R., Rahimov A.B., "Solution to classes of inverse coefficient problems and problems with nonlocal conditions for parabolic equations", *Dif. Eq.*, Vol.51, No.1, pp. 83-93, 2015.

[3] Aida-zade K.R., Rahimov A.B., "Numerical solution to a class of inverse source problems for a parabolic equation with integral overdetermination conditions", *Abstracts of the Eighth International Conference "Inverse Problems: Modeling and Simulation' (Fethiye, Turkey, May 23-28, 2016)*, Ed. A. Hasanoglu, A. Neubauer et al., Izmir: Izmir University Publication, p. 83, 2016.

# Implementation of scalar filtered density function for large eddy simulation of turbulent reacting flow using a high-order discontinuous Galerkin method

Aidyn AITZHAN [1], Medet INKARBEKOV [1], Asetzhan KOLDAS [1], Aidarkhan KALTAYEV [1]

[1] *Department of Mechanics, Al-Farabi Kazakh National University, Almaty, Kazakhstan*
*E-mail: aitzhan.aidyn.b@gmail.com*

**Abstract:** Turbulence is interesting and useful flow phenomena of nature. In case of reacting flows, turbulence promotes effective mixing of reactants. Due to fine mixing a chemical reaction is going to be highly productive. Hence, turbulent reactive flows are especially needed in industry. Nowadays engineering problems require highly accurate simulation of turbulent reacting flow. Therefore mathematical models and numerical methods must be sufficiently accurate.

In present work, a Filtered Density Function methodology (FDF) coupled with high-order Discontinuous Galerkin (DG) method is applied for Large Eddy Simulation (LES) of turbulent reacting flow. The FDF method has proven to be very effective for LES of turbulent reactive flows [1, 2] from the other side DG method is highly accurate and useful method, and at the same time is compact and relative easy [3]. In order to compare results there are provided results of computations of discontinuous Galerkin LES/FDF and finite difference direct numerical and large eddy simulations. In addition a DG-LES/FDF numerical code is parallelized with CUDA technology, which accelerated computations more that 10 times.

**Keywords:** filtered density function, high-order method

**2010 Mathematics Subject Classification:** 76F25, 76F65

**References:**

[1] Colucci P.J., Jaberi F.A., Givi P., Pope S.B., "Filtered density function for large eddy simulation of turbulent reacting flows", *Phys. Fluids*, Vol.10, No.2, pp. 499-515, 1998.

[2] Jaberi F.A., Colucci P.J., James S., Givi P., Pope S.B., "Filtered mass density function for large-eddy simulation of turbulent reacting flows", *J. Fluid Mech.*, Vol.401, pp. 85-121, 1999.

[3] Schaal K., Bauer A., Chandrashekar P., Pakmor R., Klingenberg C., Springel V., "Astrophysical hydrodynamics with a high-order discontinuous Galerkin scheme and adaptive mesh refinement", *MNRAS*, Vol.453, No.4, pp. 4278-4300, 2015.

## Dynamics of the massif in the vicinity of the tunnel any profile of section under action of transport loadings

Lyudmila ALEXEYEVA [1], Makpal AHMETZHANOVA [1], Gulnar KAYSHIBAYEVA [1]

[1] *Institute of Mathematics and Mathematical Modelling, Almaty, Kazakhstan*
*E-mail: alexeeva47@mail.ru, makpal11@gmail.com, gulnarkaishi@mail.ru*

**Abstract:** The mathematical model has been considered of dynamics of not supported tunnels of any cross section of a deep pawning under the action of the transport loadings of the established profile moving with a constant subsonic speed (a speed is smaller then the speeds of elastic waves in the massif ). For the circular tunnels such problems were considered in [1]. Here with use General Functions Method [2,3] the computer programs of calculation of the motion of the surrounding pedigree massif are developed and some calculations of dynamics of the surrounding massif are given in the vicinity of tunnels of circular and vaulted profiles of cross section.

The considered model and a computer programs allow to research the deflected mode of the massif in vicinities of transport tunnels taking into account a look and the speed of transport loading, mechanical parameters of the pedigree massif and design features of the construction. Especially it is important at design of lines of subways in an urban environment for an assessment of their seismic impact on land constructions and buildings.

**Keywords:** elastodynamics, boundary value problem, transport loading

**2010 Mathematics Subject Classification:** 74B05,45E99

**References:**

[1] Erzhanov Zh.S., Aitaliyev Sh.M., Alexeyeva L.A., *Dynamics of tunnels and underground pipelines*, Alma-Ata: Science, 240 p., 1989.

[2] Alexeyeva L.A., Boundary Element Method of Boundary Value Problems of Elastodynamics by Stationary Running Loads // Int. J. Engineering Analysis with Boundary Element, No.11., pp. 37-44, 1998.

[3] Alexeyeva L.A., The singular boundary integral equations of BIE of an elastodynamics in case of the subsonic running loadings // Differential equations, Vol.46, No.4., pp. 512-519, 2010.

# An asymptotical method for determining hydraulic resistance coefficient of gas-lift process

F.A. ALIEV [1], N.S. HAJIEVA [1], N.A. SAFAROVA [1], M.F. RAJABOV [2]

[1] *Institute of Applied Mathematics, Baku State University, Baku, Azerbaijan*
*E-mail: f_aliev@yahoo.com, nazile.m@mail.ru, narchis2003@yahoo.com*

[2] *Institute of Control Systems of ANAS, Baku, Azerbaijan*
*E-mail: mmamed@yahoo.com*

**Abstract:** A mathematical model in oil production is formulated to determine the coefficient of hydraulic resistance [1], during the motion of the gas-liquid mixture in the lift. As it is known, the motions of the gas and gas-liquid mixture in the tubes are described by the system of partial differential equations of hyperbolic type in the asymptotic case [2]:

$$\left\{ \begin{array}{l} \frac{\partial P}{\partial t} = -\frac{c^2}{F}\frac{\partial Q}{\partial z}\varepsilon, \\ \frac{\partial Q}{\partial t} = -F\frac{\partial P}{\partial z}\varepsilon - 2aQ, \end{array} \right. \tag{1}$$

where the parameters in (1) defined as [3]. Applying the lines method, we obtain from (1)

$$\dot{x} = (A_0(\lambda_c) + A_1\varepsilon)x + B\varepsilon u + V\varepsilon,$$

with initial condition $x_0 = \left[P_1^0, Q_1^0,\ P_2^0,\ Q_2^0\right]'$. It is required to minimize the functional

$$f(\lambda_c) = \sum_{i=1}^{N}\left[\tilde{Q}_2^i - Q_2^i\right]^2.$$

On a concrete example the comparison of the values of the obtained hydraulic resistance coefficient with the statistical value of hydraulic resistance is given. It is shown that they differ from each other to the order $10^{-3}$.

**Keywords:** gas-lift, the coefficient of hydraulic resistance, lines method

**2010 Mathematics Subject Classification:** 49J15, 49J35

**References:**

[1] Aliev F.A., Ismailov N.A., "Inverse problem to determine the hydraulic resistance coefficient in the gaslift process", *Appl. Comput. Math.*, Vol.12, No.4, pp. 306-331, 2013.

[2] Mutallimov M.M, Askerov I.M., Ismailov N.A., Rajabov, M.F., "An asympotical method to construction a digital optimal regime for the gaslift process", *Appl. Comput. Math.*, Vol.9, No.1, pp. 77-84, 2010.

[3] Mukhtarova N.S., "Algorithm to solution the identification problem for finding the coefficient of hydraulic resistance in gas-lift process", *Proc. of IAM*, Vol.4, No.2, pp. 206-213, 2015.

## Modeling and calculation of flotation process in one-dimensional formulation

Tulegen AMANBAEV [1], Gamidulla TILLEUOV [2], Bibigul TULEGENOVA [2]

[1] *Pushkin 125, 050010 Almaty, Kazakhstan, Institute of Mathematics and Mathematical Modeling*
*E-mail: tulegen_amanbaev@mail.ru*
[2] *M. Auezov South Kazakhstan State University, Kazakhstan*
*E-mail: gamidulla_nca@mail.ru, tulegenovaba@mail.ru*

**Abstract:** In the framework of the assumptions of the mechanics of the multiphase media is constructed a mathematical model of the flotation process in the dispersed mixture of liquid, solid and gas phases, taking into account the degree of mineralization of the surface of the bubbles. Application of the constructed model is demonstrated on the example of one-dimensional stationary flotation and it is shown that the equations describing the process of ascent of the bubbles are singularly perturbed ("hard"). Analyzed the effect of size and concentration of bubbles and the volumetric content of dispersed particles on the flotation process.

Nowadays the flotation method is widely used for division of disperse mixes in various industries [1-7] that stipulates an urgency of a considered problem. Modeling of the flotation process in the three-phase environment is a difficult task. Theoretical study of the process of flotation is based, in the main, on the kinetic model [1-5] where many important hydrodynamic factors are not considered. Consequently, it is necessary to construct the mathematical model of the flotation process with considering the influence of the hydrodynamic factors [8].

**Keywords:** Flotation process, multiphase media, bubble

**2010 Mathematics Subject Classification:** 76T10, 35Q35

**References:**

[1] Klassen V.I., and Mokrousov V.A., *Introduction in the flotation theory*, Metallurgizdat, Moskow, 1959.
[2] Klassen V.I., and Glembotsky V.A., *Flotation*, Nedra, Moskow, 1973.
[3] Rubinshtejn J., and Philippov J., *Kinetiks of a flotation*, Nedra, Moskow, 1980.
[4] Bogdanova O.S., *The theory and technology of flotation of ores*, Nedra, Moskow, 1990.
[5] Derjagin B.V., Duhin S., and Rulev N.N., *Successes of chemistry* Vol.51(1), pp. 92-118, 1982.
[6] A. Sulaymon, and A. Mohammed, *Inter. Journal of Chem.Reac.Ing.* Vol.8, 48, 2010.
[7] M. Kostoglou, T. Karapantsios, and K. Matis, *Ind. Eng. Chem. Res.* Vol.46, pp. 6590-6599, 2007.
[8] T. Amanbaev and S. Antony, *Theor. Found. Chem.Eng.* Vol.45(6), pp. 876-884, 2011.

## The Fujita type a critical exponent for a double nonlinear parabolic equation and system

Mersaid ARIPOV [1]

[1] *Department of Informatics and Computer Analysis, National University of Uzbekistan, Tashkent, Uzbekistan*
*E-mail: mirsaidaripov@mail.ru*

**Abstract:** Consider in $Q = \{(\mathrm{t}, \mathrm{x}) : \mathrm{t} > 0,\ x \in R^N\}$the following problem Cauchy to a degenerate parabolic system with double nonlinearity

$$\begin{cases} \frac{\partial u}{\partial t} = \nabla \left( v^{m_1 - 1} |\nabla u|^{p-2} \nabla u \right) + div(c(t)u) + u^{\beta_1}, \\ \frac{\partial v}{\partial t} = \nabla \left( u^{m_2 - 1} |\nabla v|^{p-2} \nabla v \right) + div(c(t)v) + v^{\beta_2} \end{cases} \tag{1}$$

$$u(0,x) = u_0(x) \geq 0,\ v(0,x) = v_0(x) \geq 0,\ x \in R^N \tag{2}$$

This problem describes the different nonlinear processes [1-4]. The solution of the problem (1), (2) depending on value of numerical parameters $m_i$, $\beta_i$, $i = 1, 2, p$and functions $c(t)$, $\gamma(t)$ have a property of a finite speed perturbation, space localization, blow-up and so on [1,2]. Now these properties of the solution to the problem Cauchy for a different nonlinear equation in the case single equation and system intensively studied by many authors (see [1-4] and the literature therein). Many works (see [1-3] and literature therein) are devoted to the Fujita type critical exponents for semi-linear system (1) ($m_i = 1,\ \beta_i,\ i = 1, 2, p = 2$). In this case in [4] the following condition $(\beta_i + 1)/\beta_1\beta_2 - 1 = N/2,\ i = 1, 2$ of the critical exponents were established.

In this work an algorithm for establishing value of a critical exponent for the system (1) and for some other degenerate system based on approximately self similar approach is suggested. Using comparison principle the condition of a global solvability, property a finite speed of perturbation, space localization of the solution, blow-up is established. The following condition of a global solvability
$\tau(t)\bar{u}^{\beta_1 - (p-2)}\bar{v}^{-(m_1-1)} < (\beta_1 - 1)N/p$, $\tau(t)\bar{u}^{-(m_2-1)}\bar{v}^{\beta_2-(p-2)} < (\beta_2 - 1)N/p$,
where $\bar{u}(t) = (T + \int \gamma(t)dt)^{-1/(\beta_1-1)}$, $\bar{v}(t) = (T + \int \gamma(t)dt)^{-1/(\beta_2-1)}$, $T \geq 0$,
$\tau(t) = \int [\bar{u}(t)]^{(p-2)}[\bar{v}(t)]^{(m_1-1)}dt$
which consists in particular the result of th work [4] is established.

**Keywords:** double nonlinear, parabolic system, property, critical exponent

**2010 Mathematics Subject Classification:** 35B20, 35B33

**References:**

[1] Samarskii A.A., Galaktionov V.A., Kurdyomov S.P., Mikhailov A.P., "Blow-up in quasilinear parabolic equations", *Berlin, Walter de Grueter*, 4, 535 p., 1995.

[2] Aripov M., Sadullaeva Sh.A., "An asymptotic analysis of a self-similar solution for the double nonlinear reaction-diffusion system", *J. Nanosystems: physics, chemistry, mathematics*, 6(6), pp. 793-802, 2015.

[3] Aripov M., Sadullaeva Sh.A., "Qualitative properties of solutions of a doubly nonlinear reaction-diffusion system with a source", *Journal of Applied Mathematics and Physics*, 3, pp. 1090-1099, 2015.

[4] Escobedo M., Herrero M. A, "Boundedness and Blow Up for a Semilinear Reaction-Diffusion System", *Journal of Differential Equations*, 89, pp. 176-202, 1991.

## The features of a non-stationary state of stress in the elastic multisupport construction

Nurgali ASHIRBAYEV[1], Zhansaya ASHIRBAYEVA[1],
Azimkhan ABZHAPBAROV[1], Manat SHOMANBAYEVA[1]

[1] *M. Auezov South Kazakhstan State University, Kazakhstan*
*E-mail: ank_56@mail.ru, saya_270681@mail.ru, azeke_55@mail.ru, mtshomanbaeva@mail.ru*

**Abstract:** The paper deals with the problem of propagation of unsteady elastic waves in an elastic multisupport construction, which is a rectangular strip. The mixed problem is formulated in terms of the stress and velocity and is numerically modeled using an explicit difference scheme through computation based on the method of spatial characteristics. The main objective of this study is to analyze the impact of the gap in the boundary conditions on the propagation of wave processes in the internal points of the studied elastic medium. The concentration of dynamic stresses was investigated in the vicinity of the gap of the boundary conditions. The results of the study were brought to the numerical solution.[1-8]

**Keywords:** Speed, voltage, load, plane strain, stress concentration, numerical solution, wave propagation

**2010 Mathematics Subject Classification:** 65L12, 74G70

**References:**

[1] V.N. Kukudzhanov, *Bulletin of the Nizhny Novgorod University. N.I. Lobachevsky* **4**(4), 1557–1558 (2011).

[2] L.A. Alexeeva, and B.T. Sarsenov, *AIP Conference Proceedings* **1676**, 020055 (2015); http://dx.doi.org/10.1063/1.4930481.

[3] N. Ashirbayev, Zh. Ashirbayeva, and M. Shomanbayeva, *AIP Conference Proceedings* **1676**, 020067(2015); http://dx.doi.org/10.1063/1.4930493.

[4] E.L. Kuznetsova, D.V. Tarlakovskii, and G.V. Fedotenkov, *Mech. Solids* **46** (5), 779–787 (2011).

[5] M.V. Dolotov, I.D. Kill', and Yu.G. Limonchenko, *J. Appl. Math. Mech.* **79** (2), 179–185 (2015).

[6] Zh.S. Erzhanov, T.D. Karimbayev, and T.B. Bayteliyev, *The stress waves in homogeneous and heterogeneous environments.* - Almaty: Gylym, 1998.

[7] R.J. Clifton, *Quart. Appl. Math.* **25**(1), 97–116 (1967).

[8] N.K. Ashirbayev, *Bulletin of KazNPU. Abay, "Physics and mathematics" series* **1** (29), 50–54 (2010).

## Modeling and solving the two-dimensional non-stationary problem in an elastic body with a rectangular hole

Nurgali ASHIRBAYEV[1], Zhansaya ASHIRBAYEVA[1],
Turlybek SULTANBEK[1], Raina BEKMOLDAYEVA[1]

[1] *M. Auezov South Kazakhstan State University, Kazakhstan*
*E-mail: ank_56@mail.ru, saya_270681@mail.ru, sultur_55@mail.ru, raina_b@mail.ru*

**Abstract:** In this work we consider the problem of the propagation of non stationary stress waves in an elastic body with a rectangular hole in the linear formulation. The wave process is caused by applying an external dynamic load on the front boundary of the rectangular region and the lateral boundaries are free of the stress. The lower boundary of the rectangular region is rigidly fixed, and the contour of the rectangular hole is free from the stress. The problem is solved by using the difference method of the spatial characteristics. On the basis of the developed numerical methods it is obtained the computational finite - difference relations of the dynamic problems at the corner points of the rectangular hole, where the first and second derivatives of the unknown functions have a discontinuity of the first kind. We analyze the dynamic stress fields in an elastic body with a rectangular hole and we studied the concentration of dynamic stresses in the vicinity of the corner points of the rectangular opening.[1-8]

**Keywords:** speed, stress, load, plane strain, stress concentration, numerical solution, wave propagation

**2010 Mathematics Subject Classification:** 65L12, 74G70

**References:**

[1] Bayanov E.V., Gulidov A.I., *Journal of Applied Mechanics and Technical Physics* 5, pp. 155-162, 2011.

[2] Alexeeva L.A., Sarsenov B.T., *AIP Conference Proceedings*, Vol.1676, 020055, 2015. http://dx.doi.org/10.1063/1.4930481.

[3] Ashirbayev N., Ashirbayeva Zh., and Shomanbayeva M., *AIP Conference Proceedings*, Vol.1676, 020067, 2015. http://dx.doi.org/10.1063/1.4930493.

[4] Kochaev A.L., Brazhe R.A., *Acta mechanika* Vol.1-2, pp. 193-198, 2011.

[5] Frishter L.Y., *Proceedings of the International scientific-practical conference "Engineering Systems - 2011", Moscow* Vol.2, pp. 3-9, 2011.

[6] Clifton R.J., *Quart. Appl. Math.*, Vol.25, pp. 97-116, 1967.

[7] Ashirbayev N.K., Nyssanov E.A., and Ashirbayeva Zh.N., *Scientific journal Bulletin of ENU. L.N. Gumilev, Astana*, Vol.2, Part 1, 3, pp. 1-36, 2015.

## On the stability estimation of differential-difference analogue of the integral geometry problem with a weight function

Galitdin B. BAKANOV [1]

[1] *Department of Mathematics, Akhmet Yasawi University, Turkestan, Kazakhstan*
*E-mail: galitdin.bakanov@ayu.edu.kz*

**Abstract:** A great deal of the work has been devoted to problems of integral geometry, that is, the problems of determining a function from its integral along a given family of curves. One of the stimuli for studying such problems is their connection with multidimensional inverse problems for differential equations [1]. In some inverse problems for hyperbolic equations were shown to reduce to integral geometry problems and, in particular, a problem of integral geometry was considered in the case of shift-invariant curves. Mukhometov [2] showed the uniqueness and estimated the stability of the solution of a two-dimensional integral geometry on the whole. His results were mainly based on the reduction of the two-dimensional integral geometry problem

$$V(\gamma, z) = \int\limits_{K(\gamma,z)} U(x,y)\rho(x,y,z)ds, \quad \gamma \in [0,l], \quad z \in [0,l] \tag{1}$$

where $U \in C^2(\overline{D}), \rho(x,y,z)$ is a known function to the boundary value problem

$$\frac{\partial}{\partial z}\left(\frac{\partial W}{\partial x}\frac{\cos\theta}{\rho} + \frac{\partial W}{\partial y}\frac{\sin\theta}{\rho}\right) = 0, \quad (x,y,z) \in \Omega_1, \tag{2}$$

$$W(\xi(\gamma), \eta(\gamma), z) = V(\gamma, z), \quad V(z,z) = 0, \quad \gamma, z \in [0,l]. \tag{3}$$

Differential - difference analogue of the problem (2)-(3) are studied. The stability estimate for the considered problem (2)-(3) are obtained.

**Keywords:** integral geometry problem, differential-difference problem, stability, uniqueness.

**2010 Mathematics Subject Classification:** 34K29

**References:**

[1] Lavrent'ev M.M., and Romanov V.G. "Three linearized inverse problems for hyperbolic equations", *Soviet Math. Dokl.*, Vol.7, pp. 1650-1652, 1966.

[2] Mukhometov R.G., "On a problem of integral geometry" *In: Mathematical Problems of Geophysics*, Vol.6:2, pp. 212-242, 1975 (in Russian).

## Soft matrices on soft multisets in an optimal decision process

Arzu ERDEM COSKUN [1], Cigdem GUNDUZ ARAS [1], Ayse SONMEZ [2], Huseyin CAKALLI [3]

[1] *Department of Mathematics, Kocaeli University, Kocaeli, Turkey*
*E-mail: erdem.arzu@gmail.com, caras@kocaeli.edu.tr*
[2] *Department of Mathematics, Gebze Institute of Technology, Kocaeli, Turkey*
*E-mail: asonmez@gyte.edu.tr*
[3] *Department of Mathematics, Maltepe University, İstanbul, Turkey*
*E-mail: huseyincakalli@maltepe.edu.tr, hcakalli@gmail.com*

**Abstract:** In this paper, we introduce a concept of a soft matrix on a soft multiset, and investigate how to use soft matrices to solve decision making problems. An algorithm for a multiple choose selection problem is also provided. Finally, we demonstrate an illustrative example to show the decision making steps.

**Keywords:** Soft matrix, soft multiset, soft max-min decision making

**2010 Mathematics Subject Classification:** 03E75, 03E99

**References:**

[1] Molodtsov D., "Soft set theory first result", *Computers and Mathematics with Applications*, Vol.37, pp. 19-31, 1999.

[2] Gunduz C., and Bayramov S., "Some Results on Fuzzy Soft Topological Spaces", *Mathematical Problems in Engineering*, pp. 1-10, 2013.

[3] Gunduz C., Sonmez A., and Cakalli H., "On Soft Mappings", *arXiv*, 1305.4545, pp. 1-12, 2013.

[4] Erdem A., Gunduz C., Sonmez A., and Cakalli H., "The use of soft matrices on soft multisets in an optimal decision process", *arXiv*, :1401.7543, [math.GM], pp. 1-16, 2014.

[5] Basu T., Mahapatra N., and Mondal S., "Different Types of Matrices in Fuzzy Soft Set Theory and Their Application in Decision Making Problems", *IRACST - Engineering Science and Technology: An International Journal ESTIJ*, Vol.2, No.3, pp. 389-398, 2012.

[6] Cagman N., and Enginoglu S., "Soft matrix theory and its decision making", *Computers and Mathematics with Applications*, Vol.59, pp. 3308-3314, 2010.

[7] Mondal J., and Roy T., "Theory of Fuzzy Soft Matrix and its Multi Criteria in Decision Making Based on Three Basic t-Norm Operators", *International Journal of Innovative Research in Science Engineering and Technology*, Vol.2, No.10, pp. 5715-5723, 2013.

[8] Vijayabalaji S., and Ramesh A., "A New Decision Making Theory in Soft Matrices", *International Journal of Pure and Applied Mathematics*, Vol.86, No.3, pp. 927-939, 2013.

[9] Alkhazaleh S., and Salleh A., "Soft Multisets Theory", *Applied Mathematical Sciences*, Vol.5/72, pp. 3561-3573, 2011.

## On a problem of reconstruction of a discontinuous function by its Radon transform

Evgeny DEREVTSOV[1], Svetlana MALTSEVA[2],
Ivan SVETOV[3], Murat SULTANOV[4]

[1,2,3] *Sobolev Institute of Mathematics, Russia*
[1,2,3] *Novosibirsk State University, Russia*
[4] *H.A.Yassawe International Kazakh-Turkish University, Kazakhstan*
*E-mail:* [1]*dert@math.nsc.ru,* [2]*sv_maltseva@mail.ru,*
[3]*svetovie@math.nsc.ru,* [4]*murat.sultanov@iktu.kz*

**Abstract:** A problem of reconstruction of a discontinuous function by its Radon transform is considered. Almost all of mathematical models of tomography include objects with discontinuous properties, but the vast majority of the algorithms applying for reconstruction of a searched-for function require its smoothness or at least continuity. Thus an employment of such algorithms for reconstruction of discontinuous functions leads to errors that increase the errors arising usually at the reconstruction of smooth functions in 3–10 times. Hence a development of the algorithms and numerical tools for reconstruction of discontinuous objects by tomographic data is an important task. An expression "breaks recovery" includes few problems [1–3]. First of them is the problem of visualization of a set of breaking points. This problem consists in obtaining the image, on which the set of breaking points of a function can be recognize easily. A result of the second problem (the identification of discontinuities) should be mathematical description of the points or lines of discontinuities. The third problem consists in determination of jump values.

The problem of reconstruction of a set of breaking points is connected closely with the problem of reconstruction of discontinuous functions. One of approaches to the numerical solution for the problem of reconstruction of discontinuous functions reconstruction consists in the next sequential steps: a) a visualization of a set of breaking points; b) an identification of this set; c) a determination of jump values; d) an eliminations of discontinuities. We consider three of listed problems except for the problem of jump values determining. The problems are investigated by mathematical modeling with usage of numerical experiments. The descriptions of simulations and their results are represented. The results of simulation are satisfactory and allow us to hope for the further development of the approach.

The authors were supported partially by Ministry of Education and Science of the Republic of Kazakhstan (project 0115PK00681).

**Keywords:** tomography, Radon transform, discontinuous function, visualization of breaking points, identification, elimination of breaks, simulation

**2010 Mathematics Subject Classification:** 44A12, 65R32, 94A08

**References:**

[1] Louis, A.K., "Feature reconstruction in inverse problems", *Inverse Problems*, Vol.27, No.6, art. no 065010, 2011.

[2] Derevtsov, E.Yu., "Some approaches to a reconstruction of a singular support of scalar, vector and tensor fields by their known tomographic data", *Siberian Electronic Mathematical Reports*, Vol.5, pp. 632–646, 2008. (in Russian).
[3] Derevtsov, E.Yu., Pickalov, V.V., "Reconstruction of vector fields and their singularities from ray transform", *Numerical Analysis and Applications*, Vol.4, No.1, pp. 21–35, 2011.

## A new combined DG-MC solver for large eddy simulation of reacting turbulent flows

Medet INKARBEKOV [1], Aydyn AYTZHAN [1], Aidarkhan KALTAYEV [1]

[1] *Institute of Mathematics and Mechanics*
*Al-Faraby Kazakh National University, Kazakhstan*
*E-mail: inkarbekovm@gmail.com*

**Abstract:** In the presented article, a new computational methodology is developed and implemented to simulate turbulent reacting mixing layers of two opposite parallel flows. The large eddy simulation (LES) approach and the methodology named "filtered density function" (FDF) are used for turbulence modelling. In the FDF methodology, the effects of the unresolved scalar fluctuations are taken into account by considering the probability density function of subgrid scale scalar quantities. A big advantage of the methodology is that in the FDF transport equation the effect of chemical reactions appears in a closed form. The base filtered transport equations are solved numerically by a Discontinuous Galerkin (DG) method. The FDF transport equation is solved by a particle based lagrangian Monte-Carlo (MC) method. The power of DG methods is that it can provide high order accuracy with fewer degrees of freedom. It is shown that the developed DG-MC solver is a powerful tool for large eddy simulation of reacting turbulent flows.

Throughout this note we use techniques from the works [1], [2].

**Keywords:** : turbulence, filtered density function, large-eddy simulation, Monte Carlo method, discontinuous Galerkin methods

**2010 Mathematics Subject Classification:** 76F10, 76F65, 76D06, 65C05, 65L60

**References:**

[1] Givi P., "Filtered density function for subgrid scale modeling of turbulent combustion", in *AIAA Journal*, Vol.44, No.1, pp. 16-23, Jan. 2006.
[2] Yilmaz S.L., Ansari N., Pisciuneri P.H., Nik M.B., Otis C.C., and Givi P., "Advances in FDF Modeling and Simulation", in *47th AIAA/ASME/SAE/ASEE Joint Propulsion Conference & Exhibit*, pp. 1-11, San Diego, CA, AIAA, AIAA-2011- 5918, 2011.

## On a method of mathematical modeling of the pyrolysis process to produce nanomaterials from natural gas

Bakhtiyar ISMAILOV [1], Khairulla ISMAILOV [2], Saule MELDEBEKOVA [3]

*Department of Mathematics, South Kazakhstan State University of M. Auezov, Shymkent, Kazakhstan*
*E-mail: ismailb@mail.ru*

**Abstract:** In this paper we developed a method for implementing a mathematical model of the methane catalytic pyrolysis to produce nanomaterials in periodic and continuous mode. It is shown that the Laplace transform method is mostly appropriate for this case, which is applicable both for the differential equations for the solid-phase components and for calculating the periodic mode. The adequacy of the resulting solution with known experimental data has been checked. Mathematic formulation is of the following mode :

$$\frac{\int \partial C_i}{\partial t} + v_1 \frac{\partial C_i}{\partial x} = D_i \frac{\partial^2 C_i}{\partial x^2} + D_i \frac{\partial^2 C_i}{\partial r^2} + \frac{D_i}{r} \frac{\partial C_i}{\partial r} + \alpha_{i-1} C_{i-1} + \beta_{i+1} C_{i+1} + J_i,$$

$$C_i(t = 0, x, r) = C_i^0(x, r),$$

$$C_i(t = 0, x, r) = C_i^L(r),$$

$$\frac{\partial C_i}{\partial x} \mid_{x=1} = 0,$$

$$D_i \frac{\partial C_i}{\partial r} \mid_{r=0} = \sum_{j=1}^{M_2} v_j^i W_j, if x = I_{ap}/2,$$

$$D_i \frac{\partial C_i}{\partial r} \mid_{r=0} = 0, if x \neq I_{ap}/2,$$

$$D_i \frac{\partial C_i}{\partial r} \mid_{r=D_{ap}/2} = 0,$$

where $i = 1, 2, ..., n; I_{ap}, D_{ap}$- device length and diameter, $m; \tilde{N}_i^L$ - gas constituent original concentration at the reactor inlet, mole/ $m^3; v_i^j$ gas phase $i$-component stoichiometric coefficient in the $j$-surface reaction; $W_j$- $j$-surface reaction velocity mole/$(m^3, s)$ (1) - modified equation system for the gas constituent concentrations with the optional term, allowing for the interim radicals concentrations impact; $\alpha_0; \alpha_{n+1} = 0$; (2) - initial conditions. In this Article we prove the solvability of the model (1) - (7), solubility and the iterative method stability in solving the system differential equations (1).

**Keywords:** catalytic pyrolyze, differential equation, Laplace transform. **2010 Mathematics Subject Classification:** 94B05, 94B15

# Mathematical modeling of the thermal effect on the aquatic environment from thermal power plant by using two water discharged pipes

Alibek ISSAKHOV [1]

[1] *Department of Mathematical and Computer Modelling, al-Farabi Kazakh National University, Almaty, Kazakhstan*
*E-mail: alibek.issakhov@gmail.com*

**Abstract:** In this work presents a mathematical modeling of the thermal load on the aquatic environment by using two water discharged pipes of TPP. It is solved by the Navier-Stokes and temperature transport equations for an incompressible fluid in a stratified medium. [1], [3], [4]. Numerical method is based on the projection method [2] which was approximated by the finite volume method. The numerical solution of the equation system is divided into four stages. At the first step it is assumed that the momentum transfer was carried out only by convection and diffusion. Intermediate velocity field is solved by 5-step Runge-Kutta method. At the second stage, the pressure field is solved by found the intermediate velocity field. Poisson equation for the pressure field is solved by Jacobi method. The third step is assumed that the transfer is carried out only by pressure gradient. The fourth step of the temperature transport equation is also solved as momentum equations. The obtained numerical results of stratified flow for two water discharged pipes were compared with experimental data and with numerical results for one water discharged pipe. General thermal load in the reservoir-cooler decreases comparing with one water discharged pipe and revealed qualitatively and quantitatively approximately the basic laws of hydrothermal processes.

**Keywords:** Stratified medium, Navier-Stokes equation, Thermal discharge, Operational capacities of thermal power plant

**2010 Mathematics Subject Classification:** 76D05, 76D50, 65N08

**References:**

[1] Fossati M., Piedra-Cueva I., "Numerical modelling of residual flow and salinity in the Rio dela Plata", *App. Math. Modelling*, Vol.32(6), pp. 1066–1086, 2008.

[2] Chorin A.J., "Numerical solution of the Navier–Stokes Equations", *Math.Comp*, Vol.22, pp. 745-762, 1968.

[3] Issakhov A., "Mathematical modeling of the discharged heat water effect on the aquatic environment from thermal power plant", *Int. J. of Nonlin. Sc. and Num. Simulation*, Vol.16(5), pp. 229–238, 2015.

[4] Issakhov A., "Mathematical modeling of the discharged heat water effect on the aquatic environment from thermal power plant under various operational capacities", *App. Math. Modelling*, Vol.40(2), pp. 1082–1096, 2016.

## Mathematical modeling of non-equilibrium sorption

Ibragim KALIEV [1], Saltanbek MUKHAMBETZHANOV [1]
Gulnara SABITOVA [2], Anghyz SAKHIT [1]

[1] *Al-Farabi Kazakh National University, Almaty, Kazakhstan*
*E-mail: kalievia@mail.ru*
[2] *Sterlitamak Branch of Bashkir State University, Russia*
*E-mail: sabitovags@mail.ru*

**Abstract:** We consider the system of equations modeling the process of non-equilibrium sorption. Difference approximation of differential problem by the implicit scheme is formulated. The solution of the difference problem is constructed using the sweep method. Based on the numerical results we can conclude the following: when the relaxation time decreases to 0, then the solution of non-equilibrium problem tends with increasing time to solution of the equilibrium problem.

**Keywords:** The system of equations of non-equilibrium sorption, difference approximation, the implicit scheme, sweep method, numerical experiments.

**2010 Mathematics Subject Classification:** 35Q35, 65M06, 76S05

## Exact solution of two phase spherical Stefan problem with two free boundaries

Alex KAVOKIN [1,2], Targyn NAURYZ [3], Nazerke BIZHIGITOVA [3]

[1] *Institute of Mathematics and Mathematical Modelling, Almaty, Kazakhstan*
*E-mail: kavokin_alex@yahoo.com*
[2] *Al-Farabi Kazakh National University, Kazakhstan*
[3] *Department of Mathematics and Natural Science, Suleyman Demirel University, Almaty, Kazakhstan*

**Abstract:** Solution of the heat equation in a spherical domain with two free boundaries (two-phase Stefan problem) when one of the subdomains degenerates at the initial time is considered. The use of conventional finite-difference methods in these cases is not expedient because of the degenerate domain. The solution is found in the form of combination of Integral Error functions series, [1] and then recurrent solvability of nonlinear algebraic equations for determining the coefficients of the series is proved.

The mathematical model includes the spherical heat distribution with constant coefficients for each of the phases

$$\frac{\partial\theta_1}{\partial t} = a_1^2\big(\frac{\partial^2\theta_1}{\partial r^2} + \frac{2}{r}\frac{\partial\theta_1}{\partial r}\big), \qquad \alpha(t) < r < \beta(t), \tag{1}$$

$$\frac{\partial\theta_2}{\partial t} = a_2^2\big(\frac{\partial^2\theta_2}{\partial r^2} + \frac{2}{r}\frac{\partial\theta_2}{\partial r}\big), \qquad \beta(t) < r < \infty, \tag{2}$$

with initial condition

$$\theta_1(r,0) = 0, \tag{3}$$

$$\theta_2(r,0) = f(r), \tag{4}$$

subjected to free boundary conditions

$$\theta_1(\alpha(t),t) = \theta_m, \tag{5}$$

$$-\lambda_1\frac{\partial\theta_1\big(\alpha(t),t\big)}{\partial r} = Q, \tag{6}$$

where $Q = \frac{P(t)}{2\pi\alpha^2(t)}$

$$\theta_1(\beta(t),t) = \theta_2(\beta(t),t) = \theta_m, \tag{7}$$

the Stefan's condition

$$-\lambda_1\frac{\partial\theta_1(\beta(t),t)}{\partial r} = -\lambda_2\frac{\partial\theta_2(\beta(t),t)}{\partial r} + L\gamma\frac{d\beta}{dt}, \tag{8}$$

as well as the condition at infinity

$$\theta_2(\infty,0) = 0. \tag{9}$$

To solve this problem we firstly reduce it to the linear problem. Then suggest exact solution in the form of Heat polynomials and series combination of integral error functions whose coefficients have to be determined.

**Keywords:** Stefan problem, degenerate domain, Integral Error functions.
**Mathematics subject classification:** 80A22

**References:**

[1] Sarsengeldin M.M., Kharin S.N., *Method of the Integral Error Functions for the Solution of the One- and Two-phase Stefan Problems and Its Application*, Filomat (accepted).

## Mathematical modeling of the temperature dependence of molten viscosity of metal according to the concept of chaotized particles

Aigul Sh. KAZHIKENOVA, D.B.ALIBIYEV, E.S. IBRAYEVA

[1] *Ye.A. Buketov Karaganda State University, Karaganda, Kazakhstan*
*E-mail: aigul-kazhikenova@mail.ru*

**Abstract:** Employees of the Chemical and Metallurgical Institute J.Abisheva proposed a new approach based on the Boltzmann distribution [1]. According to this approach, all three aggregate states can be viewed from the perspective of subordination Boltzmann distribution and to connect virtually every state with important characteristics of matter on the basis of excess or excess energy barriers melting and boiling. As in all cases considered particles, differing only in the amount of energy of chaotic motion, then their union and differential display can be described as the concept of chaotized particles. According to the concept of chaotized particles, crystal sliding and steam sliding, liquid sliding particles are present in each of the states of aggregation of matter.

The temperature dependence of viscosity fusion is defined on the basis of the concept of randomized particles in the full range of the liquid state on the uniform model considering degree of association of the elementary clusters of dynamically existing crystal mobility particles. The received form of temperature dependence of viscosity can be used for calculation of energy of activation of a viscous current fusion in a combination with Frenkel's equation. A new semi-empirical model of viscosity tested on 28 common metals for which there are reference data on the temperature dependence of viscosity. Obtained high values of correlation coefficients for the proposed model points to its functional characteristics. On this basis there were recommended calculated dependence for each metal.

**Keywords:** viscosity, chaotic particles, degree of associative clusters, the temperature dependences of the viscosity, liquid metals, fixed point

**2010 Mathematics Subject Classification:** 97M10

**References:**

[1] Malyshev V.P. , Turdukozhaeva A.M., Kazhikenova A.Sh., "The viscosity of the molten metal on the concept of randomized particles", *Heavy Engineering*, No.6, pp. 37-39, 2009.

# The mathematical models of electromagnetic field dynamics and heat transfer in closed electrical contacts including Thomson effect

Stanislav N.KHARIN [1,2], Merey M.SARSENGELDIN [1,2,3], Samat KASSABEK [1,2,3]

[1] *Kazakh-British Technical University, Almaty, Kazakhstan*
[2] *Institute of Mathematics and Mathematical Modelling, Almaty, Kazakhstan*
[3] *Suleyman Demirel University, Almaty, Kazakhstan*

**Abstract:**We represent mathematical models of electromagnetic field dynamics and heat transfer in closed symmetric and asymmetric electrical contacts including Thomson effect, which are essentially nonlinear due to the dependence of thermal and electrical conductivities on temperature [1]. Suggested solutions are based on the assumption of identity of equipotential and isothermal surfaces, which agree with experimental data [2] and valid for both linear and nonlinear cases. Well known Kohlrausch temperature-potential relation is analytically justified and agrees with [3-4].

**Keywords:** Thomson effect, symmetric and asymmetric electrical contacts

**Mathematics subject classification:** 74Nxx

**References:**

[1] Kharin S.N., Nouri H., Davies T., *The Mathematical Models of Welding Dynamics in Closed and Switching Electrical Contacts*, Proc. 49th IEEE Holm Conf. on Electrical Contacts, Washington, USA, pp. 107-123, 2003.

[2] Holm R., *Electric Contacts*, Springer Verlag, Berlin, 1981

[3] Diesselhorst H., *Ãœber das Problem eines elektrisch erwarmten Leiters*, Ann. Physics,Lpz., Vol.1, No.312, 1900

[4] Greenwood J., and Williamson J.B.P., *Electrical conduction in solids. II. Theory of temperature- dependent conductors*, Proc. Of the Royal Society (London), A246, pp.13-31, 1958.

## Solvability of a stationary problem of magnetohydrodynamics

Khonatbek KHOMPYSH [1], Sharypkhan SAKHAEV [1]

[1] *Al-Farabi Kazakh National University, Almaty, Kazakhstan*
*E-mail: konat_k@mail.ru, sakhaev_sh@mail.ru*

**Abstract:** This work is devoted to study the following stationary problem of magnetohydrodynamics consisting in finding the functions $\vec{v}(x)$, $p(x)$, $\vec{H}(x)$ and $\vec{E}(x)$ in $\Omega \subset R^3$:

$$-\nu\Delta\vec{v} + (\vec{v}\cdot\nabla)\vec{v} - \frac{\mu}{\rho}(\vec{H}\cdot\nabla)\vec{H} + \frac{1}{\rho}\nabla(p(x) + \frac{\mu}{2}|\vec{H}|^2) = \vec{f}(x), \quad x \in \Omega, \tag{1}$$

$$div\vec{v}(x) = 0, \quad x \in \Omega, \tag{2}$$

$$rot\vec{H}(x) - \sigma(\vec{E}(x) + \mu[\vec{v}\times\vec{H}]) = \vec{j}(x), \quad x \in \Omega, \tag{3}$$

$$div\mu\vec{H}(x) = 0, \quad x \in \Omega, \tag{4}$$

$$rot\vec{E}(x) = 0, \quad x \in \Omega, \tag{5}$$

$$\vec{v}(x)|_S = 0, \vec{E}_\tau(x)|_S = 0, \vec{H}\cdot\vec{n}|_S = 0. \tag{6}$$

Here $\vec{n}$ is the unit outward normal to $S$, and $\vec{E}_\tau = \vec{E} - \vec{n}\left(\vec{n}\cdot\vec{E}\right)$. $\Omega \subset R^3$ is the bounded domain with smooth boundary $S$.

Using the results in [1]- [3], we prove unique solvability of (1)-(6) in Sobolev and Hölder spaces.

**Keywords:** magnetohydrodynamics, generalized solution, stationary problem

**2010 Mathematics Subject Classification:** 76W99, 35D30, 60G10

**References:**

[1] Sakhaev Sh.S., Solonnikov V.A., "On Some Stationary Problems of Magnetohydrodynamics in Multy-Connected Domains", *Zap. Nauch. Sem. POMI*, Vol.397, pp. 126-149, 2011.

[2] Sakhaev Sh.S., Khompysh Kh., "On estimates of solutions of the linear stationary problem of magnetohydrodynamics problem in Sobolev spaces", *Advancements in Mathematical Sciences*, AIP Conference Proceedings 1676, pp. 020030-1-0200305, 2015.

[3] Solonnikov, V.A., "Some Stationary Boundary-Value Problems of Magnetohydrodynamics", *Trudy Math. Inst. Steklov*, Vol.59, pp. 5-36, 1960.

# Mathematical modelling and control theory analysis of artificial satellite

Sonu LAMBDA
[1] *National Institute of Technology Jamshedpur, India*
*E-mail: lambasonu1996@gmail.com*

**Abstract:** In this paper, an analysis of Mathematical Control Theory is carried out which concludes controllability of artificial satellites. Control Theory is application-oriented mathematics that deals with the basic principles underlying the analysis and design (control) systems. To control means that one has to influence the behaviour of the system in a desirable way. A preliminary discussion is provided which acts as stairs for reaching final conclusions and includes an analytical approach to model the physical system into dynamical equations of motions which is termed as Mathematical Modelling. Since most of the real life systems are nonlinear in nature therefore an approach for linearization is described. By doing so, we will able to apply numerous linear analysis methods to study further behaviour of the system. Then a mathematical approach used to compute Transition Matrix for getting solution of linear system by using variation of parameter method. Throughout the analysis we are looking for minimum norm (optimal) steering controls,we will also introduce linear oprator technique for calculating the steering controls of the system, after then we made some interesting conclusions by using Kalman's Controllability test. Finally, a graphical simulation is provided using MATLAB which includes computations of controllability Grammian and controller with related MATLAB codes.

**Keywords:** Control Theory, Transition Matrix, Optimal Control, Kalman's Controllability test.

# New effects for high speed collision of blunt bodies with atmospheric inhomogeneities

Vladimir LEVIN [1], Pavel GEORGIEVSKIY [2]

*Lomonosov Moscow State University, Russia*
*E-mail:* [1] *levin@imec.msu.ru,* [2] *georgi@imec.msu.ru*

**Abstract:** The results of numerical investigation of collision of blunt bodies moving in the atmosphere with supersonic speeds with local inhomogeneities – low density and high density gas bubbles are presented. The effect of appearing of anomalous pressure and density jumps in a critical point of a body essentially exceeding values calculated for 1D Riemann problem for the interaction of a bow shock wave with plain layer is studied. In case of the interaction of blunt bodies with low density bubbles the effect was previously observed but was not enough clearly explained in [1]. Similar results were obtained both experimentally and theoretically for the interaction of blunt bodies with high temperature plasma formations induced by pulsed laser energy deposition in supersonic upstream flow.

Now it is shown that the main reason for appearing of anomalous pressure and density jumps in a critical point of a body is the preceding fast process of focusing of transversal shock wave in small zone on the symmetry axes ahead of a body which is accompanied in some cases by formation of thin axial high speed cumulative jets. The mechanism of focusing processes was in general agreement with well investigated shock focusing scenarios for the interaction of plane shock with low density and high density gas bubbles [2] providing of course taking into account some specific of this particular case due to the curvature of a bow shock wave and the existence of a body closely near the focusing region.

For low density gas bubbles according to the "divergent" scenario of the interaction the "weak focusing" regime was realized. The secondary weak convergent-type shock wave propagating inside a shock layer was focused along the symmetry axes. Shortly after that the pressure and density jumps in a critical point of a body were induced by a retransmitted shock wave reflected from the rear interface of a gas bubble which was amplified by defocusing shock wave propagating from the shock focusing region.

For high density gas bubbles the "strong focusing" regime was realized. According to "convergent" scenario of the interaction the bow shock wave was diffracted around the bubble. Then the strong converging diffracted bow shock wave and the transmitted shock wave were collapsed in a very small zone (almost a point) at the rear pole of the bubble on the symmetry axes. The collapsing process was accompanied by formation of thin supersonic upstream and downstream cumulative jets. The interaction of downstream jet with a body surface was resulted in dramatic pressure and density jumps.

Parametric investigations for ellipsoidal gas bubbles of different density ratio, size and elongation were carried out. It was concluded that to realize the cumulation effect the density ratio should be more than some critical value, the gas bubble shape should be closed to spherical and the longitudinal size – about

a bow shock wave distance. There were no focusing or cumulation effects for the interaction with a flat layer or ellipsoidal gas bubbles of too large cross size.

The investigations were carried out in 2016 in the Institute of Mechanics of Moscow State University (project AAAA-A16-116021110195-4) and were financially supported partially by the Council on Grants of the President of Russian Federation (project NSh-8425.2016.1), Russian Science Foundation (project 14-11-00773) and the Russian Foundation for Basic Research (projects 14-01-00891-a, 16-29-01092-ofi).

**Keywords:** supersonic flow, numerical simulation, blunt body, gas bubble, shock focusing, cumulation

**2010 Mathematics Subject Classification:** 76M20

**References:**

[1] Georgievskii P.Yu., Levin V.A. "Unsteady Interaction of a Sphere with Atmospheric Temperature Inhomo-geneity at Supersonic Speed", *Fluid Dynamics*, Vol.28, No.4, pp. 568-574, 1993.

[2] Georgievskiy P.Yu., Levin V.A., Sutyrin O.G. "Interaction of a Shock with Elliptical Gas Bubbles", *Shock Waves*, Vol.25, No.4, pp. 357-369, 2015.

# Initiation of 3D Detonation in Supersonic Flows in Channels of Special Geometry

Vladimir LEVIN [1], Ivan MANUYLOVICH [2], Vladimir MARKOV [3]

[1 2] *Lomonosov Moscow State University, Russia*
*E-mail:* [1] *levin@imec.msu.ru,* [2] *manuylovich@imec.msu.ru*
[3] *Steklov Mathematical Institute of Russian Academy of Sciences, Russia*
*E-mail: markov@mi.ras.ru*

**Abstract:** The problem of the formation of detonation in 3D channel with square variable cross-section have been investigated. Detailed stationary and unsteady flow patterns with and without detonation have been found. The critical values of gasdynamic parameters separating different flow regimes were obtained. Hypersonic analogy of flat and 3D flows, allowing the use of 2D solutions to estimate the parameters of three-dimensional supersonic flows, have been confirmed. The limitations of the analogy concerned with chocking effect of the burning flow have been determined. A numerical study of detonation of the combustible mixture in 3D cylindrical channels with helical obstacles of various shapes have been conducted: 1) with a twisted lateral surface of elliptic cross-section; 2) with a screw body of elliptic cross-section, located at the axis of the channel; 3) with the stellate body composed of a number of twisted thin blades located at the axis of the channel; 4) with a number of twisted thin blades attached to the lateral surface of the channel, inducing an analogy with a rifled barrel in the artillery. All of these forms of static 3D channels with the obstacles can be obtained from the 2D rotating regions through their simultaneous movement along the axis of the channel and rotation. According to the hypersonic analogy, the distribution of parameters, which arise in the solution of plane problems with rotation, should be close to the distributions of the parameters in cross-sections of the 3D channel with helical elements upon injection of supersonic flow of combustible mixture in the channel. As a result of numerical calculations of unsteady flows in the specified channels, detailed stationary flow patterns with detonation waves near the obstacles have been obtained. These waves had complicated structure and interacted with each other and with the channel walls. The comparison with 2D calculation was made and the above-mentioned analogy was confirmed. The study is carried out in a framework of one-step combustion kinetics by numerical method based on the S.K. Godunov scheme. The original parallelized software package with graphical interface was developed and used to perform numerical investigation. The calculations have been performed on the MSU "Lomonosov" supercomputer.

This research has been supported by the Russian Foundation for Basic Research (Grant No. 14-01-00742-a, No. 16-29-01092 ofi_m), by the President Grant for Support of Leading Science Schools of Russian Federation (No. NSh-8425.2016.1), by the Presidium of the Russian Academy of Sciences (special program) and by the Supercomputing Center of Lomonosov Moscow State University.

**Keywords:** detonation of the combustible mixture, combustion kinetics

**2010 Mathematics Subject Classification:** 76M20

## Supersonic multi-species flow with particle dispersion on a mixing layer

Altyn MAKASHEVA [1], Altynshash NAIMANOVA [2]

*Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan*
[1] *E-mail: altyn-mak@mail.ru*
[2] *E-mail: alt_naimanova@yahoo.com*

**Abstract:** Particle-laden flows play an important role in high-speed technologies such as solid rocket propulsion systems and high-speed fuel combustors. The flow physics in such devices is very complex due to shock dynamics, turbulence and particle dispersion in mixing layers. Direct numerical simulation (DNS) has been successfully used for compressibility effects in the turbulent shear layer [1], single-phase and two-phase flows [2]. The particle dispersions in the mixing layers for the different Stokes numbers have been obtained in [3]. In spite of that there are a few investigations of a compressible multispecies shear layer flow with the dispersion of particles.

This work is our first stage of the simulation in a comprehensive study of the flow-particle interactions in a three-dimensional multispecies turbulent medium. The direct numerical simulations of the flowfield structures and the properties of the particle dispersion in the quasi-2D turbulent mixing layer (hydrogen-air) are performed by solving the time-dependent, compressible Euler equations. The 3D numerical code using the high-order essentially non-oscillatory (ENO) scheme is developed. The dispersion of the particles is studied by following their trajectories in the mixing layer with the Lagrangian method. In detail, the effect of the initial mass fraction of hydrogen and the number of particles on the growth of vortices and their thickness is studied. The simulation reveals that the capturing of the particles by the vortices essentially depends on the density of particles.

**Keywords:** supersonic shear flow, mixing layer, particle dispersion, multi-species flow, ENO-scheme.

**2010 Mathematics Subject Classification:** 02.60.Cb, 02.70.Bf

**References:**

[1] Pantano C., and Sarkar S., "A study of compressibility effects in the high-speed turbulent shear layer using direct simulation", *Fluid Mechanics*, Vol.451, pp. 329-374, 2002.

[2] Moin P., and Mahesh K., "Direct numerical simulation: a tool in turbulence research", *Fluid Mechanics*, Vol.30, pp. 539-578, 1998.

[3] Zhiwei Hu., Xiaoyu Luo, and Kai H., Luo, "Numerical simulation of particle dispersion in a spatially developing mixing layer", *Theoret. Comput. Fluid Dynamics*, Vol.15, pp. 403-420, 2002.

## Modeling and solving an optimization problem of supply of chemical raw materials

Ernazar NYSSANOV[1], Marina SAPRYGINA[2]

[1] *M. Auezov South Kazakhstan State University, Kazakhstan*
*E-mail: nyssanb@mail.ru*
[2] *South Kazakhstan State Pharmaceutical Academy, Shymkent, Kazakhstan*
*E-mail: marina-iv@mail.ru*

**Abstract:** In this article the optimization problem of supply of the chemical raw materials is solved in the following formulation. The trading company is specialized in the supply of the chemical raw materials to the industrial enterprises. The delivery is carried out by self - the transport of buyers. It is given the nomenclature data set of the chemical raw materials, the cost of its purchase, delivery and warehousing. The profit from the sale of each type of the chemical raw materials is fixed. The financial resources for the planning period are limited. In addition, on the basis of existing and potential customer's orders for the planning period, the limits on the number of purchased raw materials of every kind were defined.How many of each type of the chemical raw materials is necessary to buy for the company to get the best profit under the condition that all the purchased raw materials will be sold in the planning period? The problem is solved by the method of the linear programming. At the same time a mathematical formulation of the problem is carried out. This means that the controlled variables and the target function of the problem are identified. Then, by means of these variables the objective function and constraints are described in the form of linear relations. [1-3].

**Keywords:** Quantitative methods in management, linear programming, numerical optimization, objective function, simplex method, Mathcad, Qbasic

**2010 Mathematics Subject Classification:** 65K05, 90C05, 90B50

**References:**

[1] Zaitsev M.G., Varyukhin S.E., *Methods to optimize the management and decision-making: examples, problems, cases. Tutorial.* M., "Business", 664 p., 2008.

[2] Chase R.B., Ekvilayn N.J., Jacobs R.F., *Production and Operations Management.* Moscow-St Petersburg-Kiev: "Williams", 532 p., 2001.

[3] Eddous M., Stenfild R., *Methods of decision-making.* M., "UNITY Audit", 426 p., 1997.

## Fragmenting of algorithm for solving the problem of clustering miRNA families

Anna PYRKOVA [1], Anatoly IVASHCHENKO [2], Raigul NIYAZOVA [2]

[1] *Faculty of Mechanics and Mathematics, , Al-Farabi Kazakh National University,*

[2] *Faculty of Biology, Al-Farabi Kazakh National University, Almaty, Kazakhstan*

*E-mail: Anna.Pyrkova@kaznu.kz, a_ivashchenko@mail.ru, raiguln@mail.ru*

**Abstract:** The division of miRNAs into families does not adequately reflect the degree of nucleotide sequence similarity, and thus the categorisation of miRNAs into families requires quantitative criteria defining the differences between miRNA families. The genomes of different organisms have orthologous miRNAs that should be distributed into families. Than it is necessary to establish the degree of similarity of orthologous miRNAs and their belonging to different families. The present research is aimed at solving the problem of clustering miRNAs based on their nucleotide sequence similarity using the fragmented programming method.

Using this developed method, the miRNA nucleotide sequences of Arabidopsis thaliana were clustered. The results identified a conserved polynucleotide, which serves as the Ath-miR156a-j binding site in paralogous SPL mRNAs. These nucleotides encode the conserved ALSLLS motif and the miR156 and miR157 subfamilies belong to the same family. The human miR-1273 family includes miRNAs with very different nucleotide sequences; therefore, based on our program they belong to different miRNAs families.

**Keywords:** miRNA family, algorithm, fragmented programming method.

**References:**

[1] Eric Londina et al., Proc Natl Acad Sci USA. 2015 112: E1106 [PMID: 25713380]

[2] Wan L, Ding J, Jin T, Guan J, Zhou S., Automatically clustering large-scale miRNA sequences: methods and experiments. BMC Genomics. 2012;13 Suppl 8:S15. doi: 10.1186/1471-2164-13-S8-S15. Epub 2012 Dec 17. PubMed PMID: 23282099; PubMed Central PMCID: PMC3535721.

[3] Kaczkowski B., Torarinsson E., Reiche K., Havgaard JH., Stadler PF., Gorodkin J., Structural profiles of human miRNA families from pairwise clustering. Bioinformatics. 2009 Feb 1;25(3):291-4. doi: 10.1093/bioinformatics/btn628. Epub 2008 Dec 4. PubMed PMID: 19059941.

## Iterative method of finding hydraulic conductivity characteristics of soil moisture

Bolatbek RYSBAIULY [1], Abilmazhin ADAMOV [2]

[1] *Department of Mathematical and Computer Modeling, International Information Technologies University, Almaty, Kazakhstan*
*E-mail: b.rysbaiuly@mail.ru*
[2] *L. N. Gumilyov Eurasian National University, Kazakhstan*
*E-mail: adam1955@mail.ru*

**Abstract:** The work considers the initial boundary value problem for a nonlinear equation of hydraulic conductivity. A method of finding a nonlinear diffusion coefficient was developed and hydraulic conductivity of soil moisture was found. Numerical calculations were conducted.

**Keywords:** moisture conduction, inverse problem, conjugate problem, difference scheme, diffusion coefficient

**2010 Mathematics Subject Classification:** 80A20, 35K55, 65M06, 65M32

**References:**

[1] Childs E.C., Collis-George N., *Proceedings the Royal Society A*, Vol.201, pp. 392-405, 1950.
[2] Nerpin S.I.,Chudnovsky A.F., *The soil physics*, M.:Nauka, p.584, 1967.
[3] Lykov A.V., *Heat and mass transfer*, M.: Energiya, p. 560, 1982.
[4] Jackson R.D., Regina R.S., e. a., *Soil Science Society of America*, Vol.38, pp. 861-866, 1984.
[5] Reynolds A.J., *Turbulent flows in Engineering*, Bruner University, London, p. 408, 1974.
[6] Globus F.M., *Experimental hydro physics of the soil* , L.: Gydrometeoizdat, p. 355, 1969.
[7] Rysbaiuly B., Baymankulov A.T., *DAN NASc RK*, Vol.3, pp. 5-8, 2009.
[8] Rysbaiuly B., Baymankulov A.T., *International Journal of Academic Research*, Vol.5, pp. 84-91, 2010.
[9] Rysbaiuly B., *Wulfenia Journal, Austria*, Vol.20, 12, pp. 311-335, 2014.
[10] Gardner W.R., *Highway Research board, Rep*, 40, pp. 78-87, 1958.
[11] Fedyaeva E.A., "Mathematical modeling of thermal and moisture transfer in dispersed unsaturated soils", in *Lomonosov Readings, 16-22 April of 2014, Moscow State University, section Engineering and environmental geology.*, M., 2014.
[12] Gray W.G., Miller C.T., *Advances in Water Resources*, Vol.28, pp. 161-180, 2005.
[13] Rohde A.A., *Fundamentals of soil moisture*, L.:Hydrometeoizdat. Vol.1, 1985.
[14] Blackwell B., Charles St Clair Jr., *Ill-posed inverse problems of the heat conductivity.*, M.: Mir, p. 312, 1989.
[15] Kabanihin S.I., *Inverse and ill-Posed Problems. Theory and Applications.* Degruter, Germany, p. 459, 2011.
[16] Tikhonov A.N., *Reports of ASc USSR*, Vol.39, 5, pp. 195-198, 1943.
[17] Rysbaiuly B., *Calculation of thermo physical characteristics of the soil.* in International conference "Advanced Mathematics, Computations & Applications-2014", Akademgorodok, Novosibirsk, Russia, June 8-11 2014, 60.
[18] Samarsky A.A., *Theory of difference schemes.* M.: Nauka, p. 616, 1983.

# Spectral properties of a Laplace operator with Samarskii–Ionkin type boundary conditions in a disk

Makhmud SADYBEKOV [1], Abdizhahan SARSENBI [1,2],
Aizhan TENGAYEVA [1,3]

[1] *Institute of Mathematics and Mathematical Modelling, Kazakhstan*
*E-mail: sadybekov@math.kz, sarsenbi@math.kz*
[2] *Auezov South Kazakhstan State University, Kazakhstan*
[3] *Kazakh National Agrarian University, Kazakhstan*
*E-mail: aijan0973@mail.ru*

**Abstract:** In this paper we consider a spectral problem for a model differential operator of the second order with involution. The operator is given by a differential expression

$$Lu = -u''(-x), -1 < x < 1 \tag{1}$$

and boundary conditions of the general form:

$$\alpha_j u'(-1) + \beta_j u'(1) + \alpha_{j1} u(-1) + \beta_{j1} u(1) = 0, \;\; j = 1, 2. \tag{2}$$

Coefficients of the boundary condition are numbers, which can be complex.

A criterion of the basis property of systems of eigenfunctions of the operator is set in the sense of coefficients of the boundary conditions. The interest to studying the basis properties of operators of the form (1), (2) has arose relatively recently. In works of T.Sh. Kal'menov and his disciples the *self-adjointness* property of the Cauchy operator:

$$L_K u = -u''(-x), \;\; -1 < x < 1, \quad u'(-1) = 0, \;\; u(-1) = 0 \tag{3}$$

is successfully applied in solving initial-boundary value problems for a heat equation with inverse time direction, in finding a criterion of well-posedness of the Cauchy problem for the Poisson operator, in formulating an asymptotic solution to problems with a small parameter.

**Keywords:** generalized spectral problem, involution, eigenfunctions, eigenvalues, basis property

**2010 Mathematics Subject Classification:** 34L10, 34L15

## Spectral properties of a Laplace operator with Samarskii–Ionkin type boundary conditions in a disk

Makhmud SADYBEKOV [1], Nurgissa YESSIRKEGENOV [1,2]

[1] *Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan*
[2] *Department of Mathematics, Imperial College London, London, United Kingdom*
*E-mail: sadybekov@math.kz, n.yessirkegenov15@imperial.ac.uk*

**Abstract:** One of the most important problems in the theory of harmonic functions are Dirichlet and Neumann boundary value problems. In the one-dimensional case, or when considering the problem in a multidimensional parallelepiped, the main problems include also the periodic boundary value problems.

In this paper we consider spectral properties of a Laplace operator with boundary conditions of Samarskii-Ionkin type in a disk and prove the completeness of eigenfunctions.

Let $\Omega = \{z = (x, y) = x + iy \in C : |z| < 1\}$ be a unit disk, $r = |z|, \varphi = \arctan(y/x)$, $\Omega^+ = \Omega \cap \{y > 0\}$, and $\Omega^- = \Omega \cap \{y < 0\}$. We consider the spectral problem corresponding to the Laplace operator

$$-\Delta u(z) = \lambda u(z), |z| < 1$$

with local boundary conditions

$$u(1, \varphi) = 0, \ \ 0 \le \varphi \le \pi,$$

and with one of the nonlocal boundary conditions

$$\frac{\partial u}{\partial r}(1, \varphi) - \frac{\partial u}{\partial r}(1, 2\pi - \varphi) = 0, \ \ 0 \le \varphi \le \pi,$$

or

$$\frac{\partial u}{\partial r}(1, \varphi) + \frac{\partial u}{\partial r}(1, 2\pi - \varphi) = 0, \ \ 0 \le \varphi \le \pi.$$

We note that unlike the one-dimensional case the system of root functions of the problems consists only of eigenfunctions.

**Keywords:** Poisson equation, Samarskii–Ionkin type problem, eigenfunctions, eigenvalues

**2010 Mathematics Subject Classification:** 35J05, 35J25, 35P10

## Analytical solution of two-phase spherical Stefan problem by heat polynomials and integral error functions

Merey M.SARSENGELDIN [1,2,3], Stanislav N.KHARIN [1,2], Hassan NOURI [4]

[1] *Kazakh-British Technical University, Almaty, Kazakhstan*
[2] *Institute of Mathematics and Mathematical Modelling, Almaty, Kazakhstan*
[3] *Suleyman Demirel University, Almaty, Kazakhstan*
[4] *University of the West of England (UWE) Bristol, UK*

**Abstract:** On the base of the Holm model [1], we represent two phase spherical Stefan problem and its analytical solution, which can serve as a mathematical model for diverse thermo-physical phenomena in electrical contacts [2]. Spherical Stefan problem is reduced to linear problem with degenerate domain. Suggested solution is obtained from Integral Error Function and its properties which are represented in the form of series whose coefficients have to be determined:

$$u_1(x,t) = \sum_{n=0}^{\infty} A_{2n}(2a_1\sqrt{t})^{2n}\left[i^{2n}erfc\frac{-x}{2a_1\sqrt{t}} + i^{2n}erfc\frac{x}{2a_1\sqrt{t}}\right]$$
$$+\sum_{n=0}^{\infty} A_{2n+1}(2a_1\sqrt{t})^{2n+1}\left[i^{2n+1}erfc\frac{-x}{2a_1\sqrt{t}} - i^{2n+1}erfc\frac{x}{2a_1\sqrt{t}}\right], \qquad (1)$$

$$u_2(x,t) = \sum_{n=0}^{\infty}(2a_2\sqrt{t})^{n}\left[B_n i^{n}erfc\frac{-x}{2a_2\sqrt{t}} + C_n i^{n}erfc\frac{x}{2a_2\sqrt{t}}\right]. \qquad (2)$$

We find free interphase boundary, unknown coefficients $A_{2n+1}, A_{2n}, B_n$ and $C_n$ from boundary conditions utilizing Faa Di Bruno formula, Bell polynomials and Lebniz differentiation formula. To simplify calculations, we use Hermite polynomials to represent series (1) in the following form:

$$u_1(x,t) = \sum_{n=0}^{\infty} A_{2n} \sum_{m=0}^{n} x^{2n-2m}t^m B_{2n,m} + \sum_{n=0}^{\infty} A_{2n+1} \sum_{m=0}^{n} x^{2n-2m+1}t^m B_{2n+1,m}.$$

Convergence of solution series is proved.

**Keywords:** Stefan problem, integral error function, heat polynomials

**Mathematics subject classification:** 80A22

**References:**

[1] Holm R., *Electric Contacts*, Fourth Edition, Springer Verlag , 482 p., 1981.
[2] Sarsengeldin M.M., Kharin S.N., *Method of the Integral Error Functions for the Solution of the One- and Two-phase Stefan Problems and Its Application*, Filomat (accepted)

## The mathematical modeling of grouping the dipole water clusters

V. SHAIDUROV [1], Viktoriya KORNIENKO [2], A. VYATKIN [1]

[1] *Institute of Computational Modeling of Russian Academy of Sciences, Krasnoyarsk, Russia,*
*E-mail: shaidurov04@mail.ru*
[2] *Siberian Federal University, Krasnoyarsk, Russia*
*E-mail: vika-svetlakova@yandex.ru*

**Abstract:** In the paper, a physical-mathematical model and a computational algorithm implementing the model are proposed to study the behavior of particles having an electric dipole moment in an external electric field. Computational experiments demonstrate the orientation dynamics of water clusters with the increase of the generated field. The dipole properties of some water clusters were previously determined using Hyperchem program.

**Keywords:** electric dipole, water clusters, physical-mathematical model, computational experiment, Runge-Kutta methods

**2010 Mathematics Subject Classification:** 93A30, 70-08

**References:**

[1] Miyake T., Aida M., H-bond patterns and structure distributions of water octamer $(H_2O)_8$ at finite temperatures. *Chemical Physics Letters*. Vol.424, pp. 215-220, 2006.

[2] Feynman R.P., Leighton R.B., Sands M., The Feynman Lectures on Physics. Vol. 2. London, Addison-Wesley Publishing Company, 1964.

[3] Kittel Ch., Introduction to Solid State Physics. Wiley, 2004.

[4] Vyatkin A., Analyzing the Behavior of the Explicit Runge-Kutta Methods at the Singular point in the Two-body Problem. *Automation and Remote Control*. Vol.74(2), pp. 325-330, 2013.

## Simulation of shock wave boundary layer interaction in flat channel with jet injection

Nurtoleu SHAKHAN[1], Asel BEKETAEVA[2], Altynshash NAIMANOVA[2]

[1] *Al-Farabi Kazakh National University, Almaty, Kazakhstan*
*E-mail: shakhan-nurtoleu@mail.ru*
[2] *Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan*
*E-mail: azimaras@mail.ru*

**Abstract:** The flow around jets has been comprehensively investigated experimentally [1] and simulated numerically. Nowadays the interactions between boundary layer and shock wave in the channel flow field with perpendicular jet injection aren't well understood. Due to complexity of these interactions, usually they are studied by parts. One part of these studies focuses on the properties of the shock wave structures in the region of the jet injection [2], other considers the zone of the shock wave boundary layer interaction [3]. In the present we simulate and study the shock wave structures in the supersonic channel flow with jet injection where both processes will be present. For describing the process of the multispecies supersonic gas flow in the flat channel with perpendicular jet injection, Favre-averaged Navier-Stokes equations coupled with $k-\omega$ turbulence model are applying. High order Weighted Essentially Non-Oscillatory (WENO) scheme is used to approximate convective terms. During the investigation of flow physics in detail, the three shock wave structures are observed: in the region of the jet (barrel, bow, oblique and closing shocks), on the upper boundary layer (reflection, transmitted and reattachment shocks), and new structures behind the jet on the lower boundary layer, which are analogous to the structures on the upper boundary layer.

**Keywords:** supersonic flow, SWBLI, flow control, separation, Navier-Stokes equations, WENO-scheme, shock wave

**2010 Mathematics Subject Classification:** 02.60.Cb, 02.70.Bf

**References:**

[1] Glagolev A.I., and Zubkov A.I., "Interaction of a gas jet exhausting from an orifice in a flat plate with a supersonic flow", *Izv. Akad. Nauk SSSR, Mekh. Zhidk. Gaza*, pp. 99-102, 1978.

[2] Viti V., and Wallis S., "Jet interaction with a primary jet and an array of smaller jets", *AIAA Journal*, Vol.42, pp. 1358-1368, 2004.

[3] Pasquariello V., and Grilli M., "Large-eddy simulation of passive shock-wave/boundary-layer interaction control", *International Journal of Heat and Fluid Flow*, Vol.49, pp. 116-127, 2014.

## Application of of fuzzy logic for matching modes use of wind power plant with an electrical load schedule

R.Y. SHIKHLINSKAYA [1], S.M. SALIMOV [2]

[1] *Institute of Applied Mathematics, Baku State University, Baku, Azerbaijan*
*E-mail: reyhanshikhli@gmail.com*
[2] *Institute of Economics of ANAS, Baku, Azerbaijan*

**Abstract:** This article proposed method, based on the theory of fuzzy sets, taking into account the uncertainty of wind power, as an energy carrier, it is a random uncontrollable natural process, for for matching modes use of wind power plant with an electrical load schedule. - *in Absheron, Typical for Caspian coast.* The average annual of wind speed of this zone is more than $4m/sec$.

Wind power unit runs on unmanaged schedule, in connection with the many uncertainties and unpredictable parameters. The consumer often does not allow for interruptions in the supply of energy.

Bases of the analysis mode of operation the wind turbine in fuzzy-defined conditions based on the principle of constructing a fuzzy model based on binary and conventional fuzzy relationship. The first of these fuzzy relations based on two basic sets X and Y, and the second – on two basic sets $Y$ and $Z$. Here $X$ – it describes a number of characteristics of wind power, taken on the Beaufort scale, $Y$ - the set of characteristics of power output, and $Z$ - set of daily intervals load. Fuzzy relation L - describes development of capacity at a certain wind turbine, and $N$ - a covering daily load. For the construction of fuzzy relation N of a covering daily load function used concept of a conditional fuzzy subset:

$$\mu(y_i, z_j) = max(min(\mu(y_i, z_j), \mu(y_i))). \tag{1}$$

Using the method of construction of fuzzy relations, provided possibility of comparing produced cabacity to the load schedule, based on which it can be concluded about the use of electric energy.

**Keywords:** alternative energy resources, wind power plant, solar baterries, wind power, wind speed, wind energy, wind linguistic change, construction method of uncertain relation, the theory of fuzzy sets.

**2010 Mathematics Subject Classification:** 94D05, 00A71

# Numerical solution of $2D$-vector tomography problem using the method of approximate inverse

Ivan SVETOV[1], Svetlana MALTSEVA[2], Anna POLYAKOVA[3]

[1,2,3] *Sobolev Institute of Mathematics, Russia*
[1,2,3] *Novosibirsk State University, Russia*
*E-mail:* [1] *svetovie@math.nsc.ru,* [2] *sv_maltseva@mail.ru,*
[3] *apolyakova@math.nsc.ru*

**Abstract:** The problem of vector tomography is considered in this work in the following formulation. Let a certain vector field $\mathbf{v}$ be given in a bounded domain of $\mathbb{R}^2$ filled by a medium in which the probing radiation propagates along straight lines. We have to reconstruct this field from its known longitudinal and (or) transverse ray transforms.

We propose an algorithm for solution of the vector tomography problem. The algorithm based on so called method of approximate inverse developed by A.K. Louis and his pupils [1–3]. The idea of the method of approximate inverse is as follows. Let $A : H \to K$ be a linear bounded operator. It is required to find an approximate solution (a function $f$) of the operator equation $Af = g$ for given $g \in K$. Mollifiers $e_\gamma^y$ are used for solving. This functions have a properties $\langle e_\gamma^y, e_\gamma^y \rangle_H = 1$ and $\langle f, e_\gamma^y \rangle_H \approx f(y)$. Let $A^*$ be an adjoint operator for $A$. Hence, equation $A^* \psi_\gamma^y = e_\gamma^y$ has a solution $\psi_\gamma^y \in K$ and

$$f(y) \approx \langle f, e_\gamma^y \rangle_H = \langle f, A^* \psi_\gamma^y \rangle_H = \langle Af, \psi_\gamma^y \rangle_K = \langle g, \psi_\gamma^y \rangle_K.$$

The authors were supported partially by the Russian Foundation for Basic Research (project 14-01-31491-mol_a)

**Keywords:** approximate inverse, vector tomography, ray transform, solenoidal vector field, potential vector field

**2010 Mathematics Subject Classification:** 44A12, 65R10, 65R32, 94A08

**References:**

[1] Louis, A.K., Maass, P., "A mollifier method for linear operator equations of the first kind", *Inverse Problems*, Vol.6, pp. 427–440, 1990.

[2] Louis, A.K., "Approximate inverse for linear and some nonlinear problems", *Inverse Problems*, Vol.12, pp. 175–190, 1996.

[3] Rieder, A., Schuser, T., "The Approximate Inverse in Action with an Application to Computerized Tomography", *SIAM Journal on Numerical Analysis*, Vol.37, No.6, pp. 1909–1929, 2000.

# Duality property of the noncommutative $\ell_\infty$ and $\ell_1$ valued symmetric Hardy spaces

Kanat TULENOV [1,2]

[1] *Institute of Mathematics and Mathematical Modeling, Kazakhstan.*
*E-mail: tulenov@math.kz*
[2] *Al-Farabi Kazakh National University, Kazakhstan*
*E-mail: kanat.tulenov@gmail.com*

**Abstract:** In this paper we consider the noncommutative $H_E(\mathcal{A};\ell_\infty)$ and $H_E(\mathcal{A};\ell_1)$ spaces and obtain some result on duality for these spaces, i.e we obtain the following result:

**Theorem.** Let $E$ be an $r$-convex symmetric Banach function space on $[0;1]$ for some $0<r<\infty$ and $E$ do not contain $c_0$ or separable. Then

$$(i)\ \ (H_E(\mathcal{A};\ell_1))^*=L_{E^\times}(\mathcal{M};\ell_\infty)/J(H^0_{E^\times}(\mathcal{A};\ell_\infty))$$

isometrically via the following duality bracket

$$((x_n),(y_n))=\sum_{n=1}^{\infty}\tau(y_n^*x_n)$$

for $x\in H_E(\mathcal{A};\ell_1)$ and $y\in H_{E^\times}(\mathcal{A};\ell_\infty)$, where $J(H^0_{E^\times}(\mathcal{A};\ell_\infty))=\{x^*:\ x\in H^0_{E^\times}(\mathcal{A};\ell_\infty)\}$.

$$(ii)\ \ (L_E(\mathcal{M};\ell_1)/J(H^0_p(\mathcal{A};\ell_1)))^*=H_{E^\times}(\mathcal{A};\ell_\infty)$$

isometrically via the following duality bracket

$$((x_n),(y_n))=\sum_{n=1}^{\infty}\tau(y_n^*x_n)$$

for $x\in H_E(\mathcal{A};\ell_1)$ and $y\in H_{E^\times}(\mathcal{A};\ell_\infty)$, where $J(H^0_{E^\times}(\mathcal{A};\ell_1))=\{x^*:\ x\in H^0_{E^\times}(\mathcal{A};\ell_1)\}$.

This theorem is analogue of Theorem 5 in [1].

**Keywords:** von Neumann algebra, subdiagonal algebras, noncommutative vector valued symmetric Hardy spaces, duality.

**2010 Mathematics Subject Classification:** 46L51, 46L52.

**References:**

[1] T. N. Bekjan, K. Tulenov, and D. Dauitbek, The Noncommutative $H_p^{(r,s)}(\mathcal{A};l_\infty)$ and $H_p(\mathcal{A};l_1)$ spaces. *Positivity*, **19** 877–891 (2015).

# Lateral boundary conditions for the Klein-Gordon-Fock equation

Kanat TULENOV [1], Dostilek DAUITBEK [2]

[1] *Institute of Mathematics and Mathematical Modeling*
*E-mail: tulenov@math.kz*
[2] *Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: dauitbek@math.kz*

**Abstract:** In this paper we consider an initial-boundary value problem for the Klein-Gordon-Fock equation. In [**?**], Authors considered one dimentional potential

$$u(x,t)=\int_{\Omega}(-\frac{1}{2}|x-y|)f(y)dy \tag{1}$$

in $\Omega=(0,1)$ and the equation

$$-u^{''}(x)=f(x),\ \ x\in\Omega, \tag{2}$$

and they showed that if they are solved the equation (2) with the boundary conditions

$$u^{'}(0)+u^{'}(1)=0,\ \ -u^{'}(1)+u(0)+u(1)=0. \tag{3}$$

then they were finded an unique solution of that boundary value problem in the form (1).

This simple method finds equivalent boundary value problems of ODE for one dimensional potential integrals. However this task becomes tedious for PDE, and in works [1, 2] are obtained boundary conditions of the volume potentials for elliptic and hyperbolic equations and showed some their applications. In particular, in [2], by using a new non-local boundary value problem, which is equivalent to the Newton potential, Authors found explicitly all eigenvalues and eigenfunctions of the Newton potential in the 2-disk and the 3-ball. The aim of this work is to find lateral boundary conditions for the Klein-Gordon-Fock equation as shown in [**?**].

**Keywords:** Potential, fundamental solution, Klein-Gordon-Fock equation

**2010 Mathematics Subject Classification:** 35J05, 35J08, 35J25, 35P05

**References:**

[1] Kal'menov T.Sh., Suragan D., "Initial-boundary value problem for the wave equation", *Electron. J. Diff. Equ.*, Vol. 48, pp. 1-6, 2014.
[2] Kal'menov T.Sh., Suragan D., "To spectral problems for the volume potential", *Doklady Mathematics*, Vol. 80, No. 2, pp. 646-649, 2009.

## A modelling the dynamics of in-situ leaching process at the microscale

Ainur ZHUMALI [1]

[1] *Center of Mathematics and Cybernetics, Kazakh-British Technical University, Almaty, Kazakhstan*
*E-mail: ainura_.89@mail.ru*

**Abstract:** Some physical and chemical processes take place at interfaces between the solid and the liquid. One of them is in-situ leaching, an effective and promising method of extraction of valuable components from ores by their selective dissolution. To explore the main features of the movement of fluids media and dissolution mechanism of rocks by acids, it is considered a mathematical models at the microscopic level [1], using equations of continuum mechanics [2] and well-known chemical laws. The important point here is a derivation of new conditions at the free boundary and the dynamics of this boundary.

The efficient and accurate numerical simulation at the microscale plays an important role in understanding of the basic physical properties of rocks leaching processes. The application of complex numerical algorithms on a huge grids leads to the high demands on computer resources. Therefore, the numerical simulation of exact mathematical models is based on parallel computing.

**Keywords:** solid-liquid interface, microscopic dynamics, parallel computing

**2010 Mathematics Subject Classification:** 76D45, 76S05

**References:**

[1] Meirmanov A., Mathematical models for poroelastic flows, Atlantis Press, Paris, 2013.
[2] Malvern L.E., Introduction to Mechanics of a Continuum Medium, Prentice–Hall, Inc. Englewood Cliffs, N.J. (1969).

# MS2

# SPECTRAL THEORY OF DIFFERENTIAL EQUATIONS

# The first regularized trace of integro-differential Sturm-Liouville operator on the segment with punctured points at integral perturbation of transmission conditions

Aizat AIMAKHANOVA [1], Saltanat SHALGINBAYEVA [2], Lyazzat ZHUMANOVA [3]

[1] *Institute of Mathematics and Mathematical Modeling, Kazakhstan, and Asfendiyarov Kazakh National Medical University, Kazakhstan*
*E-mail: aizat.68@mail.ru*
[2] *Kazakh Ablai Khan University of International Relations and World Languages, Kazakhstan*
*E-mail: salta_sinar@mail.ru*
[3] *Al-Farabi Kazakh National University, Kazakhstan*
*E-mail: lzhumanova@mail.ru*

**Abstract:** The report is devoted to calculating a first regularized trace of one integro-differential operator with the main part of the Sturm-Liouville type on a segment with punctured points at integral perturbation of "transmission" conditions. The Sturm-Liouville operator

$$-y''(x) + q(x)y(x) + \gamma \int_0^\pi y(t)dt = \lambda y(x)$$

given on the segments $\frac{\pi}{n}(k-1) < x < \frac{\pi}{n}k$, $k = \overline{1,n}$; $n \geq 2$ is considered. Boundary conditions of the Dirichlet type:

$$y(0) = 0, \quad y(\pi) = 0$$

are given on the left-hand and right-hand ends of the segment $[0, \pi]$. The functions continuous on $[0, \pi]$, the first derivatives of which have jumps at the points $x = \frac{\pi}{n}k$, are solutions. The value of jumps is expressed by the formula

$$y'\left(\frac{\pi k}{n} - 0\right) = y'\left(\frac{\pi k}{n} + 0\right) - \beta_k \int_0^\pi y(t)dt, \quad k = \overline{1, n-1}.$$

The basic result of the report is the exact formula of the first regularized trace of the considered differential operator.

**Keywords:** spectral problem, first regularized trace, integro-differential operator, inner-boundary condition

**2010 Mathematics Subject Classification:** 34L10, 34L15, 34L20

# On the solvability of differential equations in logarithmic scale of Hilbert spaces

Shavkat ALIMOV

*MIMOS, Kuala Lumpur, Malaysia*
*E-mail: shavkat.alimov@mimos.my*

**Abstract:** Consider a Hilbert space $H$ and selfadjoint semibounded operator $\Lambda$ with lower bound $\mu > 1$. This operator generates the scale of Hilbert spaces $H^\alpha = D(\Lambda^\alpha)$ with the norm $\|u\|^{(\alpha)} = \|\Lambda^\alpha u\|$.

With $H^\alpha$ we consider the logarithmic scale of Hilbert spaces $L^s$ with norms $\|u\|_{(s)} = \|(\ln \Lambda)^s u\|$.

Consider the Cauchy problem

(1) $u''(t) - Au(t) = f(t), \quad u(0) = u_0, \quad u'(0) = u_0',$

where $A : H \to H$ is an unbounded linear operator. We assume that $A$ is a continuous operator from $L^s$ to $L^{s+2}$.

Let $C([0,T] \to H^\alpha)$ denote the space of all continuous functions $u : [0,T] \to H^\alpha$.

**Theorem 1.** *Let $\alpha > 0$ and $T > 0$. For any $f \in C([0,T] \to H^\alpha)$ and $u_0, u_0' \in H^\alpha$ there exists unique solution of the Cauchy problem (1) from $C([0,T] \to H^\alpha)$.*

As application to the hypersingular integro-differential equations consider $H = L_2(\mathbb{R}^n)$ and $\Lambda f(x) = F^{-1}(1+|\xi|^2)^{1/2} F f(x)$, where $F$ is Fourier transform. In this case $H^\alpha$ coincides with Sobolev space $W_2^\alpha(\mathbb{R}^n)$.

Let $A$ be the singular integral operator with kernel (see [1], page 50)

(2) $K(x,y) = [(x-y) \otimes (x-y)]|x-y|^{-n-2}, \quad x, y \in \mathbb{R}^n, n \geq 3.$

The integral operator with kernel (2) maps $L^s$ to $L^{s+2}$ continuously (see [2], Proposition 3.5). Moreover, the convolution $K * K$ has the same property. Hence, it follows from theorem 1 that the corresponding equation has solution in $W_2^\alpha(\mathbb{R}^n)$ for any $\alpha > 0$.

**Keywords:** integro-differential equation, singular integral, peridynamics.

**2010 Mathematics Subject Classification:** 45J05, 46N20.

**References:**

[1] E. Emmrich,E R. Lehoucq and D. Puhst, Peridynamics: A nonlocal continuum theory, Lect. Notes Comp. Sci. Eng. 89 (2013), 45-65.

[2] S. A. Alimov, Y. Cao, and O. A. Ilhan. *On the problem of peridynamics with special convolution kernels*, Journal of Integral Equations and Applications, Vol. 26, No 3 (2014), pp. 301-321.

## On spectral problems for loaded two-dimension Laplace operator

Meiramkul AMANGALIYEVA [1], Muvasharkhan JENALIYEV [1], Kanzharbek IMANBERDIYEV [1], Murat RAMAZANOV [2]

[1] *Institute of Mathematics and Mathematical Modeling, Àlmaty, Kazakhstan*
*E-mail: muvasharkhan@gmail.com*
[2] *Buketov Karaganda Stated University, Karaganda, Kazakhstan*
*E-mail: ramamur@mail.ru*

**Abstract:** In domain $Q = \{x, y : -\pi < x < \pi,\ -\pi < y < \pi\}$ we consider following two spectral problems:

$$(1)\quad \begin{cases} -\triangle\varphi(x,y) + \alpha\varphi(0,y) = \lambda\varphi(x,y), \ \ \{x,y\} \in Q, \\ \dfrac{\partial^j \varphi(-\pi,y)}{\partial x^j} = \dfrac{\partial^j \varphi(\pi,y)}{\partial x^j}, \ \ \dfrac{\partial^j \varphi(x,-\pi)}{\partial y^j} = \dfrac{\partial^j \varphi(x,\pi)}{\partial y^j}, \ \ j = 0,\ 1; \end{cases}$$

$$(2)\quad \begin{cases} -\triangle\varphi(x,y) + \alpha\varphi(0,y) + \beta\varphi(x,0) = \lambda\varphi(x,y), \ \ \{x,y\} \in Q, \\ \dfrac{\partial^j \varphi(-\pi,y)}{\partial x^j} = \dfrac{\partial^j \varphi(\pi,y)}{\partial x^j}, \ \ \dfrac{\partial^j \varphi(x,-\pi)}{\partial y^j} = \dfrac{\partial^j \varphi(x,\pi)}{\partial y^j}, \ \ j = 0,\ 1; \end{cases}$$

where $\Delta$ is Laplace operator, $\alpha$, $\beta \in \mathbb{C}$ are given complex numbers, $\lambda \in \mathbb{C}$ is a spectral parameter.

Note that the need to investigate of spectral problems (1) and (2) arise in solving of the stabilization problem for a loaded heat equation with help of the boundary control actions.

Depending on the values of $\alpha$ and $\beta$, we give a complete description of the eigenvalues, eigenfunctions and associated functions of the set of spectral problems (1) and (2).

**Keywords:** Loaded Laplace operator, spectrum, eigenfunction

**2010 Mathematics Subject Classification:** 35J05, 35P05, 35R10

**References:**

[1] M.T. Jenaliyev, M.I. Ramazanov, "Stabilization of the solution of heat equation, loaded on a zero-dimensional manifolds with using boundary conditions (in Russian)", *Matematicheskii zhurnal*, Vol. 15, No. 4 (58) (2015), p. 33–53.

# Spectrum of Volterra integral operator of the second kind

Meiramkul AMANGALIYEVA [1], Muvasharkhan JENALIYEV [1], Madi ERGALIYEV [1], Murat RAMAZANOV [2]

[1] *Institute of Mathematics and Mathematical Modeling, Àlmaty, Kazakhstan*
*E-mail: muvasharkhan@gmail.com*
[2] *Buketov Karaganda Stated University, Karaganda, Kazakhstan*
*E-mail: ramamur@mail.ru*

**Abstract:** In this paper we consider the singular Volterra integral equation with spectral parameter $\lambda \in \mathbb{C}$ of form

$$\varphi(t) - \lambda \int_0^t K(t,\, \tau)\, \varphi(\tau)\, d\tau = f(t), \quad t > 0, \tag{1}$$

where

$$K(t,\, \tau) = K^{(1)}(t,\, \tau) + K^{(2)}(t,\, \tau), \tag{2}$$

$$K^{(1)}(t,\, \tau) = \frac{1}{2a\sqrt{\pi}} \cdot \frac{t^{\omega} + \tau^{\omega}}{(t-\tau)^{\frac{3}{2}}} \exp\left(-\frac{(t^{\omega} + \tau^{\omega})^2}{4a^2(t-\tau)}\right), \tag{3}$$

$$K^{(2)}(t,\, \tau) = \frac{1}{2a\sqrt{\pi}} \cdot \frac{t^{\omega} - \tau^{\omega}}{(t-\tau)^{\frac{3}{2}}} \exp\left(-\frac{(t^{\omega} - \tau^{\omega})^2}{4a^2(t-\tau)}\right), \ \omega\, 1/2. \tag{4}$$

We call such equations as the Volterra integral equations with 'incompressible' kernel [1]. It is shown that the corresponding homogeneous equation on $|\lambda| \geq \exp\{|\arg\lambda|\}, \quad \arg\lambda \in [-\pi, \pi]$ has a continuous spectrum, and the multiplicity of the characteristic numbers grows with increasing $|\lambda|$. We use the Carleman-Vekua regularization method. We introduce the characteristic integral equation. We prove that the initial integral equation has eigenfunctions, the multiplicity of which depends on the value of the spectral parameter $\lambda$. We prove the solvability theorem of the nonhomogeneous equation (1)–(4) in a case when the right-hand side of the equation belongs to a certain class.

**Keywords:** Volterra integral equation, spectrum, eigenfunction

**2010 Mathematics Subject Classification:** 45C05, 45D05, 45E10

**References:**

[1] M.T. Jenaliyev, M.M. Amangaliyeva, M.T. Kosmakova, and M.I. Ramazanov, "On a Volterra equation of the second kind with 'incompressible'", *Advances in Difference Equations*, Vol. 2015:71 (2015), 14p.

# On a regular problem for an elliptic-parabolic equation with a potential boundary condition

Gaukhar AREPOVA

*Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: arepovag@mail.ru*

**Abstract:** In this paper we construct a lateral boundary condition for an elliptic-parabolic equation in a finite domain. Theorem on existence and uniqueness of the solution of the problem considered are proved by method of theory potential. For construction of problem we use the Bitsadze-Samarskii problem with boundary conditions.

Bitsadze-Samarskii conditions binds internal traces $u(x,0)$ and $\frac{\partial u}{\partial \tau}(x,0)$ with boundary conditions $u(x,t)$ and $\frac{\partial u}{\partial \tau}(x,t)$ on the boundary of the domain.

**Theorem** *For each*

$$f(x,t) \in C^{\alpha}(\overline{D^{-}}) \bigcap C^{\alpha}(\overline{D^{+}})$$

*Bitsadze-Samarskii problem has a unique solution* $u(x,t)$ *and*

$$u(x,t) \in C^{\alpha}(D) \bigcap C^{2+\alpha}(\overline{D^{-}}) \bigcap C_{x,t}^{2+\alpha,1+\alpha}(\overline{D^{+}}).$$

Throughout this note we mainly use techniques from our works [1-3].

**Keywords:** an elliptic-parabolic equation, Bitsadze-Samarskii problem, lateral boundary condition

**2010 Mathematics Subject Classification:** 35Q79, 35K05, 35K20

**References:**

[1] T.Sh. Kal'menov, D. Suragan, On spectral problems for the volume potential. *Doklady Mathematics*. **80**:2. 646–649 (2009).

[2] T. Sh. Kalmenov and D. Suragan, *A boundary condition and Spectral Problems for the Newton Potentials,* Operator Theory: Advances and Applications, Vol. 216 (2011), pp.

[3] T. Sh. Kal'menov and N. E. Tokmagambetov, *On a nonlocal boundary value problem for the multidimensional heat equation in a noncylindrical domain*, Siberian Mathematical Journal, **54** (6) (2013), pp. 1023–1028.

## Structure of fractional spaces generated by the two dimensional neutron transport operator

Allaberen ASHYRALYEV [1], Abdulgafur TASKIN

[1] *Emeritus Professor, Turkey & Turkmenistan*
*E-mail: aallaberen@gmail.com, gafurtaskin@hotmail.com*

**Abstract:** In this study, the structure of fractional spaces generated by the two-dimensional neutron transport operator $A$ defined by formula $Au = \omega_1 \frac{\partial u}{\partial x} + \omega_2 \frac{\partial u}{\partial y}$ is investigated.The positivity of $A$ in $C\left(R^2\right)$ and $L_p\left(R^2\right)$, $1 \leq p < \infty$ is established. The structure of fractional spaces generated by the operator $A$ is studied. It is established that for any $0 < \alpha < 1$ and $1 \leq p < \infty$ the norms in the spaces $E_{\alpha,p}\left(L_p\left(R^2\right),\ A\right)$ and $E_\alpha\left(C\left(R^2\right),\ A\right)$, $W_p^\alpha\left(R^2\right)$ and $C^\alpha\left(R^2\right)$ are equivalent, respectively. The positivity of the neutron transport operator in Hölder space $C^\alpha\left(R^2\right)$ and Slobodeckij space $W_p^\alpha\left(R^2\right)$ is proved.

**Keywords:** neutron transport operator, positive operator, fractional spaces, Hölder spaces, Slobodeckij spaces

**2015 Mathematics Subject Classification:** 94B05, 94B15

**References:**

[1] A. Ashyralyev, N. Nalbant, and Y. Sozen, J. Franklin Inst. Eng. Appl. Math. 351(2), 713-731 (2014).

[2] A. Ashyralyev, S. Akturk, and Y. Sozen, Bound. Value Probl. 2014, 1-17 (2014), (DOI: 10.1186/1687-2770-2014-3).

[3] A. Ashyralyev, and F. S. Tetikoglu, Bound. Value Probl. 2015, 1-17 (2015), (DOI: 10.1186/s13661-015-0293-9).

[4] A. Ashyralyev, *Fractional spaces generated by the positive differential and difference operators in a Banach Space*, ISMME, Springer, 2006,10-19.

[5] G. E. Semenova, *Positivity of elliptic difference operators and its applications*, AIP Conference Proceedings, 1470, 232-235 (2012).

[6] A. Ashyralyev, *A survey of results in the theory of fractional spaces generated by positive operators*, TWMS J. Pure Appl. Math. V.6, N.2, 2015, pp.129-157.

[7] A. Ashyralyev, and A. Yakubov, In: Modeling the Proceesses in Exploration of Gas Deposits and Applied Problems of Theoretical Gas Hydrodynamics, Ilim, Ashgabat, 1995, pp. 162-172.

[8] A. Ashyralyev, and A. Taskin, *Stable Difference Schemes for the Neutron Transport Equation* AIP Conference Proceedings 1389, 570 (2011); (DOI: 10.1063/1.3636794).

[9] A. Ashyralyev, and A. Taskin, *Structures of the fractional spaces generated by the difference neutron transport operator* AIP Conference Proceedings 1676, 020051 (2015); (DOI: 10.1063/1.4930477).

# Green's function of the second order differential operator with involution

Allaberen ASHYRALYEV [1], Abdisalam SARSENBI [2]

[1] *Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan*
*E-mail: aallaberen@gmail.com*
[2] *Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan*
*M. Auezov South-Kazakhstan State University, Shymkent, Kazakhstan*
*E-mail: abdisalam@mail.ru*

**Abstract:** In the present paper, we are interested in the Green's function of the differential operator $L$ defined by formula

$$Lu = \alpha u''(x) - u''(-x) - \lambda u(x), -1 < x < 1 \tag{1}$$

with Dirichlet conditions

$$u(-1) = 0, u(1) = 0 \tag{2}$$

Note that a function $u(x)$ is called a *solution* of boundary value problem with the positive parameter $\lambda$ for the differential equation

$$Lu = f(x), -1 < x < 1 \tag{3}$$

with Dirichlet conditions (2), if it satisfies this equation (3) and boundary conditions (2). Here $f(x)$ is a continuous function defined on [-1,1].

A function $G(x,t;\lambda)$ is called the *Green's function* of the differential operator $L$ defined by formula (1) and conditions (2), if the function

$$\int_{-1}^{1} G(x,t;\lambda) f(t)\, dt$$

is a solution of boundary value problem (2)-(3).

The main findings of the report: we have the following point wise estimates for the Green's function $G(x,t;\lambda)$:

$$|G(x,t;\lambda)| \le \frac{M(\alpha)}{\sqrt{\lambda}} \begin{cases} e^{-c_0(\alpha)\sqrt{\lambda}(-t-|x|)}, -1 \le t \le -|x|, \\ e^{-c_0(\alpha)\sqrt{\lambda}(|x|-|t|)}, -|x| \le t \le |x|, \\ e^{-c_0(\alpha)\sqrt{\lambda}(t-|x|)}, |x| \le t \le 1. \end{cases}$$

**Keywords:** differential equation with involution, Green's function

**2010 Mathematics Subject Classification:** 34B27, 34L10, 34L15

## Bounded perturbations of the correct restrictions and extensions

Bazarkan N. BIYAROV [1], Gulnara K. ABDRASHEVA [2]

[1] *Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: bbiyarov@gmail.com*
[2] *L. N. Gumilyov Eurasian National University, Astana, Kazakhstan*
*E-mail: gulnara.abdrash@gmail.com*

**Abstract:** This paper considers the correct restrictions of the maximal operator. We obtain a criterion of relative-boundedness of the perturbations of the inverse operator to a fixed correct restrictions. Given an applications of this criterion to some differential operators.

Under a weak perturbation of the compact operator $T$ will be called the operator $T(I+S)$, where $S$ is such a compact operator that the operator $I+S$ is continuously invertible. A lot of work (see [1], [2], [3]– [7]) is devoted to the spectral properties of a weak perturbation of a compact operator $T$.

If $S$ is a bounded operator, then the perturbed operator will be called the bounded perturbation. This work is devoted to describe among the correct restrictions and extensions of the bounded perturbations of the fixed operator.

**Keywords:** bounded perturbations, correct restrictions, correct extensions

**2010 Mathematics Subject Classification:** 46Nxx, 47Fxx, 35P05

**References:**

[1] Keldysh, M. V., "On the eigenvalues and eigenfunctions of certain classes of non-selfadjoint equations", *Dokl. Akad. Nauk SSSR*, Vol. 77, No.1, pp. 11–14, 1951.

[2] Matsaev, V. I., Mogul'skii, E. Z., "On the possibility of a weak perturbation of a complete operator to a Volterra operator", *Dokl. Akad. Nauk SSSR*, Vol. 207, No. 3, pp. 534–537, 1972; (English transl.: *Soviet Math. Dokl.*, Vol. 13, pp. 1565–1568, 1972).

[3] Kal'menov, T. Sh., Suragan, D., "Boundary conditions for the volume potential for the polyharmonic equation", *Differential Equations*, Vol. 48, No.4, pp. 604–608, 2012.

[4] Kal'menov, T. Sh., Sadybekov, M. A., Sarsenbi, A. M., "Biorthogonal Expansions in Root Functions of Differential Operators", *Differential Equations* Vol. 47, No.1, pp. 144–148, 2011.

[5] Kal'menov, T. Sh., Suragan, D., "To spectral problems for the volume potential", *Doklady Mathematics*, Vol. 80, No.2, pp. 646–649, 2009.

[6] Kal'menov, T. Sh., Kanguzhin, B. E., Koshanov, B. D., "On integral representations of correct restrictions and regular extensions of differential operators", *Doklady Mathematics*, Vol. 81, No.1, pp. 94–96, 2010.

[7] Kal'menov, T. Sh., Suragan D. , "Determination of the structure of the spectrum of regular boundary value problems for differential equations by VA Il'in's method of anti-a priori estimates", *Doklady Mathematics*, Vol. 78, No.3, pp. 913–915, 2008.

# On spectrum of perturbed two-dimensional harmonic oscillator in a strip

Ziganur FAZULLIN [1], Irina NUGAEVA [2]

[1] *Department of Mathematics and Information Technologies, Bashkir State University, Ufa, Russia*
*E-mail: fazullinzu@mail.ru*
[2] *Department of Mathematics and Information Technologies, Bashkir State University, Ufa, Russia*
*E-mail: nuga-irina@yandex.ru*

**Abstract:**

Let $\Pi = \{(x,y), x \in \mathbb{R}, 0 \le y \le \pi\}$, $L_0 = -\frac{\partial^2}{\partial x^2} - \frac{\partial^2}{\partial y^2} + x^2$. We denote by $V$ the operator of multiplication by bounded measurable real-valued function $V(x,y)$, which is finite on the variable $x$. That is, there exists $r > 0$ with the property: $V(x,y) \equiv 0$, for any $x$, $|x| \ge r$. In our further considerations such functions $V(x,y)$ are said to be *finite in the strip* $\Pi$. We denote $\operatorname{supp} V$ by $\Pi_0(r)$. Here, we deal with the Dirichlet problem operator $L = L_0 + V$ in the space $L^2(\Pi)$. Notice that the title of the paper is just prompted by the possibility to consider the expression $(\widetilde{V}(x,y) + y^2)$, $y \in [0,\pi]$, instead of $V(x,y)$, without loss of generality.

Let $P_s^{(1)}$ $P_l^{(2)}$ be orthogonal projectors onto eigensubspaces of one-dimensional Laplace operators of Dirichlet problem and harmonic oscillator corresponding to the eigenvalues $s^2$, $s = 1, 2...$, and $2l+1$, $l = 0, 1, 2...$, correspondingly. That is, $P_s^{(1)} f = (f, f_s) f_s$, $f_s(y) = \sqrt{\frac{2}{\pi}} \sin sy$ $P_l^{(2)} f = (f, \varphi_l)\varphi_l$, $\varphi_l(x) = (2^l l! \sqrt{\pi})^{-\frac{1}{2}} e^{-\frac{x_2}{2}} H_l(x)$, where $H_l(x)$ are Hermitian polynomials.

Then, we have the following assertion on the eigenvalues $\lambda_{sl} = s^2 + 2l + 1$, $s = 1, 2, ...$, $l = 0, 1, 2, ...$ of the operator $L_0$

**Theorem 1.** The spectrum of the operator $L_0$ consists of the eigenvalues $\lambda_n = n$, $n \in \mathbb{N} \setminus \{1, 3\}$, with multiplicities $\nu_n$

$$\nu_n = \begin{cases} [\frac{\sqrt{n}}{2}], & \lambda_n \in \left((2[\frac{\sqrt{n}}{2}])^2; (2[\frac{\sqrt{n}}{2}]+1)^2\right) \\ [\frac{\sqrt{n}}{2}] + \frac{(-1)^n+1}{2}, & \lambda_n \in \left((2[\frac{\sqrt{n}}{2}]+1)^2; (2[\frac{\sqrt{n}}{2}]+2)^2\right) \end{cases}.$$ The corresponding projector operators are $P_n = \sum\limits_{s=1}^{\nu_n} P_s^{(1)} \otimes P^{(2)}_{(\frac{n}{2} - \frac{s^2+1}{2})}$.

For each $n \in \mathbb{N} \setminus \{1, 3\}$ we denote by $\mu_s^{(n)}$, $s = 1, \dots, \nu_n$, the eigenvalues of the operator $L$ corresponding to the eigenvalue $\lambda_n$ of the operator $L_0$. We have established that there exists $d > 0$ with the property: for $n >> 1$ the inequalities $\left|\mu_s^{(n)} - \lambda_n\right| \le d n^{-\frac{1}{2}}$ hold for any $s = 1, \dots, \nu_n$.

Using the method of our paper [1] we get our main result:

**Theorem 2.** Assume that $V(x,y) \in C^4(\Pi_0(r))$. Then,

$$\sum_{k\in\mathbb{N}\setminus\{1,3\}}\left[\sum_{s=1}^{\nu_k}\left(\lambda_k-\mu_s^{(k)}\right)+\operatorname{sp}P_kV\right]=\frac{1}{12\pi}\iint\limits_{\Pi}V^2(x,y)dxdy.$$

First author was supported by the Ministry of Education and Science of Russian Federation, project no. 01201456408.

**References:**

[1] Fazullin, Z.Yu., Murtazin, Kh.Kh., "Regularized trace of a two-dimensional harmonic oscillator", *Sbornik Mathematics*, Vol.192, No.5, p. 725, 2001.

---

## Dirac systems that contain discontinuity conditions

Tuba GULSEN [1], Etibar S. PANAKHOV [1,2]

[1] *Department of Mathematics, Firat University, Elazig, Turkey*
*E-mail: tubagulsen87@hotmail.com, epenahov@firat.edu.tr*
[2] *Institute of Applied Mathematics, Baku State University, Azerbaijan*

**Abstract:** Boundary value problems which contain transmission conditions inside a finite interval have been studied in [1–3]. This type of problems may have discontinuities in the solution or its derivative at an inner point of the interval and arise in many areas of natural sciences. For example, heat and mass transfer, mechanics, electronics, radio, geophysics. Direct and inverse problems for Dirac differential equation with transmission conditions have been studied in [4–6]. The aim of this paper is to study a Dirac system which contain transmission conditions inside a finite interval.

**Keywords:** Dirac systems, transmission conditions, discontinuous boundary-value problems, eigenvalues, eigen-vector functions

**2010 Mathematics Subject Classification:** 34L05, 34L40.

**References:**

[1] N. Altinisik,M. Kadakal,O. Sh. Mukhtarov,*Acta Mathematica Hungarica* **102**, 159–175 (2004).
[2] M. Kadakal,O. Sh. Mukhtarov,*Computers and Mathematics with Applications* **54**, 1367–1379 (2007).
[3] A. A. Nabiev,R. K. Amirov,*Math. Methods Appl. Sci.* **36**, 1685–1700 (2012).
[4] N. B. Kerimov,*Differential Equations* **38**, 164–174 (2002).
[5] B. Keskin,A. S. Ozkan,*Acta Mathematica Hungarica* **130**, 309–320 (2011).
[6] M. M. Tharwat,A. Yildirim,A. H. Bhrawy,*Numerical Functional Analysis and Optimization* **34**, 323–348 (2013).
[7] B. M. Levitan,I. S. Sargsjan,*Sturm-Liouville and Dirac Operators*, Kluwer Academic, Dordrecht, The Netherlands, 1991, pp. 1–2.

## Stability of basis property of a type of problems with nonlocal perturbation of boundary conditions

Nurlan IMANBAEV [1], Makhmud SADYBEKOV [2]

[1] *Institute of Mathematics and Mathematical Modeling, Kazakhstan, and South Kazakhstan State Pedagogical Institute, Kazakhstan*
*E-mail: imanbaevnur@mail.ru*
[2] *Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: sadybekov@math.kz*

**Abstract:** This report is devoted to a spectral problem for a multiple differentiation operator with an integral perturbation of boundary conditions of one type which are regular, but not strongly regular:

$$l(u) \equiv -u''(x) = \lambda u(x),\ 0 < x < 1, \tag{2}$$

$$U_1(u) \equiv u'(0) - u'(1) + \alpha u(0) = \int_0^1 \overline{p(x)} u(x) dx,\ p(x) \in L_2(0,1), \tag{3}$$

$$U_2(u) \equiv u(0) - u(1) = 0. \tag{4}$$

Here $\alpha \neq 0$ is an arbitrary complex number.

The unperturbed problem ($p(x) \equiv 0$) has an asymptotically simple spectrum, and its system of normalized eigenfunctions creates the Riesz basis. We construct the characteristic determinant of the spectral problem with an integral perturbation of the boundary conditions. The perturbed problem can have any finite number of multiple eigenvalues. Therefore, its root subspaces consist of its eigen and (maybe) adjoint functions. It is shown that the Riesz basis property of a system of eigen and adjoint functions is stable with respect to integral perturbations of the boundary condition.

Throughout this note we mainly use techniques from our works [1].

**Keywords:** Riesz basis, regular boundary conditions, eigenvalues, root functions, spectral problem, integral perturbation of boundary condition, characteristic determinant

**2010 Mathematics Subject Classification:** 35J05, 35J08, 35J25, 35P05

**References:**

[1] Sadybekov M.A., and Imanbaev N.S., "On the basis property of root functions of a periodic problem with an integral perturbation of the boundary condition", *Differential Equations*. Vol.48, No.6, pp. 896–900, 2012.

# On a homogeneous parabolic problem in an infinite corner domain

Muvasharkhan JENALIYEV [1], Murat RAMAZANOV [2]

[1] *Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan*
*E-mail: muvasharkhan@gmail.com*
[2] *Buketov Karaganda Stated University, Karaganda, Kazakhstan*
*E-mail: ramamur@mail.ru*

**Abstract:** It is considered the homogeneous boundary problem

$$\frac{\partial u(x,t)}{\partial t} - a^2 \frac{\partial^2 u(x,t)}{\partial x^2} = 0,\ \{x,t\} \in G = \{x,t:\ 0 < x < t,\ t > 0\}; \tag{1}$$

$$\frac{\partial u(x,t)}{\partial x}|_{x=0} = 0,\ \frac{\partial u(x,t)}{\partial x}|_{x=t} + \frac{d\tilde{u(t)}}{dt} = 0; \tag{2}$$

where $\tilde{u}(t) = u(t,t)$.

Note that problem (1)–(2) is homogeneous case of the problem, studied in [1], which stated that the case of non-homogeneous boundary value problem "... it appears to be useful in the study of some free boundary value problems. "In the paper [1] was obtain a theorem on the unique solvability of the non-homogeneous boundary value problem in weighted Holder spaces.

In this paper, we set in class of essentially bounded functions with a given weight the existence of a nontrivial solution for a constant factor and a constant term. We introduce the class as follows:

$$(x + t^{1/2})^{-1}\, u(x,t) \in L_\infty(G),\ \text{i.e. } u(x,t) \in L_\infty(G;\, (x + t^{1/2})^{-1}).$$

THEOREM 1. *The boundary value problem (1)–(2) has a nontrivial solution* $u(x,t) = C_2\, \tilde{u}(x,t) + C_1$, *where* $\tilde{u}(x,t) \in L_\infty(G;\, (x + \sqrt{t})^{-1})$, $C_1, C_2 = const$.

THEOREM 2. *In the class of functions* $L_\infty\left(G;\, [x^{1+\alpha} + t^{(1+\alpha)/2}]^{-1}\right)$ *the boundary value problem (1)–(2) has only the trivial solution* $u(x,t) \equiv 0$.

**Keywords:** parabolic equation, boundary value problem, integral equation

**2010 Mathematics Subject Classification:** 94B05, 94B15

**References:**

[1] V. A. Solonnikov, and A. Fasano, "One-dimensional parabolic problem arising in the study of some free boundary problems", *Zapiski nauchnykh seminarov POMI (in Russian)*, Vol. 269, pp.322–338, 2000.

## Spectral geometry inequalities for Schatten $p$-norms of compact operators

Tynysbek Sh. KALMENOV, Durvudkhan SURAGAN and Bolys SABITBEK

*Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: suragan@math.kz*

**Abstract:** In this talk we give answers to some spectral geometry questions of Schatten $p$-norms of convolution type compact operators on complete, connected, simply connected Riemannian manifolds of constant sectional curvature. For example, we show that among all domains of a given measure the geodesic ball is a maximizer of Schatten $p$-norms of some convolution type integral operators. The main reason why the results are useful, beyond the intrinsic interest of geometric extremum problems, is that they produce a priori bounds for spectral invariants of operators on arbitrary domains. This talk is mainly based on the papers [1]- [5].

**Keywords:** Convolution operators, Schatten $p$-norm, $n$-sphere, real hyperbolic space, Rayleigh-Faber-Krahn inequality, Hong-Krahn-Szego inequality

**2010 Mathematics Subject Classification:** 35P99, 47G40, 35S15

**References:**

[1] Kal'menov T.Sh., and Suragan D., "To spectral problems for the volume potential", *Doklady Mathematics (translated from Doklady Akademii Nauk)*, doi: 10.1134/S1064562409050032, Vol.80(2), pp. 646–649, 2009.

[2] Rozenblum G., Ruzhansky M., and Suragan D., "Isoperimetric inequalities for Schatten norms of Riesz potentials", *Journal of Functional Analysis*, doi: 10.1016/j.jfa.2016.04.023, 2016.

[3] Ruzhansky M., and Suragan D., "On first and second eigenvalues of Riesz transforms in spherical and hyperbolic geometries", *Bulletin of Mathematical Sciences*, doi: 10.1007/s13373-016-0082-5, 2016.

[4] Ruzhansky M., and Suragan D., "On Schatten norms of convolution-type integral operators", *Russian Mathematical Surveys (translated from Uspekhi Matematicheskikh Nauk)*, doi: 10.1070/RM9665, Vol.71(1), 2016.

[5] Ruzhansky M., and Suragan D., "Isoperimetric inequalities for the logarithmic potential operator", *Journal of Mathematical Analysis and Applications*, doi:10.1016/j.jmaa. 2015.07.041, Vol.434(2), pp. 1676-1689, 2016.

# On a heat transfer model for the locally inhomogeneous initial data

Tynysbek KAL'MENOV [1], Gaukhar AREPOVA [2]

[1] *Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: kalmenov@math.kz*
[2] *Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: arepovag@mail.ru*

**Abstract:** We consider a model case of the problem of heat diffusion in a homogeneous body with a special initial state. The peculiarity of this initial state is its local inhomogeneity. That is, there is a closed domain $\Omega$ inside a body, the initial state is constant out of the domain. Mathematical modeling leads to the problem for a homogeneous multi-dimensional diffusion equation.

Let $\Omega \subset \mathbb{R}^n$ be a bounded domain with sufficiently smooth boundary $\partial\Omega$. Through $D$ we denote a cylindrical domain $D = \Omega \times (0, T)$. In $D$ we consider a surface heat potential

$$u = \int_{\Omega} \varepsilon_n(x - \xi, t) u_0(\xi) d\xi, \tag{1}$$

where $\varepsilon_n(x,t) = \theta(t)(2a\sqrt{\pi t})^{-n} e^{-\frac{|x|^2}{4a^2 t}}$ is a fundamental solution of the heat equation

$$\Diamond u \equiv \left(\frac{\partial}{\partial t} - a^2 \triangle_x\right) u = 0. \tag{2}$$

We construct the boundary conditions on the boundary of the domain $\Omega$, which can be characterized as "transparent" boundary conditions . We separately consider a special case – a model of redistribution of heat in a uniform linear rod, the side surface of which is insulated in the absence of (internal and external) sources of heat and of locally inhomogeneous initial state.

Throughout this note we mainly use techniques from our works [1, 2].

**Keywords:** diffusion equation, homogeneous body, initial state, local inhomogeneity, transparent boundary conditions

**2010 Mathematics Subject Classification:** 35Q79, 35K05, 35K20

**References:**

[1] Kal'menov T.Sh., and Suragan D., "Boundary conditions for the volume potential for the polyharmonic equation" *Differential Equations*. Vol.48, No.4, pp. 604–608, 2012.

[2] Kal'menov T.Sh. and Otelbaev M., "Boundary Criterion for Integral Operators", *Doklady Mathematics*. Vol.93, No.1, pp. 58–61, 2016.

# A uniqueness theorem of a boundary inverse problem of a differential operator on an interval with integro–differential boundary conditions

Baltabek KANGUZHIN [1a,2], Niyaz TOKMAGAMBETOV [1b,3]

[1] *Al–Farabi Kazakh National University, Almaty, Kazakhstan*
[a] *E-mail: kanbalta@mail.ru,* [b] *E-mail: niyaz.tokmagambetov@gmail.com*
[2] *Institute of Mathematics and Mathematical Modeling, Kazakhstan*
[3] *Imperial College London, United Kingdom*
*E-mail: n.tokmagambetov@imperial.ac.uk*

**Abstract:** In this work we research a boundary inverse problem of spectral analysis of a differential operator with integral boundary conditions in the functional space $\mathrm{L}_2(0,b)$, where $b < \infty$. A uniqueness theorem of the inverse boundary problem in $\mathrm{L}_2(0,b)$ is proved. Note that a boundary inverse problem of spectral analysis is the problem of recovering boundary conditions of the operator by its spectrum and some additional data.

In this paper we investigate the inverse problem of spectral analysis of high order differential operators with integro–differential boundary conditions. In this case, it is necessary to find from the spectral data not only coefficients of the differential expression, also, we need to find boundary functions of the integro–differential boundary conditions. Coefficient inverse problems are well studied. Therefore in this paper we study the issue of reconstruction of boundary functions.

**Keywords:** boundary inverse problem, uniqueness theorem, differential operator, integro–differential boundary condition

**2010 Mathematics Subject Classification:** 47A55, 35P25

## On identities for eigenvalues of a well-posed pertubation of the Laplace operator in a punctured domain

Baltabek KANGUZHIN [1a,2], Gulzat NALZHUPBAYEVA [1b,3]

[1] *Department of Fundamental Mathematics,Al–Farabi Kazakh National University, Almaty, Kazakhstan*
[a] *E-mail: kanbalta@mail.ru,* [b] *E-mail: nalzhuppa@list.ru*
[2] *Institute of Mathematics and Mathematical Modeling, Kazakhstan*
[3] *Imperial College London, United Kingdom*
*E-mail: n.gulzat@imperial.ac.uk*

**Abstract:** In this paper we consider a class of finite perturbed well-posed problems for the Laplace operator. And, identities for eigenvalues of the considering problems are found.

In this paper we are going to continue research of the chain of papers [1–5], where a class of well-posed problems in a Hilbert space for the m-Laplace operator in a punctured domain is considered and an analog of the Green formula and a class of self-adjoint problems are obtained by the first author and his co–authors. Our aim is to obtain some identities for eigenvalues of well-posed problems for the Laplace operator in a punctured domain.

**Keywords:** Laplace operator, well-posed solvable boundary value problem, punctured domain, Dirichlet boundary condition, finite perturbation, identities for eigenvalues, resolvent, non-local type boundary value problem, nuclear operator

**2010 Mathematics Subject Classification:** 35P15

**References:**

[1] B. E. Kanguzhin, and A. A. Aniyarov, "Well-posed problems for the Laplace operator in a punctured disk", *Mathematical Notes*, Vol. 89, No. 5–6, pp. 819–829, 2011.

[2] B. E. Kanguzhin, D. B. Nurakhmetov, and N. E.Tokmagambetov, "Laplace operator with $\delta$–like potentials", *Russian Mathematics*, Vol. 58, No. 2, pp. 6–12, 2014.

[3] B. E. Kanguzhin, and N. E. Tokmagambetov, "A Regularized Trace Formula for a Well-Perturbed Laplace Operator", *Doklady Mathematics*, Vol. 91, No. 1, pp. 1–4, 2015.

[4] B. E. Kanguzhin, and N. E. Tokmagambetov, "On Regularized Trace Formulas for a Well-Posed Perturbation of the m-Laplace Operator", *Differential Equations*, Vol. 51, No. 12, pp. 1583–1588, 2015.

[5] B. E. Kanguzhin, and N. E. Tokmagambetov, "Resolvents of well–posed problems for finite–rank perturbations of the polyharmonic operator in a punctured domain", *Siberian Mathematical Journal*, Vol. 57, No. 2, pp. 265–273, 2016.

## On first eigenvalue of Laplace operator

Baltabek KANGUZHIN [1], Dostilek DAUITBEK [2]

[1] *Institute of Mathematics and Mathematical Modeling, Kazakhstan*
[1] *Al-Farabi Kazakh National University, Kazakhstan*
*E-mail: kanbalta@mail.ru*
[2] *Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: dauitbek@math.kz*

**Abstract:** We consider a self-adjoint differential operator in a Hilbert space. Then the domain of the operator is changed by the perturbation of the boundary conditions so that a given neighborhood "there is no eigenvalues on neighborhood of zero" from the points of the spectrum of the perturbed operator. For the Sturm-Liouville operator on the segment and the Laplace operator on the square such a possibility is achieved through integral perturbations of boundary conditions. These statements are given with full proves and with possible extension. The G.G. Islamov's paper [1] solves the following problem. Suppose that the closed operator $A$ of the Banach space $X$ has eigenvalues in a "prohibited" area of $\Omega$ the complex plane. The requirement is for a $K$ finite-dimensional $V = A - K$ operator, wherein the operator will not have points in the spectrum in $\Omega$ area. Our main result is following theorem:

**Theorem 1.** If the parameter $\alpha$ chosen so, that inequality holds

$$(\lambda_2 - \lambda_1) \le \alpha \int_0^{\pi} \frac{\partial v_1(x_1, \pi)}{\partial x_2} dx_1, \tag{1}$$

the eigenvalues $\{\eta_k\}_{k=1}^{\infty}$ of the operator $B_1$ defined by the formula

$$\eta_k = \lambda_k \text{ when } k \ge 2,$$

and $\eta_1$ is the only real root of the equation.

**Keywords:**Laplace operator, eigenfunction, eigenvalue.

**2010 Mathematics Subject Classification:** 35J05, 35J08, 35J25, 35P05

**References:**

[1] G.G. Islamov "Eksperemental'nye vozmuchenia zamknutyh operatorov", *Izv. vuzov Math.*, 33, No.1, pp. 35-41 (1989).

# Estimates of the eigenvalues of operator arising in swelling pressure model

Baltabek KANGUZHIN [1], Lyailya ZHAPSARBAYEVA [2]

[1] *Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan, E-mail: kanbalta@mail.ru*

[2] *Al-Farabi Kazakh National University, Almaty, Kazakhstan, E-mail: leylazhapsarbaeva@rambler.ru*

**Abstract:** Swelling pressures from materials confined by structures can cause structural deformations and instability. Due to the complexity of interactions between expansive solid and solid-liquid equilibrium, the forces exerting on retaining structures from swelling are highly nonlinear. In 1994, Mesri and etc. [1], also [2], and [3], developed a simplistic equation for swelling pressure as a function of mobilized volume strain which can be used to show ( [4]) that the pressure $p$ acting on a wall can be modeled by $p(x) = p_{sl}e^{-\alpha v(x)}$, where $p_{sl}$ is the swelling pressure against an unyielding wall, $\alpha$ a constant depending upon the solid-liquid equilibrium, and $v(x)$ the deflection of the wall modeled as a cantilever beam at location $x$ along the beam from the fixed end. We consider the initial/boundary value problem of an Euler-Bernoulli elastic beam subject to the swelling pressure with one end clamped and another end free.

In this paper two sided estimates of eigenvalues of the operator corresponding to some nonlinear equation, which is the model of expansive clay are established.

**Keywords:** Euler-elastic beam equation, Swelling Pressure Model, Estimates of the eigenvalues

**2010 Mathematics Subject Classification:** 31B20, 35P20

**References:**

[1] Mesri, G., Pakbaz, M. C., Cepeda-Diaz, A. F., "Meaning, measurement and field application of swelling pressure of clay shales", *Géotechnique*, Vol. 44, No.1, pp.129–145, 1994.

[2] Grob, H., "Schwelldrucim Belchentunnel", *Proceedings of the International Symposium on Underground Openings, Lucerne, Switzerland*, pp. 99–119, 1972.

[3] Gysel, M., "Design of Tunnels in Swelling Rock", *Rock Mechanics and Rock Engineering*, Vol.20, No.4, pp. 219–242, 1987.

[4] Endley, S., "A Commentary on Active Pressure in Clays by Classical Method and Effect of Swelling", *Louisiana Civil Engineering Conference and Show*, 2011.

## The resolvent equation of nonlinear Fredholm integral equations

Akylbek KERIMBEKOV

*Department of Applied Mathematics and Informatics, Kyrgyz-Russian Slavic university, Bishkek, Kyrgyzstan*
*E-mail: akl7@rambler.ru*

**Abstract:** It is considered the new method solving of nonlinear Fredholm integral equation. It is found the sufficient condition for the existence of a solution of the nonlinear integral equation and it is proved uniqueness of the found solution.

In the article we consider the nonlinear Fredholm integral equation [1, 2]

$$\varphi(x) = \lambda \int_a^b K\left(x, t, \varphi(t)\right)\, dt + f(x), \tag{1}$$

where $f(x)$ is known continuous function defined on the $[a, b]$; Kernel $K\left(x, t, \varphi(t)\right)$ nonlinearly depends on the unknown function $\varphi(x)$ and continuous on set of arguments $x \in [a, b]$, $t \in [a, b]$,$\varphi \in (c, h)$, and it has a continuous derivative with respect to functional variable $\varphi$; $\lambda$ is a parameter.

By the method described in [3, 4] we have established that if the kernel $K\left(x, t, \varphi(t)\right)$ satisfies

$$0 \equiv \int_a^b \int_a^b K'_{\varphi}\left[x, t, \psi(t, \lambda)\right] u(t)\, dt\, dx, \tag{2}$$

the equation (1) admits a solution of the form

$$\varphi(x) = \varphi_0(x) + \lambda u(x), \tag{3}$$

where $\varphi_0(x)$, $u(x)$ , $\psi(t, \lambda)$ are known functions . This method is the most effective in the case where identity (2) can be represented in the form

$$0 \equiv A(x) \int_a^b G\left[t, \psi(t, \lambda)\right] \bar{u}(t)\, dt. \tag{4}$$

In this case, if the parameter $\lambda$ is *the resolvent number*, i.e. is the root of *the resolvent algebraic equation*

$$B(\lambda) = \int_a^b G\left[t, \psi(t, \lambda)\right] \bar{u}(t)\, dt = 0, \tag{5}$$

correlation (2) holds for any $x \in [a, b]$.

Substituting the found value for resolvent number $\lambda_1, ..., \lambda_p,\ p \leq \infty$ into (3) we obtain the solution of equation (1). We note that if $\lambda$ is not a characteristic number of kernels $K'_{\varphi}\left[x, t, \psi(t, \lambda)\right]$, the solution is unique.

**Keywords:** nonlinear integral equation, uniqueness of solution, characteristic number, orthogonality

**2010 Mathematics Subject Classification:** 45B05

**References:**

[1] Smirnov, N.S. *Vvedenie v teoriyu nelineynih integralnih uravneniy.* M.: ONTI, 1936. (in Russian)

[2] Krasnov, M. L. *Integralnie uravneniya.* M.: Nauka, 1975. (in Russian)

[3] Kerimbekov, A. "About the one method of solving nonlinear Fredgolm integral equations", *Abstract. V Congress of the Turkic World mathematicians, Kyrgyzstan, "Issyk-Kul Aurora", 5-7 june* , P.84, 2014.
[4] Kerimbekov, A. "Metod resheniya nelineynih integralnih uravneniy Fredgolma", *Vestnik KRSU* , Vol.16, No.1, pp. 23-26, 2016. (in Russian)

## On Spectral Properties of Perturbations of the Operator $-u''(-x)$ with Initial Data

Leonid KRITSKOV [1], Abdizhahan SARSENBI [2,3]

[1] *Moscow State University named after M.V.Lomonosov, Moscow, Russia*
*E-mail: kritskov@cs.msu.su*
[2] *South-Kazakhstan State University, Shymkent, Kazakhstan*
[3] *Institute for Mathematics and Mathematical Modeling, Almaty, Kazakhstan*
*E-mail: abzhahan@mail.ru*

**Abstract:** In the recent years there appears a rising interest to the problems that contain involution in the main terms of its differential operations [1–3].

The peculiarity of the initial value problem

$$L_0 u = -u''(-x),\ -1 < x < 1, \quad u(-1) = u'(-1) = 0,$$

for the simplest second-order differential operation with involution (reflection) is the non-emptiness of its spectrum. As the related spectral expansion could be constructed explicitly, it is interesting to study the spectral expansion for the perturbed operator $L = L_0 + Q$ where $Qu = q(x)u(x)$ is the multiplication on a square integrable complex-valued potential $q(x)$.

Let $S_m(x, f)$ and $\sigma_m(x, f)$ be $m$-th partial sums of the spectral expansions for $L_0$ and $L$ respectively. Using asymptotics of the Green's function for $L$, the equiconvergence property is obtained, i.e.

$$S_m(x, f) - \sigma_m(x, f) = o(1), \quad m \to \infty,$$

uniformly for $x \in [-1, 1]$ and any $f(x) \in L_1(-1, 1)$. It directly provides the claim that $\sigma_m(x, f)$ converges to $f(x)$ in the $L_2$-metric for any $f(x) \in L_2(-1, 1)$, or, in other words, the root functions of $L$ constitute basis.

The research is supported by the grant 0971/GF4 and by the target program 0085/PTSF-14 from the Ministry for Education and Science of the Republic of Kazakhstan.

**Keywords:** ODE with involution, spectral expansion, equiconvergence

**2010 Mathematics Subject Classification:** 34K08, 34L10

**References:**

[1] Kurdyumov, V.P., Khromov, A.P., "Riesz bases formed by root functions of a functional-differential equation with a reflection operator", *Differ. Equations*, **44** (2008), 2, 203–212.
[2] Cabada, A., Tojo, F.A.F., "Solutions and Green's function of the first order linear equation with reflection and initial conditions", *Boundary Value Problems*, **2014**:99 (2014).
[3] Kritskov, L.V., Sarsenbi, A.M., "Basicity in $L_p$ of root functions for differential equations with involution", *Electr. J. Diff. Equ.*, 2015. **2015**:278 (2015).

## Well-posed problems for the Laplace operator in the multiply connected domains

Daulet NURAKHMETOV [1,2]

[1] *Department of Information System, S. Seifullin Kazakh Agrotechnical University, Astana, Kazakhstan*
*E-mail: dauletkaznu@gmail.com*
[2] *Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan*
*E-mail: nurakhmetov@math.kz*

**Abstract:** We give a complete description of well-posed solvable boundary-value problems for the Laplace operator in the multiply connected domains.

In [1] are studied the spectral properties of the Schrödinger operator with point-like interactions. In [2] are given of well-posed solvable boundary-value problems for the Laplace operators in the disk and in the punctured disk. The practical application of these kinds of operators can be found in [3], [4]. In [5] are obtained an analog of the Green formula and a class of self-adjoint extensions of the Laplacian. In [6] are studied of the first regularized of the Laplace operator in a plane bounded domain.

This report is a continuation of the paper [2].

**Keywords:** Laplace operator, interior boundary problem, Green function.

**2010 Mathematics Subject Classification:** 35A2

**Acknowledgments:** This publication is supported by the target program 0085/PTSF-14 from the Ministry of Science and Education of the Republic of Kazakhstan.

**References:**

[1] Neiman-Zade M.I., Savchuk A.M., "Schrödinger Operator with Singular Potentials", *Tr. Mat. Inst.Steklova*, Vol.236, pp. 262-271, 2002.
[2] Kanguzhin B.E., Aniyarov A.A., "Well-Posed Problems for the Laplace Operator in a Punctured Disk", *Mathematical Notes*, Vol.89, No.6, pp. 819-829, 2011.
[3] Albeverio S., Gesztesy F., Hoegh-Krohn, R., Holden, H., Solvabel models in quantum mechanics, New Yourk: Springer, 1988 (Second edition: AMS, 2005).
[4] Goloshchapova N.I., Zastavnyi V.P., Malamud M.M., "Positive definite functions and spectral properties of the Schrödinger operator with point interactions", *Mathematical Notes*, Vol.90, No.1, pp. 149-154, 2011.
[5] Kanguzhin B.E., Nurakhmetov D.B., Tokmagambetov N.E., "Laplace Operator with $\delta$-Like Potentials", *Russian Mathematics*, Vol.58, No.2, pp. 6-12, 2014.
[6] Kanguzhin B.E., Tokmagambetov N.E.,"A Regularized Trace Formula for a Well-Perturbed Laplace Operator", *Doklady Mathematics*, Vol.91, No.1, pp. 1-4, 2015.
[7] Bitsadze A.V., Equations of Mathematical Physics, Moscow: Nauka, 1976 (in Russian). (Second edition: AMS, 2005).
[8] Kalmenov T.Sh., Koshanov B.D., Nemchenko M.Yu., "Green function representation in the Dirichlet problem for polyharmonic equations in a ball", *Doklady Mathematics*, Vol.78, No.1, pp. 528-530, 2008.

## Very weak solutions to wave equations

Michael RUZHANSKY

*Imperial College London, UK*
*E-mail: m.ruzhansky@imperial.ac.uk*

**Abstract:** In this talk we discuss a notion of a very weak solution to hyperbolic equations with irregular coefficients which was introduced for the case of the wave equation in [1]. Using methods of nonharmonic analysis developed in [2] this allows one to prove existence and uniqueness for Cauchy problems of the type

$$u''(t) - a(t)Au(t) = f(t), \quad u(0) = u_0, \quad u'(0) = u_1,$$

where $A$ is a densely defined positive operator on some Hilbert space, and $a(t)$ is a non-negative distribution. Examples will be given for the classical wave equation (based on joint work with Claudia Garetto), and for the wave equation for the Landau Hamiltonian describing the movement of the particle in irregular electromagnetic field (based on joint work with Niyaz Tokmagambetov).

**References:**

[1] Garetto C., Ruzhansky M., "Hyperbolic second order equations with non-regular time dependent coefficients", *Arch. Ration. Mech. Anal.*, Vol. 217, pp.113–154, 2015.

[2] Ruzhansky M., Tokmagambetov N., "Nonharmonic analysis of boundary value problems", *Int. Math. Res. Notices*, Vol. 2016, No. 12, pp.3548–3615, 2016.

---

## An embedding theorem of Sobolev type spaces on stratified Lie groups

Bolys SABITBEK [1], Durvudkhan SURAGAN [2]

[1] *Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: b.sabitbek@math.kz*
[2] *Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: suragan@math.kz*

**Abstract:** In this talk, we present a version of horizontal weighted Hardy-Rellich type inequality on stratified Lie groups and study some of their consequences. Our results reflect on many results previously known in special cases. Moreover, new Sobalev type spaces are defined on stratified Lie groups and proved an embedding theorem for these functional spaces.

**Keywords:** Stratified Lie groups, Hardy-Rellich type inequality, Sobolev space

**2010 Mathematics Subject Classification:** 94B05, 94B15

## An isoperimetric inequality for heat potential and heat equation

Makhmud A. SADYBEKOV [1], Aidyn A. KASSYMOV [2]

*E-mail: sadybekov@math.kz*[1]*, kassymov@math.kz*[2]
*Institute of Mathematics and Mathematical Modeling, Kazakhstan*

**Abstract:** In this paper we prove that the ball is a minimizer of the first characteristic number of heat potential and heat operator among all Euclidean domains of a given measure.

Historically, for the first time of scientific literature, in Rayleigh's famous book "The Theory of Sound" (first published in 1877), by using some explicit computation and physical interpretations, he stated that a disk minimizes (among all domains of same area) the first eigenvalue of the Dirichlet Laplacian and the norm of an inverse operator is maximized in a disk among all Euclidean domains of a given area. The proof of this conjecture was obtained after about 50 years later, simultaneously (and independently) by G. Faber and E. Krahn. Nowadays, the Rayleigh-Faber-Krahn inequality has been extended to many other operators. In this paper we prove the Rayleigh-Faber-Krahn theorem for the heat potential operator $H$, i.e. it is proved that the ball is a minimizer of the first characteristic number of the integral operator $H$ among all domains of a given measure in $R^d$. The main reason why the result is useful, beyond the intrinsic interest of geometric extremum problems, is that it gives a priori bound for operator norm of the heat potential on arbitrary cylindric domains.

**Keywords:** heat potential, heat equation, Rayleigh-Faber-Krahn inequalty, Schatten p-norm, isoperimetric inequality.

**2010 Mathematics Subject Classification:** 35K10, 31B35, 47G10.

## On some spectral inequalities for a nonlocal elliptic problem

Makhmud A. SADYBEKOV, Berikbol T. TOREBEK

*Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan*
*E-mail: sadybekov@math.kz, torebek@math.kz*

**Abstract:** Let $\Omega \subset R^2$ be a bounded domain, symmetric with respect to the origin and with a smooth boundary $\partial\Omega$. This symmetry means that alongside with a point $(x_1, x_2)$ her "opposite" point $x* = (-x_1, -x_2)$ also belongs to the domain. Let us denote $\partial\Omega_+ = \partial\Omega \cap \{x_1 \geq 0\}$, $\partial\Omega_- = \partial\Omega \cap \{x_1 < 0\}$.

In this paper we consider the following problem:

$$-\Delta u\left(x\right) = f\left(x\right), x \in \Omega,$$

in the domain $\Omega$, and satisfying the following boundary conditions

$$u\left(x\right) = -u\left(x*\right), \frac{\partial u\left(x\right)}{\partial n_x} = \frac{\partial u\left(x*\right)}{\partial n_x}, x \in \partial\Omega_+.$$

Here $n_x$ is a derivative in the direction of an outer normal to $\partial\Omega$.

Investigated problem is an analogue of the classical periodic boundary value problems in the case of non-rectangular region. Note that the problem P in the case of a circle was first formulated and investigated in [1].

We prove self-adjointness of the problem and show a method of constructing eigenfunctions.

We obtain an analogue of the Rayleigh type inequality and some spectral inequality for the first eigenvalue of the nonlocal problem.

**Keywords:** Laplace operator, nonlocal problem, eigenvalue, eigenfunctions, Rayleigh type inequality

**2010 Mathematics Subject Classification:** 35P15, 35J05, 49R50

**References:**

[1] M. .A. Sadybekov, and B. K. Turmetov, *Differential Equations*. **50**:2, 268–273 (2014).

# Spectral properties of a Laplace operator with Samarskii–Ionkin type boundary conditions in a disk

Makhmud SADYBEKOV [1], Abdizhahan SARSENBI [1,2],
Aizhan TENGAYEVA [1,3]

[1] *Institute of Mathematics and Mathematical Modelling, Kazakhstan*
*E-mail: sadybekov@math.kz, sarsenbi@math.kz*
[2] *Auezov South Kazakhstan State University, Kazakhstan*
[3] *Kazakh National Agrarian University, Kazakhstan*
*E-mail: aijan0973@mail.ru*

**Abstract:** In this paper we consider a spectral problem for a model differential operator of the second order with involution. The operator is given by a differential expression

$$Lu = -u''(-x), \ -1 < x < 1 \tag{1}$$

and boundary conditions of the general form:

$$\alpha_j u'(-1) + \beta_j u'(1) + \alpha_{j1} u(-1) + \beta_{j1} u(1) = 0, \ \ j = 1, 2. \tag{2}$$

Coefficients of the boundary condition are numbers, which can be complex.

A criterion of the basis property of systems of eigenfunctions of the operator is set in the sense of coefficients of the boundary conditions. The interest to studying the basis properties of operators of the form (1), (2) has arose relatively recently. In works of T.Sh. Kal'menov and his disciples the *self-adjointness* property of the Cauchy operator:

$$L_K u = -u''(-x), \ -1 < x < 1, \quad u'(-1) = 0, \ \ u(-1) = 0 \tag{3}$$

is successfully applied in solving initial-boundary value problems for a heat equation with inverse time direction, in finding a criterion of well-posedness of the Cauchy problem for the Poisson operator, in formulating an asymptotic solution to problems with a small parameter.

**Keywords:** generalized spectral problem, involution, eigenfunctions, eigenvalues, basis property

**2010 Mathematics Subject Classification:** 34L10, 34L15

# Spectral properties of a Laplace operator with Samarskii–Ionkin type boundary conditions in a disk

Makhmud SADYBEKOV [1], Nurgissa YESSIRKEGENOV [1,2]

[1] *Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan*
[2] *Department of Mathematics, Imperial College London, London, United Kingdom*
*E-mail: sadybekov@math.kz, n.yessirkegenov15@imperial.ac.uk*

**Abstract:** One of the most important problems in the theory of harmonic functions are Dirichlet and Neumann boundary value problems. In the one-dimensional case, or when considering the problem in a multidimensional parallelepiped, the main problems include also the periodic boundary value problems.

In this paper we consider spectral properties of a Laplace operator with boundary conditions of Samarskii-Ionkin type in a disk and prove the completeness of eigenfunctions.

Let $\Omega = \{z = (x, y) = x + iy \in C : |z| < 1\}$ be a unit disk, $r = |z|, \varphi = \arctan(y/x)$, $\Omega^+ = \Omega \cap \{y > 0\}$, and $\Omega^- = \Omega \cap \{y < 0\}$. We consider the spectral problem corresponding to the Laplace operator

$$-\Delta u(z) = \lambda u(z), |z| < 1$$

with local boundary conditions

$$u(1, \varphi) = 0, \ \ 0 \le \varphi \le \pi,$$

and with one of the nonlocal boundary conditions

$$\frac{\partial u}{\partial r}(1, \varphi) - \frac{\partial u}{\partial r}(1, 2\pi - \varphi) = 0, \ \ 0 \le \varphi \le \pi,$$

or

$$\frac{\partial u}{\partial r}(1, \varphi) + \frac{\partial u}{\partial r}(1, 2\pi - \varphi) = 0, \ \ 0 \le \varphi \le \pi.$$

We note that unlike the one-dimensional case the system of root functions of the problems consists only of eigenfunctions.

**Keywords:** Poisson equation, Samarskii–Ionkin type problem, eigenfunctions, eigenvalues

**2010 Mathematics Subject Classification:** 35J05, 35J25, 35P10

## The theorem on the basis property of eigenfunctions of second order differential operators with involution

Abdizhahan SARSENBI [1]

[1] *Department of Mathematical methods and modeling, M. Auezov South Kazakhstan State University, Shymkent, Kazakhstan*
*E-mail: abzhahan@gmail.com*

**Abstract:** In the works [1-2] was shown that the eigenfunction system

$$\left\{\cos k\pi x,\ \sin(k+\frac{1}{2})\pi x,\ \ k=0,1,2,...\right\}$$

of the spectral problem

$$-u''(-x)=\lambda u(x),\ \ -1<x<1, \tag{1}$$

$$u'(-1)=0,\ u'(1)=0. \tag{2}$$

forms a Riesz basis of the space $L_2(-1,1)$. In this connection, naturally arises the question, will the eigenfunctions system of the spectral problem

$$-u''(-x)+q(x)u(x)=\lambda u(x),\ \ -1<x<1, \tag{3}$$

with boundary conditions (2) be a basis in space $L_2(-1,1)$? Let $\sigma_m(f)$ partial sum of partial sum of eigenfunctions expansion of the spectral problem (1), (2), and $S_m(f)$ partial sum of eigenfunctions expansion of the spectral problem (3), (2). Sequence $S_m(f)$ call equconvergent with sequence $\sigma_m(f)$ on the interval $-1\le x\le 1$, if $S_m-\sigma_m\to 0$ uniformly converges on this interval at $m\to\infty$. It is possible to show the validity of the theorem.

**Theorem 1.** *Eigenfunction system of the spectral problem (3), (2) with continuous coefficient $q(x)$ forms basis of the space $L_2(-1,1)$.*

**Keywords:** differential equation with involution, Green's function, eigenfunction expansions, basis

**2010 Mathematics Subject Classification:** 34K08, 34L10, 46B15

**References:**

[1] Sarsenbi, A.M., "Unconditional bases related to a nonclassical second-order differential operator, *Differential Equations*, Vol. 46, No.4, pp.509–5014, 2014.
[2] Sadybekov, M.A., Sarsenbi, A.M., "Criterion for the basis property of the eigenfunction system of a multiple differentiation operator with an involution, *Differential Equations*, Vol. 48, No.8, pp.1112–1118, 2012.

# Solution of a singularly perturbed Cauchy problem for linear systems of ordinary differential equations by the method of spectral decomposition

Amir SHALDANBAYEV[1], Manat SHOMANBAYEVA[2], Asylzat KOPZHASSAROVA[2]

[1] *Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan*
*E-mail: shaldanbaev51@mail.ru*
[2] *M. Auezov South Kazakhstan State University, Kazakhstan*
*E-mail: mtshomanbaeva@mail.ru, asyl_k@mail.ru*

**Abstract:** In this paper a completely new method, which is essentially an operational method [1] and is based on the spectral theory of equations with deviating argument [2-7], is proposed. The method consists in the fact that the spectral decompositions of Cauchy problem solutions are constructed by the system of eigenfunctions of the self-adjoint differential operators with deviating argument. It turned out that so-called fundamental solutions of a system of linear differential equations and equations of higher order play a special role. At the asymptotic analysis of solutions of singularly perturbed problems the so-called regulatory functions of S.A. Lomov occupy the important place. For the first time it is proved that the fundamental solutions of linear differential equations act as the regularizing functions of S.A. Lomov.

**Keywords:** Spectrum, spectral decomposition, deviating argument, singular perturbation

**2010 Mathematics Subject Classification:** 34D15, 34K08

**References:**

[1] G. I. Shishkin, J. J. H. Miller, and E. O'Riordan, Fitted Numerical Methods for Singular Perturbation Problems: Error Estimates in the Maximum Norm for Linear Problems in One and Two Dimensions, World Scientific, River Edge, NJ, USA, 1996.

[2] S. M. Reed, and B. Simon, *Functional Analysis*, Academic Press, 1981.

[3] T. Sh. Kal'menov, S. T. Akhmetova, and A. Sh. Shaldanbaev, *Mat. Zh. Almaty* **4**, 41–48 (2004), (in Russian).

[4] T. Sh. Kal'menov, and A. Sh. Shaldanbaev, *Journal of Inverse and Ill-Posed Problems* **18**, 352–369 (2010).

[5] A. A. Kopzhassarova, A. L. Lukashov, and A.M. Sarsenbi, *Abstract and Applied Analysis*, vol.2012, Article ID 590781, 5 pages, 2012.

[6] A.Kopzhassarova and A.Sarsenbi, *Abstract and Applied Analysis*, vol. 2012, Article ID 576843, 6 pages, 2012.

[7] I.Orazov, A.Shaldanbayev, and M.Shomanbayeva, *Abstract and Applied Analysis* vol. 2013, Article ID 128363,(2013)

## Differential operators with singular coefficients

Andrei SHKALIKOV

*Lomonosov Moscow State University, Russia*
*E-mail: shkalikov@mi.ras.ru; ashkaliko@yandex.ru*

**Abstract:** Classical theory of symmetric ordinary differential operators allows to work with differential expressions of the form

$$\ell_{2n}[y] := \sum_{k=0}^{n}(-1)^{n-k}(p_k\, y^{(n-k)})^{(n-k)} + \\ + i\sum_{k=0}^{n-1}(-1)^{n-k-1}\{(q_k\, y^{(n-k-1)})^{(n-k)} + (q_k\, y^{(n-k)})^{(n-k-1)}\}, \tag{4}$$

provided that the coefficients $p_k$, $q_k$ and $(p_0)^{-1}$ are locally integrable functions. This theory was developed in the works of D. Shin, N. Glazman, A. Zettle, N. Everitt et al.

Our goal is to extend the frames of the classical theory and to define operators associated with non-symmetric differential expressions

$$\tau(y) = \sum_{k,s=0}^{n}\left(r_{ks}y^{(n-k)}\right)^{(n-s)}, \tag{5}$$

whose coefficients $r_{ks}$ are distributions of finite order singularity (depending on the indices $k, s$).

We shall discuss several approaches to this problem. The most important one is based on the so-called regularization procedure. This procedure provides several advantages. In particular, one can write out useful asymptotic formulae for the solutions of the equation

$$\ell_{2n}[y] - \lambda y = 0.$$

We shall also discuss similar problems for partial differential operators.

The talk is based on the joint papers with prof. K.A. Mirzoev and A.M. Savchuk.

**Keywords:** differential operators with distribution coefficients, differential operators with singular coefficients, regularization method

**2010 Mathematics Subject Classification:** 34E15

# Existence of eigenvalues of problem with shift for an equation of parabolic-hyperbolic type

Aizhan TENGAYEVA [1], Gulnar DILDABEK [2]

[1] *Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: aijan0973@mail.ru*
[2] *Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: dildabek.g@gmail.com*

**Abstract:** In the paper a spectral problem for an operator of parabolic-hyperbolic type of I kind with non-classical boundary conditions is considered. The problem is considered in a standard domain. The parabolic part of the space is a rectangle. And the hyperbolic part of the space coincides with a characteristic triangle. We consider a problem with the local boundary condition in the domain of parabolicity and with the boundary condition with displacement in the domain of hyperbolicity.

Let $\Omega \in R^2$ be a finite domain bounded for $y > 0$ by the segments $AA_0$, $A_0B_0$, $B_0B$, $A = (0,0)$, $A_0 = (0,1)$, $B_0 = (1,1)$, $B = (1,0)$, and for $y < 0$ by the characteristics $AC:\ x + y = 0$ and $BC:\ x - y = 1$ of an equation of the mixed parabolic-hyperbolic type

$$Lu = \left\{ \begin{array}{c} u_x - u_{yy}, y > 0 \\ u_{xx} - u_{yy}, y < 0 \end{array} \right\} = f(x,y). \tag{1}$$

Problem $S$. *Find a solution to Eq.* (1) *satisfying boundary conditions*

$$u|_{AA_0 \cup A_0B_0} = 0, \tag{2}$$

$$\alpha u\left(\theta_0\left(t\right)\right) = \beta u\left(\theta_1\left(t\right)\right), \quad 0 \le t \le 1, \tag{3}$$

*where* $\theta_0\left(t\right) = \left(\frac{t}{2}, -\frac{t}{2}\right)$, $\theta_1\left(t\right) = \left(\frac{t+1}{2}, \frac{t-1}{2}\right)$.

We prove the strong solvability of the considered problem. The main aim of the paper is the research of spectral properties of the problem. The existence of eigenvalues of the problem is proved.

**Keywords:** spectral problem, equation of parabolic-hyperbolic type, boundary condition with shift

**2010 Mathematics Subject Classification:** 35M10, 35M12, 35P

## Duality property of the noncommutative $\ell_\infty$ and $\ell_1$ valued symmetric Hardy spaces

Kanat TULENOV [1,2]

[1] *Institute of Mathematics and Mathematical Modeling, Kazakhstan.*
*E-mail: tulenov@math.kz*
[2] *Al-Farabi Kazakh National University, Kazakhstan*
*E-mail: kanat.tulenov@gmail.com*

**Abstract:** In this paper we consider the noncommutative $H_E(\mathcal{A};\ell_\infty)$ and $H_E(\mathcal{A};\ell_1)$ spaces and obtain some result on duality for these spaces, i.e we obtain the following result:

**Theorem.** Let $E$ be an $r$-convex symmetric Banach function space on $[0;1]$ for some $0<r<\infty$ and $E$ do not contain $c_0$ or separable. Then

$$(i)\ \ (H_E(\mathcal{A};\ell_1))^*=L_{E^\times}(\mathcal{M};\ell_\infty)/J(H^0_{E^\times}(\mathcal{A};\ell_\infty))$$

isometrically via the following duality bracket

$$((x_n),(y_n))=\sum_{n=1}^{\infty}\tau(y_n^*x_n)$$

for $x\in H_E(\mathcal{A};\ell_1)$ and $y\in H_{E^\times}(\mathcal{A};\ell_\infty)$, where $J(H^0_{E^\times}(\mathcal{A};\ell_\infty))=\{x^*:\ x\in H^0_{E^\times}(\mathcal{A};\ell_\infty)\}$.

$$(ii)\ \ (L_E(\mathcal{M};\ell_1)/J(H^0_p(\mathcal{A};\ell_1)))^*=H_{E^\times}(\mathcal{A};\ell_\infty)$$

isometrically via the following duality bracket

$$((x_n),(y_n))=\sum_{n=1}^{\infty}\tau(y_n^*x_n)$$

for $x\in H_E(\mathcal{A};\ell_1)$ and $y\in H_{E^\times}(\mathcal{A};\ell_\infty)$, where $J(H^0_{E^\times}(\mathcal{A};\ell_1))=\{x^*:\ x\in H^0_{E^\times}(\mathcal{A};\ell_1)\}$.

This theorem is analogue of Theorem 5 in [1].

**Keywords:** von Neumann algebra, subdiagonal algebras, noncommutative vector valued symmetric Hardy spaces, duality.

**2010 Mathematics Subject Classification:** 46L51, 46L52.

**References:**

[1] T. N. Bekjan, K. Tulenov, and D. Dauitbek, The Noncommutative $H_p^{(r,s)}(\mathcal{A};l_\infty)$ and $H_p(\mathcal{A};l_1)$ spaces. *Positivity*, **19** 877–891 (2015).

# Lateral boundary conditions for the Klein-Gordon-Fock equation

Kanat TULENOV [1], Dostilek DAUITBEK [2]

[1] *Institute of Mathematics and Mathematical Modeling*
*E-mail: tulenov@math.kz*
[2] *Institute of Mathematics and Mathematical Modeling, Kazakhstan*
*E-mail: dauitbek@math.kz*

**Abstract:** In this paper we consider an initial-boundary value problem for the Klein-Gordon-Fock equation. In [**?**], Authors considered one dimentional potential

$$u(x,t)=\int_{\Omega}(-\frac{1}{2}|x-y|)f(y)dy \tag{1}$$

in $\Omega=(0,1)$ and the equation

$$-u^{"}(x)=f(x),\ \ x\in\Omega, \tag{2}$$

and they showed that if they are solved the equation (2) with the boundary conditions

$$u^{'}(0)+u^{'}(1)=0,\ \ -u^{'}(1)+u(0)+u(1)=0. \tag{3}$$

then they were finded an unique solution of that boundary value problem in the form (1).

This simple method finds equivalent boundary value problems of ODE for one dimensional potential integrals. However this task becomes tedious for PDE, and in works [1, 2] are obtained boundary conditions of the volume potentials for elliptic and hyperbolic equations and showed some their applications. In particular, in [2], by using a new non-local boundary value problem, which is equivalent to the Newton potential, Authors found explicitly all eigenvalues and eigenfunctions of the Newton potential in the 2-disk and the 3-ball. The aim of this work is to find lateral boundary conditions for the Klein-Gordon-Fock equation as shown in [**?**].

**Keywords:** Potential, fundamental solution, Klein-Gordon-Fock equation

**2010 Mathematics Subject Classification:** 35J05, 35J08, 35J25, 35P05

**References:**

[1] Kal'menov T.Sh. , Suragan D., "Initial-boundary value problem for the wave equation", *Electron. J. Diff. Equ.*, Vol. 48, pp. 1-6, 2014.
[2] Kal'menov T.Sh. , Suragan D., "To spectral problems for the volume potential", *Doklady Mathematics*, Vol. 80, No. 2, pp. 646-649, 2009.

## Euler-Hilbert-Sobolev spaces on homogeneous groups

Nurgissa YESSIRKEGENOV

*Institute of Mathematics and Mathematical Modeling, Kazakhstan and Department of Mathematics, Imperial College London, 180 Queen's Gate, SW7 2AZ London, United Kingdom*
*E-mail: n.yessirkegenov15@imperial.ac.uk*

**Abstract:** In this talk we define Euler-Hilbert-Sobolev spaces and obtain embedding results on homogeneous groups using Euler operators, which are homogeneous differential operators of order zero. Sharp remainder terms of $L^p$ and weighted Sobolev and Sobolev-Rellich inequalities on homogeneous groups are given. As consequences, we obtain analogues of the generalised classical Sobolev and Sobolev-Rellich inequalities. We also discuss applications of logarithmic Hardy inequalities to Sobolev-Lorentz-Zygmund spaces.

**Keywords:** Sobolev inequality, Hardy inequality, weighted Sobolev inequality, Rellich inequality, Euler-Hilbert-Sobolev space, Sobolev-Lorentz-Zygmund space, homogeneous Lie group

**2010 Mathematics Subject Classification:** 22E30, 43A80

# MS3

# DINAMICAL

# SYSTEMS

# On a dynamical system relating to the Ricci flow on Aloff-Wallach spaces

Nurlan ABIEV [1]

[1] *M. Kh. Dulaty Taraz State University, Kazakhstan, E-mail: abievn@mail.ru*

**Abstract:** The talk is devoted to the normalized Ricci flow equation

$$\dot{\mathbf{g}}(t) = -2\mathrm{Ric}_{\mathbf{g}} + 2n^{-1}\mathbf{g}(t)S_{\mathbf{g}}$$

on Aloff-Wallach spaces $\mathrm{SU}(3)/\mathrm{SO}(2)$, where $\mathbf{g}(t)$ means a 1-parameter family of Riemannian metrics, $\mathrm{Ric}_{\mathbf{g}}$ is the Ricci tensor and $S_{\mathbf{g}}$ is the scalar curvature of $\mathbf{g}$. On Aloff-Wallach spaces the normalized Ricci flow equation can be reduced to a system of four ODE's of the form

$$\dot{x_i} = -2\,x_i\left(\mathbf{r_i} - \frac{\mathbf{r_4} + 2(\mathbf{r_1} + \mathbf{r_2} + \mathbf{r_3})}{7}\right), \tag{4}$$

where $\mathbf{r_i}$, $i = 1, \ldots, 4$, are the principal Ricci curvatures (see [3,4] for explicit expressions of $\mathbf{r_i}$). The main result is contained in the following two Lemmas.

**Lemma 1.** *The system* (4) *has exactly two singular (equilibrium) points.*

**Lemma 2.** *The volume function* $V := x_1^2 x_2^2 x_3^2 x_4$ *is a first integral of the system (4).*

It should be noted that similar questions were studied in [1,2] in the case of generalized Wallach spaces.

**Acknowledgements.** The project was supported by Grant 1452/GF4 of MES of the Republic of Kazakhstan for 2015-2017.

**Keywords:** Ricci flow, Riemannian metric, Ricci curvature, Aloff-Wallach space, generalized Wallach space.

**2010 Mathematics Subject Classification:** 53C30, 53C44, 34C05, 37C10.

**References:**

[1] Abiev N. A., Arvanitoyeorgos A., Nikonorov Yu. G., Siasos P., "The dynamics of the Ricci flow on generalized Wallach spaces", *Differential Geometry and its Applications*. Vol. 35, Supplement, pp. 26–43, 2014.

[2] Abiev N. A., Nikonorov Yu. G., "The evolution of positively curved invariant Riemannian metrics on the Wallach spaces under the Ricci flow", *Annals of Global Analysis and Geometry*. Vol. 50, No.1, pp. 65–84, 2016.

[3] Nikonorov Yu. G., "Compact homogeneous Einstein 7-manifolds", *Geometriae Dedicata*, Vol. 109, No. 1, pp. 7–30, 2004.

[4] Wang M. Y., "Some examples of homogeneous Einstein manifolds in dimension seven", *Duke Mathematical Journal*. Vol. 49, pp. 23–28, 1982.

## Investigation on absolute stability of regulated systems in a simple critical case

Serikbai A. AISAGALIEV, A.M. AYAZBAYEVA

*Al-Farabi Kazakh National University, Almaty, Kazakhstan*
*E-mail: Serikbai.Aisagaliev@kaznu.kz, a_ayazbaeva@mail.ru*

**Abstract:** A new method for studying of absolute stability of the equilibrium position of nonlinear regulated systems in a simple critical case by evaluating improper integrals along the solutions of the system is proposed. Sectors are found, where the equilibrium position of the system is absolutely stable and the Aizerman problem has a positive solution. The effectiveness of the method is shown in the example.

**Keywords:** Absolute stability, non-singular transformation, properties of solutions, improper integrals, Aizerman problem, absolute stability sectors

**2010 Mathematics Subject Classification:** 02.30.Hq, 02.30.Yy

**References:**

[1] Aisagaliev S.A. "About determination of a region of absolute stability of forced movements in non-linear systems", *Izv. AS USSR.Technique cibernetics*, No. 5, pp. 38-48, 1969.

[2] Aisagaliev S.A. "About determination of a region of absolute stability control system with several nonlinear elements", *USSR. Automation and Remote Control*, No. 12, pp. 83-94, 1970.

[3] Aisagaliev S.A. "To the theory of absolute stability of regulated systems", *Differential equations*, V. 30, No. 5, pp. 748-757, 1994.

[4] Aisagaliev S.A. "Theory of regulated systems", – *Almaty: Kazakh University*, – 234 p., 2000.

[5] Aisagaliev S.A. "Stability theory of dynamical systems", – *Almaty: Kazakh University*, – 216 p., 2012.

[6] Aisagaliev S.A., Kalimoldayev M.N. "Certain problems of Synchronization theory", *Journal Inverse Ill Posed Problems*, 21, – pp. 159-175, 2013.

## Controllability of process described by system of linear integro-differential equations with resrtictions

Serikbai AISAGALIEV [1], Ilya SEVRYUGIN [2]

[1] *Al-Farabi Kazakh National University, Almaty, Kazakhstan*
*E-mail: Serikbai.Aisagaliev@kaznu.kz*
[2] *Al-Farabi Kazakh National University, Almaty, Kazakhstan*
*E-mail: Ilya.Sevrugin@gmail.com*

**Abstract:** Studied control process described by linear integro-differential equation $\dot{x} = A(t)x + B(t)u(t) + C(t)\int\limits_a^b K(t,\tau)w(\tau)d\tau + \mu(t)$, $t \in I = [t_0, t_1]$ with boundary, phase, integral restrictions and restrictions on the control value.

For this problem offered method of solution based on passing from solving of initial problem to solving of some class of Fredholm integral equation of the first kind. For initial problem obtained necessary and sufficient conditions of it's solvability. Offered numerical method for solving control problem by constructing minimizing sequences.

This work is a continuation of researches conducted in the works [1, 2]

**Keywords:** optimal control, linear integro-differential equations

**2010 Mathematics Subject Classification:** 49N05

**References:**

[1] Aisagaliev S.A., Sevryugin I.V. "Upravljaemost' i bystrodejstvie processa, opisyvaemogo linejnoj sistemoj obyknovennyh differencial'nyh uravnenij", *Matematicheskij zhurnal*, Vol. 13, pp.5–30, 2013.

[2] Aisagaliev S.A., "Konstruktivnaja teorija kraevyh zadach obyknovennyh differencial'nyh uravnenij", -*Kazak universiteti*, Almaty, 2015.

# Localization of an air target by means of GNSS-based multistatic radar

Daulet Sh. AKHMEDOV [1], Almat S. RASKALIYEV [2]

[1] *Institute of Space Technique and Technology, Almaty, Kazakhstan*
*E-mail: lacp@mail.ru*
[2] *Department of Mechanics, Al-Farabi Kazakh National University, Kazakhstan*
*E-mail: raskaliyev@mail.ru*

**Abstract:** The research of employing GNSS satellites as transmitters of opportunity for passive bistatic and multistatic radars to detect [1, 2], track [3] and localize [4] an aircraft has aroused some interest lately. A main benefit of utilizing GNSS signals for such application is because of its global coverage. Furthermore, the presence of several GNSS satellites at various positions provides an advantage for using this application, as such geometrical arrangements enable the radar to receive reflections of the target from different angles.

In this work, we offer a contribution to multistatic GNSS-based radar methods, which allows for positioning of air targets provided that pseudoranges of reflected signals have been preliminarily measured. This novel computational method consists from decomposition of non-linear function in a Taylor series and the Least Squares method. Employing pseudorange equitions based on reflected signals from multiple satellites should make it possible to determine a position estimate of the aircraft more accurately than by means of the bistatic radar.

**Keywords:** GNSS, multistatic radar, GPS, air target, reflected signals

**2010 Mathematics Subject Classification:** 65D15, 94A12

**References:**

[1] Behar, V., Kabakchiev, C., "Detectability of Air Targets using Bistatic Radar Based on GPS L5 Signals", *Proceedings of Intern. Radar Symp. (IRS) 2011*, Leipzig, pp.212–217, 2012.

[2] Glennon, E.P., Dempster, A.G., Rizos. C., "Feasibility of air target detection using GPS as a bistatic radar", *Journal of Global Positioning Systems*, Vol.5, No.1-2, pp.119–126, 2006.

[3] Lindgren, T., "Algorithm Development for Multistatic GNSS Radar Technology", Licantiate thesis, Lule University of Technology, iSSN: 1402-1757, 2007.

[4] Pui, C., Trinkle, M., "GPS Bistatic Radar for Target Detection and Estimation using Antenna Arrays", *Proceedings of IGNSS Society 2011*, Sydney, 2011.

## Gradient in the optimal control problem described by the differential equations

Hikmet AKHUNDOV [1], Mutallim MUTALLIMOV [1]

[1] *Institute of Applied Mathematics, Baku State University, Baku, Azerbaijan*
*E-mail: axund2005@rambler.ru, mutallim@mail.ru*

**Abstract:** A gradient is calculated in the optimal control problem with terminal functional described by the second order ordinary differential equation. Using the calculated gradient of the optimal control problem a problem of numerical solving of the initial problem is considered.

Consider the minimization of the functional

$$J(u) = |R(r_1, u) - z|^2 \tag{1}$$

subject to

$$-a\frac{d}{dr}\left(r^2\frac{dR}{dr}\right) + (Q(r) - Er^2)R(r) = u(r), \quad r_0 < r < r_1, \quad r_0 > 0 \tag{2}$$

$$R(r_0) = z_0, \ \dot{R}(r_1) = z_1. \tag{3}$$

where $R(\cdot) \in H'[r_0, r_1]$, $u(\cdot) \in L_2[r_0, r_1]$, $\tilde{H}^1[r_0, r_1] = \{\psi(\cdot) \in H^1[r_0, r_1] : \psi(r_0) = 0\}$.

The following theorem is proved.

**Theorem 1.** If there exists $\alpha > 0$ such that $Q(r) - Er^2 \geq \alpha$ for $r \in [r_0, r_1]$, where $Q(\cdot) \in C[r_0, r_1]$ then for any $h(\cdot) \in L_2[r_0, r_1]$ the adjoint to the initial one problem has the only generalized solution and

$$\|\Delta R(\cdot)\| \leq \|h\|_{L_2[r_0, r_1]}.$$

On the base of these results a numerical algorithm is developed for the solution of the considered problem.

**Keywords:** gradient, optimal control, differential equation.

**2010 Mathematics Subject Classification:** 31A25, 65L10, 65M32, 80A23

**References:**

[1] Mikhaylov, V.P., "Partial Differential Equations", Moscow, Nauka, 424p, 1985.
[2] Vasil'ev, F.P., "Numerical Methods for Solving Extremal Problems", (in Russian) Moscow, Nauka, 400p, 1981.

# Time frequency method of solving one boundary value problem for a hyperbolic system and its application to the oil extraction

F.A. ALIEV [1], N.A. ALIEV [1], A.P. GULIEV [1], K.G. GASIMOVA [2]

[1] *Institute of Applied Mathematics, Baku State University, Baku, Azerbaijan*
*E-mail: f_aliev@yahoo.com*

[2] *Azerbaijan State Pedagogical University, Baku, Azerbaijan*

**Abstract:** We consider the boundary value problem, where the motion of the object is described by the two-dimensional linear system of partial differential equations of hyperbolic type [1, 2], where a discontinuity is at a point within the interval that defines the phase coordinate $x$. Using the method of series and Laplace transformation in time $t$ (time-frequency method), an analytical solution is found for the determination of debit $Q(2l, t)$ and pressure $P(2l, t)$, which can be effective in the calculating of the coefficient of the hydraulic resistance in the lift at oil extraction by gas lift method [3], where $l-$ is the well depth. For the case, where the boundary functions are exponential form, the formulas for the $P(2l, t)$ and $Q(2l, t)$ depending only on $t$ are obtained. It is shown that at constant boundary functions, these formulas allow us to determine the coefficient of hydraulic resistance in the lift of the gas lift wells, which determines the change in the dynamics of pollution [4].

**Keywords:** hyperbolic equation, boundary problems, method of series, Laplace transformation, time-frequency method, gas lift, coefficient of hydraulic resistance

**2010 Mathematics Subject Classification:** 35L04, 35L20, 65N99

**References:**

[1] Aliev, F.A., Ilyasov, M.Kh., Nuriev, N.B., "Problems of modeling and optimal stabilization of the gas lift process", *Applied Mechanics*, Vol.46, No.6, pp.115–122, 2010.

[2] Aliev, F.A., Guliev, A.P., Ilyasov, M.Kh., "Method series to solving a boundary value problem for the system of hyperbolic equations, arising in the oil production", *Proceedings of IAM*, Vol.2, No.2, pp.113–136, 2013.

[3] Aliev, F.A., Ismailov, N.A., "Optimization problems with periodic boundary conditions and boundary control for lift wells", *Journal of Mathematical Sciences August*, Vol.208, No.5, pp.467–476, 2015, Translated from Neliniini Kolyvannya, Vol.17, No.2, 2014, pp.151-160, No.2, pp.216–227, 2014.

[4] Aliev, F.A., Ismailov, N.A., "Inverse problem to determine the hydraulic resistance coefficient in the gas lift process", *Appl. Comput. Math.*, Vol.12, No.3, pp.306–313, 2013.

## On Numerical Solutions of Telegraph Equations with the Dirichlet Boundary Condition

Allaberen ASHYRALYEV[1], Kadriye Tuba TURKCAN, Mehmet Emir KOKSAL[2]

[1] *Emeritus Professor, Turkey & Turkmenistan*
*E-mail: aallaberen@gmail.com*
[2] *Department of Mathematics, Ondokuz Mayis University, 55139 Atakum, Samsun, Turkey*
*E-mail: mekoksal@omu.edu.tr*

**Abstract:** In this study, the Cauchy problem for telegraph equations in a Hilbert space is considered. Stability estimates for the solution of this problem are presented. The third order of accuracy difference scheme is constructed for approximate solutions of the problem. Stability estimates for the solution of this difference scheme are established. As a test problem to support theoretical results, one-dimensional telegraph equation with the Dirichlet boundary condition is considered. Numerical solutions of this equation are obtained by first, second and third order of accuracy difference schemes..

**Keywords:** telegraph equations, numerical solutions, error analysis

**Acknowledgment:** This work is supported by the Scientific and Technological Research Council of Turkey - TUBITAK (Project no: 115E938) as in the scope of the second work packet.

# Optimal control problem for the degenerate phase field system of equations

Guzel BAYBULATOVA [1]

[1] *Department of Mathematics, Chelyabinsk State University, Chelyabinsk, Russia*
*E-mail: baybulatova_g_d@mail.ru*

**Abstract:**

In this work, the optimal control problem for the degenerate linear system of the phase field is considered

$$v(x,0) = \varphi(x), \quad x \in \Omega, \tag{1}$$

$$\theta \frac{\partial v}{\partial n}(x,t) + (1-\theta)v(x,t) = 0, \quad (x,t) \in \partial\Omega \times (0,T), \tag{2}$$

$$\theta \frac{\partial w}{\partial n}(x,t) + (1-\theta)w(x,t) = 0, \quad (x,t) \in \partial\Omega \times (0,T), \tag{3}$$

$$v_t(x,t) + lw_t(x,t) = k\Delta v(x,t) + u_1(x,t), \quad x \in \Omega \times (0,T), \tag{4}$$

$$\Delta w(x,t) + \alpha w(x,t) + \beta v(x,t) + u_2(x,t) = 0, \quad x \in \Omega \times (0,T), \tag{5}$$

$$\|u_1\|^2_{L_2(0,T;L_2(\Omega))} + \|u_2\|^2_{L_2(0,T;L_2(\Omega))} \le R^2, \tag{6}$$

$$J(v,w) = \frac{1}{2}\|v - \tilde{v}\|^2_{L_2(0,T;L_2(\Omega))} + \frac{1}{2}\|w - \tilde{w}\|^2_{L_2(0,T;L_2(\Omega))} \to \inf, \tag{7}$$

where $\theta, l, k, \alpha, \beta, R$ are constants, $k > 0$, $\tilde{v}, \tilde{w} \in L_2(0,T;L_2(\Omega))$ are given functions, $v$, $w$ are unknown functions, $(u_1,\ u_2)$ is function of control. The goal is to minimize (7) when $(u_1, u_2)$ satisfy condition (6). Here we use results on the existence of the degenerate initial value problem and the optimal control problem which presented in [1]. Simplified version of system (1)–(7) is considered in [2]. So, we are looking through the work in order to prove the conditional gradient numerical method convergence for the control problem. Besides, we've described the algorithm of numerical solution constructing.

**Keywords:** optimal control, degenerate evolution, numerical solution

**2010 Mathematics Subject Classification:** 49J20,49M05,64K10

**References:**

[1] Plekhanova M.V., Fedorov V.E. An optimality criterion in a control problem for a Sobolev-type linear equation. Journal of Computer and Systems Sciences International, 2007, vol. 46, no. 2, pp. 248—254.

[2] Plekhanova M.V., Baybulatova G.D. Conditional gradiend method for a robust control problem to a degenerate evolution system. Chelyabinsk Physical and Mathematical Journal, 2016, vol. 1, iss. 1, pp. 81—92. (In Russ.).

## About the robustness of the middle stabilizing controller for quasi-linear state dependent coefficients discrete-time systems

Yulia DANIK [1]

[1] *Institute for Systems Analysis Federal Research Center "Computer Science and Control", Russian Academy of Sciences, Moscow, Russia*
*E-mail: yuliadanik@gmail.com*

**Abstract:** This paper is dedicated to the robustness analysis of a stabilizing controller for quasi-linear state dependent coefficients discrete systems. Two nonlinear stabilizing regulator construction algorithms based on different versions of the discrete state dependent Riccati equation are discussed, the one from [1], [2] and the new one based on the discrete maximum principle. The robustness or fault-tolerance is the property of the regulator that allows it to maintain the control quality if the parameters of the real model differ from the computational model. There are different approaches for robustness analysis, for example the Afanas'ev minimax method, the robust stability properties of Schur polynomials analogous to Kharitonov theorems, linear matrix inequalities (LMI) approach [3] - [4] etc. Here the interval parametric uncertainties in the linear part of the system are investigated. The proposed nonlinear stabilizing regulator is calculated at the average values of the uncertainty parameters and is used for all realizations of the uncertain system. The basic idea is that the existence of only weak nonlinearity in the system allows us to study its robustness based on the robustness of the corresponding unperturbed discrete linear system. First we consider the linear part of the closed-loop system obtained along the so-called middle controller and then take into account the nonlinear part of the system, by introducing some conditions. The sufficient robustness conditions are formulated in the form of linear matrix inequalities. The results of the numerical experiments that demonstrate the robustness of the closed-loop system are analyzed [5] - [6].

**Keywords:** nonlinear discrete control systems, SDRE, regulator construction, robustness, stabilization, discrete maximum principle

**2010 Mathematics Subject Classification:** 93D05, 93D09, 93D15, 93D20, 93C10, 93C55

**Acknowledgements** This work was supported by the Russian Science Foundation (Project No. 16-11-00048).

**References:**

[1] Emel'yanov, S.V., Danik, Yu.E., Dmitriev, M.G., Makarov, D.A., "Stabilization of nonlinear discrete time dynamic control systems with a parameter and state-dependent coefficients ", *Doklady Mathematics 2016*,Vol. 93, No.1,pp.121–123, 2016.

[2] Danik, Yu.E., Dmitriev, M.G., Makarov, D.A., "An algorithm for constructing regulators for nonlinear systems with the formal small parameter ", *Informatsionnye tekhnologii i vychislitel'nye sistemy* , No.4,pp.35–44, 2015. (in Russian)

[3] Ramos, D. C. W., Peres, P. L. D., "A less conservative LMI condition for the robust stability of discrete-time uncertain systems ", *System Control Lett* , Vol. 43,pp.371–378, 2001.

[4] Oliveira, R.C.L.F., Peres, P.L.D., "Stability of polytopes of matrices via affine parameter-dependent Lyapunov functions: Asymptotically exact LMI conditions ", *Linear Algebra Appl.*,Vol. 405, No.8,pp.209–228, 2005.

[5] Danik, Yu.E., Dmitriev, M. G., "The robustness of the stabilizing regulator for quasilinear discrete systems with state dependent coefficients ", *International Siberian Conference on Control and Communications (SIBCON2016). Proceedings. Moscow: The Tomsk IEEE Chapter & Student Branch* , Russia, Moscow, May 12-14, 2016.

[6] Danik, Yu.E., "Robustness to parametric disturbancies of the quasi-linear discrete-time system ", *Informatics, control & systems analysis, ICSA- 2016.* , Tver, June 8-11, 2016 (In Russian), accepted.

# On a controllability for a class of $2D$-linear modular dynamic systems with periodic parameters

F.G. FEYZIEV [1], G.H. MAMMADOVA [2], F.N. NABI-ZADE [3]

[1,3] *Sumgait State University, Sumgait, Azerbaijan*
*E-mail: FeyziyevFG@mail.ru*
[2] *Institute of Applied Mathematics, Baku State University, Baku, Azerbaijan*
*E-mail: gamar.mamadova@gmail.com*

**Abstract:** The homogeneous $2D$- linear modular dynamic systems (2D-LMDS) with periodic parameters are described by the following equations (1)-(3):

$$x[n+1,c] = \sum_{\rho\in P} A[n,\rho]\, x[n,c+\rho] + \sum_{q\in Q} B[n,q]\, u[n,c+q],\ \ GF\,(p)\,, \tag{1}$$

$$y[n,c] = \sum_{\tau\in R} G[n,\tau]\, x[n,c+\tau],\ \ GF\,(p)\,. \tag{2}$$

$$A[n,\rho] = A[n+M,\rho],\ B[n,q] = B[n+M,q],\ G[n,\tau] = G[n+M,\tau]. \tag{3}$$

Based on the definitions (4)

$$q_i[k,c] = x[k\cdot M+i-1,c],\ \ z_i[k,c] = u[k\cdot M+i-1,c],$$
$$\gamma_i[k,c] = y[k\cdot M+i-1,c],\ w_i[k,c] = z_i[k+1,c],\ i=1,...,M;$$

$$q[k,c] = (q_1[k,c],...,q_M[k,c]\,)^T\,,\ \ z[k,c] = (z_1[k,c],...,z_M[k,c]\,)^T \tag{4}$$

$$\hat{q}\,(k,c) = (q_1[k,c],...,q_M[k,c],\, z_1[k,c],...,z_M[k,c])^T\,.$$

2D-LMDS (1)- (3) is written in the form called a block model

$$\hat{q}[k+1,c] = \sum_{\rho\in D} \tilde{A}[\rho]\,\hat{q}[k,c+\rho] + \sum_{q\in L} \tilde{B}[q] w[k,c+q],\ GF\,(p)\,, \tag{5}$$

$$\gamma[k,c] = \sum_{\tau\in\Phi} \tilde{G}[\tau]\,\hat{q}[k,c+\tau],\ \ GF\,(p)\,. \tag{6}$$

**Theorem.** In order to 2D-LMDS described by the (1-3) be fully $N$- controllable it is sufficient fulfillment of the condition: the corresponding block model (5), (6) have to be $k_N$ controllable, where

$$k_N = k_1 - k_0,\ k_1 = int\,(n_1/N)\,,\ k_0 = int\,(n_0/N)\,.$$

**Keywords:** modular dynamic systems with periodic parameters, a block model

**2010 Mathematics Subject Classification:** 37N35 , 11S31

**References:**

[1] Feyziev F.G., Faradjeva M.R. The modular sequential machines, Baku, "Elm", 2006, p. 234.

## On the systems of rational difference equations

Allaguly GURBANLYYEV

*Department of Mathematics, Necmettin Erbakan University, Konya, Turkey*
*E-mail: akurbanli@yahoo.com*

**Abstract:** In this paper, we have investigated the solutions of the system of rational difference equations

$$\begin{cases} x_{n+1} = \frac{\alpha y_{n-2}}{\beta+\rho y_{n-2}x_{n-1}y_n} \\ y_{n+1} = \frac{\delta x_{n-2}}{\zeta+\eta x_{n-2}y_{n-1}x_n} \\ z_{n+1} = \frac{\theta z_{n-2}}{\lambda+\mu y_{n-2}x_{n-1}y_n} \end{cases}$$

where the parameters $\alpha, \beta, \rho, \delta, \zeta, \eta, \theta, \lambda, \mu$ and initial values $x_{-i},\ y_{-i},\ z_{-i}$, $i \in \{0, 1, 2\}$ are reel numbers.

**Keywords:** difference equation, difference equation systems, solutions

**2010 Mathematics Subject Classification:** Applied Mathematics, MS6:

**References:**

[1] Kurbanli, A. S., Çinar, C., Yalcinkaya, İ., "On the behavaior of positive solutions of the system of rational difference equations $x_{n+1} = \frac{x_{n-1}}{y_n x_{n-1}+1}$, $y_{n+1} = \frac{y_{n-1}}{x_n y_{n-1}+1}$", *MathematicalandComputerModelling*, vol. 53, No 5-6, pp. 1261-1267, 2011.

[2] Kurbanli, A. S., "On the behavaior of positive solutions of the system of rational difference equations $x_{n+1} = \frac{x_{n-1}}{y_n x_{n-1}-1}$, $y_{n+1} = \frac{y_{n-1}}{x_n y_{n-1}-1}$, $z_{n+1} = \frac{1}{y_n z_n}$", *Advances in Difference Equations,* 2011:40, 2011.

[3] Kurbanli, A. S., "On the behavior of solutions of the system of rational difference equations $x_{n+1} = \frac{x_{n-1}}{y_n x_{n-1}-1}$, $y_{n+1} = \frac{y_{n-1}}{x_n y_{n-1}-1}$, $z_{n+1} = \frac{z_{n-1}}{y_n z_{n-1}-1}$", *Discrete Dynamics in Nature and Society*, 2011, Volume 2011, Article ID 932362, 12 pages, doi:10.1155/2011/932362, 2011.

[4] Kurbanli, A. S., "On the behavior of solutions of the system of rational difference equations $x_{n+1} = \frac{x_{n-1}}{y_n x_{n-1}-1}$, $y_{n+1} = \frac{y_{n-1}}{x_n y_{n-1}-1}$". *World Applied Sciences Journal*, 10 (11), pp. 1344-1350, 2011.

[5] Kurbanli, A. S., Çinar, C., Simsek, D., "On the Periodicity of Solutions of the System of Rational Difference Equations $x_{n+1} = \frac{x_{n-1}+y_n}{y_n x_{n-1}-1}$, $y_{n+1} = \frac{y_{n-1}+x_n}{x_n y_{n-1}-1}$", *Applied Mathematics*, Vol 2, Number 4, pp. 410-413, 2011.

[6] Kurbanli, A. S., Çinar, C., Erdogan, M. E., "On the behavaior of solutions of the system of rational difference equations $x_{n+1} = \frac{x_{n-1}}{y_n x_{n-1}-1}$, $y_{n+1} = \frac{y_{n-1}}{x_n y_{n-1}-1}$, $z_{n+1} = \frac{x_n}{y_n z_{n-1}}$", *Applied Mathematics*, Vol.2, pp. 1031-1038, 2011.

[7] Gurbanlyyev, A., "On a system of difference equations". *European Journal of Mathematics and Computer Science*, 3 (1), pp. 1-14, 2016.

[8] Gurbanlyyev, A., "On the behavior of solutions of the system of rational difference equations". *European Journal of Mathematics and Computer Science*, 3 (1), pp. 23-42, 2016.

## Design of PI regulators for dynamic systems with constrained control and fixed endpoints of trajectories

Zainel MURZABEKOV

*Research Institute for Mathematics and Mechanics, al-Farabi Kazakh National University, Almaty, Kazakhstan*
*E-mail: murzabekov-zein@mail.ru*

**Abstract:** The construction of the automatic operation control for real and complex systems requires the use of new information technologies, it is actually necessary to develop new principles of design the systems with a high level of complexity. In the field of automatic control published works one can find various examples of mathematical formulation and methods for solving optimal control problems [1]. But still, the development of various methods of constructing the PI and PID controllers with the necessary properties is an urgent problem [2].

In this work, the problem of optimal control for time-varying linear systems with fixed endpoints of trajectories is considered. A correspondent quadratic objective functional depends on the control, the state of the object and on its integral. New technique of designing the PI controller for the automatic control systems with box constraints on values of control is proposed. The problem is solved by using Lagrange multipliers of a special type [3].

**Keywords:** Optimal control problem, box constraints, Lagrange multipliers, PI regulator

**2010 Mathematics Subject Classification:** 49J15, 49N35

**References:**

[1] Atans, M., Falb, P.L., *Optimal Control: an Introduction to the Theory and its Applications*, Dover Publications, Inc., New York, 2006.

[2] O'Dwyer, A., *Handboopk of PI and PID Controller Tuning Rules*, Imperial College Press, London, 2009.

[3] Murzabekov, Z.N., "The Synthesis of the Proportional-Differential Regulators for the Systems with Fixed Ends of Trajectories Under Two-Sided Constraints on Control", *Asian Journal of Control*, Vol. 18, No. 2, pp. 494-501, 2016.

## Optimal control problem for the three-sector economic model of a cluster

Zainel MURZABEKOV [1], Shamshi AIPANOV [1],
Saltanat USUBALIEVA [1]

[1] *Research Institute for Mathematics and Mechanics, Al-Farabi Kazakh National University, Almaty, Kazakhstan*
*E-mail: murzabekov-zein@mail.ru, aipanov@mail.ru, salta-74@mail.ru*

**Abstract:** Problems of optimal control for dynamic systems with fixed ends of trajectories are often used in practical applications. Various mathematical formulations of optimal control problems and their classification are given in [1].

In this work the problem of optimal control for the three-sector economic model of a cluster is considered [2]. Problem statement is to determine the optimal distribution of investment and manpower in moving the system from a given initial state to desired final state. To solve the optimal control problem with finite-horizon planning, in case of fixed ends of trajectories, with box constraints, method of Lagrange multipliers of a special type is used [3]. This approach allows to represent the desired control in the form of synthesis control, depending on state of the system and current time. The results of numerical calculations for a three-sector economic model show the effectiveness of the proposed method.

**Keywords:** Economic model, cluster, investments, manpower, optimal control problem, method of Lagrange multipliers

**2010 Mathematics Subject Classification:** 49J15, 49N35

**References:**

[1] Geering, H.P., *Optimal Control with Engineering Applications*, Springer-Verlag, Berlin-Heidelberg, 2007.

[2] Murzabekov, Z., Milosz, M. Tussupova, K., "Modeling and Optimization of the Production Cluster", *Proceedings of 36th International Conference on Information Systems and Architecture and Technology ISAT-2015, Part II: Advances in Intelligent Systems and Computing*, Springer International Publishing, Switzerland, 2016, pp. 99-108.

[3] Aipanov, Sh.A., Murzabekov Z.N., "Analytical Solution of a Linear Quadratic Optimal Control Problem with Control Value Constraints", *Journal of Computer and Systems Sciences International*, Vol. 53, No. 1, pp. 84-91, 2014.

# The method of accelerated convergence for constructing conditional-periodical solutions

Zhussip SULEIMENOV [1]

[1] *Al-Farabi Kazakh National University, Kazakhstan*
*E-mail: zh_suleimenov@mail.ru*

**Abstract:** In the present paper we take quasi-linear system of differential equations

$$\frac{dx}{dt} = Ax + \varepsilon f(t,x), \tag{1}$$

where $x = colon(x_1, x_2)$, $A = (a_{jk}), j = k = 1, 2$,
$f(t,x) = colon(f_1(t,x_1,x_2), f_2(t,x_1,x_2))$
conditionally-periodic by $t$ with frequency basis $\omega_1, \omega_2, ..., \omega_n$; analytical by $t$ and $x$ in the domain
$D = \{(t,x) \subset C^3 : ||x|| \leq h, ||Im\omega t|| \leq q\}$ function, $det|A - \lambda E| = 0$ has purely imaginary roots $i\sigma_1, i\sigma_2$, and rational numbers $\sigma_1, \sigma_2$ non-co-measurable with $\omega_1, \omega_2, ..., \omega_n$, $\epsilon$ is a small parameter.

In order to find a conditionally-periodic solutions of (1) the method of accelerated convergence [1] is applied. As an initial approximate conditionally-periodic solutions of the system (1) $x^{(0)}(t,\varepsilon) = 0 := colon(0,0)$ is chosen. Its residual denoted by $x^{(1)}(t,\varepsilon)$ and take this function as a first approximation to the original conditionally-periodic solutions of the system (1).

**Keywords:** quasi-linear system of differential equations, conditionally-periodic solution, small parameter

**2010 Mathematics Subject Classification:** 34B15

**References:**

[1] Bogolyubov N., Mitropolsky Y., Samoilenko A. *The method of accelerated convergence in nonlinear mechanics*. Kiev, "Naukova Dumka", 1969.

## Martinet-Ramis modulus for one Quadratic System

Mikhail M. TUROV[1]

[1] *Department of Mathematical, Chelyabinsk State University, Chelyabinsk, Russia*
*E-mail: turov_m_m@mail.ru*

**Abstract:** There are considered a quadratic system

$$\begin{cases} \dot{p} = p(1-v) \\ \dot{v} = v(p-v) \end{cases} \tag{1}$$

This system is in some sense, limit system for well known Jouanolou system [1]. The system (1) has a saddlenode singularity at the origin. In this work we calculate first coefficients of Martinet-Ramis' modulus [2].

Martinet-Ramis' modulus $(C, \phi)$ (for saddlenode singular point) [2] are constructed by transformations reducing initial system to its (orbital) formal normal form. Solutions of the system (1) with given initial conditions can be found as a series (with respect to initial condition). Using these solutions it is possible to find coefficients of the normalizing transformations, and then to determinate coefficients of modulus.

As a result we get

**Theorem 1.** *Let $(C, \phi)$ be Martinet-Ramis modulus for (1). Then: $C = 0$, $\phi(z) = z + 2\pi i z^2 + (2\pi i - 4\pi^2)z^3 + ...$*

**Corollary 2.** *The system (1) is not analytically orbital equivalent to its formal normal form.*

**Keywords:** Martinet-Ramis modulus, saddlenode, central manifold

**2010 Mathematics Subject Classification:** 34C20, 37F75

**References:**

[1] J.P. Jouanolou, "Equations de Pfaff Algebriques", Lecture Notes in Mathematics 708, Springer-Verlag, 1979.
[2] Martinet, J., Ramis, J.-P., "Classification analytique des equations differentielles non lineaires resonnantes du premier ordre", *Ann. Sci. Ecole Norm. Sup. (4) 16 (1983)*, No.4, pp. 571-621.

## Soliton solutions of the Hirota equation

Kuralay YESMAKHANOVA [1], Gaukhar SHAIKHOVA [1],
Guldana BEKOVA [1]

[1] *Eurasian International Center for Theoretical Physics and Department of General and Theoretical Physics,*
*Eurasian National University, Astana, Kazakhstan*
*E-mail: kryesmakhanova@gmail.com*

**Abstract:** It is well known that nonlinear integrable systems have attracted a lot of attention among researchers. This fascinating subject of nonlinear science has branched out in almost all areas of technology and science. In nonlinear science soliton solutions play an important role. There are many ways to obtain soliton solutions of the nonlinear evolution equations, such as the Painleve analysis [1], the Hirota's bilinear method [2], Darboux transformation (DT) [3] and so on. Among the various methods, the DT has been proved very successful in driving different kinds of solutions for many of the integrable equations from a trivial seed. In this work, we focus on the construction soliton solutions for the 2+1-dimensional Hirota equation [4], which is modified nonlinear Schrodinger equation. One-soliton solutions are obtained by means of the one-fold Darboux transformation for the 2+1-dimensional Hirota equation.

**Keywords:** Hirota equation, Darboux transformation, soliton solution

**2010 Mathematics Subject Classification:** 35C08, 35Q51, 37K40

**References:**

[1] Bindu S.G. , Mahalingam A. and Porsezian K. "Dark soliton solutions of the coupled Hirota eqaution in nonlinear fiber", *Physics Letters A*, Vol.286, pp.321-331, 2001.

[2] Gui M., Zhenyun Q. "Construction of Nth-order rogue wave solutions for Hirota equation by means of bilinear method", *www.arxiv.org*, 2014.

[3] Yesmakanova K.R., Shaikhova G.N., Bekova G.T. and Myrzakulova Zh.R. "Determinant Reprentation of Dardoux Transformation for the (2+1)-Dimensional Schrödinger-Maxwell-Bloch Equation", *Advances in Intelligent Systems and Computing*, Vol. 441, pp.183-198, 2016.

[4] Myrzakulov R., Mamyrbekova G. K., Nugmanova G.N and Lakshmanan M. "Integrable (2+ 1)-dimensional spin models with self-consistent potentials", *Symmetry*, Vol. 7(3), pp.1352-1375, 2015.

# AUTHOR'S INDEX

# Index


Çapan H., *40*
Çopuroğlu E., *137*

Ağirseven D., *177*
Abdenov A.Z., *210*
Abdigaliyeva A., *124*
Abdrasheva G.K., *269*
Abduakhitova G., *113*
Abdualiyeva M., *195*
Abdullaev B., *168*
Abiev N., *296*
Abishev N., *193*
Abyar A., *213*
Abylayeva A., *24, 25*
Abzhapbarov A., *138*, *230*
Adamov A., *250*
Adeyefa E., *125*
Aghayan S.M., *26*
Ahmetzhanova M., *226*
Aida-Zade K., *224*
Aimakhanova A., *262*
Aipanov Sh., *309*
Aisagaliev S.A., *297, 298*
Aitzhan A., *225*
Akanbay N., *200, 201*
Akhmedov D.Sh., *299*
Akhundov H., *300*
Akhymbek M., *27*
Akimzhan N., *188*
Akishev G., *28*
Akysh A.SH., *29*
Aldashev S. A., *30*
Alekhina V., *202*
Alexeyeva L., *226*
Ali J., *31*
Alibiyev D., *126*, *240*
Aliev F.A., *227*, *301*
Aliev N.A., *301*
Alimov Sh., *263*
Alp M., *162*
Altynbekov Sh., *195*
Amanbaev T., *228*
Amangaliyeva M., *264, 265*
Annadurdyyeva B., *186*
Aptekarev A.I., *32*
Arepova G., *266*, *275*
Aripov M., *229*
Ashirbayev N., *230, 231*
Ashirbayeva Z., *231*
Ashirbayeva Zh., *230*
Ashyralyev A., *33*, *56*, *127–132*, *135*, *177, 178*, *267, 268*, *302*
Ashyralyyeva M., *187*, *191*
Askerov I. M., *212*
Assanova A., *34*
Atakhan N., *35*
Attaev A., *36*
Auzhani Y., *170*
Ayaz S., *92*
Ayazbayeva A.M, *297*
Aydemir K., *133*, *160*
Aytzhan A., *235*
Azimov A., *134*

Başar F., *39, 40*, *115, 116*
Baituyakov Sh., *37*
Baituyakova Zh., *37*
Baizhanov B., *203*
Bakanov G.B., *232*
Balgimbayeva Sh.A., *38*
Baybulatova G., *303*
Bayramov S., *204*
Bazarkhanov D., *40*
Beketaeva A., *255*
Bekjan T.N., *41*
Bekmaganbetov K.A, *42*
Bekmoldayeva R., *231*
Bekova G., *312*
Belakroum Kh., *135*
Bidaibekov Y., *188–190*, *193*
Bilall Sh., *43*
Biyarov B.N., *269*
Bizhanova G., *136*
Bizhigitova N., *239*
Bokayev N., *44*
Borikhanov M.B., *136*
Bostanov B., *189*
Bouhadjera H., *137*
Burenkov V., *44*

Cakalli H., *45, 46*, *69*, *233*
Charyyev A., *187*, *191*
Chaudhary S., *153*

Dairbekov N., *47*
Danik Y., *304*
Darooneh A.H., *157*, *167*
Dasibekov A., *138*
Dauitbek D., *259*, *278*, *293*
Dauylbayev M., *49*
Demidenko G., *50*
Denisov V. N., *51*
Derevtsov E., *234*
Dildabek G., *52*, *291*
Djamalov G., *53*
Dmitriev D., *206*
Dmitriev M., *139*
Duisebayeva P., *138*
Dzhobulaeva Zh., *54*
Dzhumabaev D.S., *55*

Elhaffaf A., *86*
Erdem Coskun A., *233*
Erdogan A.S., *56*
Erdogan S., *140*
Ergaliyev M., *265*
Erzhanov N., *57*
Eshkuvatov K., *141*
Et M., *45*

Falaleev L., *58*
Fateev Yu., *206*
Fazullin Z., *270*
Fedorov V.E., *59*
Feyziev F.G., *306*
Folaranmi R.O., *125*

Gasimova K.G., *301*
Georgievskiy P., *244*
Gordievskikh D.M., *59*
Grinshkun V., *190*
Guezane-Lakoud A., *135*, *142*
Guliev A.P.., *301*
Gulsen T., *271*
Gunduz Aras C., *204, 205*, *233*
Gurbanlyyev A., *307*
Gurbanov B., *127, 128*

Hajieva N.S., *227*
Hamza R., *104*
Hatami H., *143*
Hosseini Shekarabi F., *143*

Ibraeva E., *126*
Ibrayeva E., *240*
Igenberlina A., *74*
Igisinov S., *82*
Ilyasova M., *37*
Imanbaev N., *272*
Imanberdiyev K., *264*
Inkarbekov M., *235*
Inkarbekov M., *225*
Iskakov K., *212*
Iskakova G., *197*
Iskakova U., *60*
Iskenderova D., *144*
Ismailov B., *236*
Ismailov Kh., *236*
Israfilov D., *61*
Issagali A., *146*
Issagali A., *145*
Issakhov A., *237*
Issakhov A., *207*
Ivashchenko A., *249*

Jenaliyev M., *264, 265*, *273*
Jussupov A., *212*

Kabdrakhova S., *62*
Kadirkulov B., *63, 64*
Kahraman T., *129*
Kal'menov T., *66*, *275*
Kaliev I., *238*
Kalimbetov B., *192*
Kalimbetov B., *65*
Kalmenov T. Sh., *274*
Kaltayev A., *225*, *235*
Kalybay A., *67, 68*
Kamalova G., *188–190*
Kanguzhin B., *276–279*
Kaplan H., *69*
Karabaeva B., *130*
Karaca B., *131*
Karachik V., *147*
Karachik V.V., *70*
Karakuş M., *71, 72*
Karimov Sh.T., *73*
Karimova U.Y., *153*
Karshygina G., *74*
Kasenov S., *134*
Kasenov S., *148*
Kaskataeva B., *193*
Kassabek S., *241*
Kassymetova M., *222*
Kassymov A., *284*
Kavokin A., *239*
Kavokin A.A., *106*
Kayshibayeva G., *226*
Kazhikenova A., *126*, *240*
Kerimbekov A., *280*
Kerimkhulle S., *208*
Khairullin E., *75*
Khakimzyanov G., *126*
Khaldi R., *142*
Khalifeh K., *157*
Khamnei H.J., *149*
Khan A., *150*
Kharin S.N., *241*, *253*
Khompysh Kh., *242*
Kitaybekov E., *30*
Klunnikova M., *194*
Kočinac L.D.R., *209*
Koksal M.E.., *302*
Koldas A., *225*
Kopzhassarova A., *289*
Korneeva A., *151*
Kornienko V., *254*
Kornilov V., *188*
Koshanov B., *152*
Koshanova M., *76*
Koshkarova B., *78*
Kovrizhkin V., *76*
Kremez N., *206*
Kritskov L., *281*
Kropotin N., *214*
Kulakhmetova A.T., *106*
Kulbaev D., *210*
Kulpeshov B., *211*
Kumar S., *153*
Kurmanbaev D., *79*
Kusherbayeva U., *113*
Kussainova A., *212*
Kussainova L., *78*
Kydyrmina N., *80*

Lambda S., *243*
Levin V., *244*, *246*

Maden s., *153*
Madiyarov N., *195*
Makarov D., *139*
Makarov R., *219*
Makasheva A., *247*
Makhdoom I., *155*
Makouyi R., *149*
Maltseva S., *234*, *257*
Mamatova G.U., *176*
Mamedov B.A., *137*, *212*
Mammadova G.N., *306*
Manapova A., *156*
Manuylovich I., *246*
Markov V., *246*
Masanov S., *147*
Matin D., *44*
Matyakubov A., *81*
Meldebekova S., *236*
Minglibayeva B.B., *55*
Mirzakulova A., *49*
Moghaddasi H., *157*
Mohammadzadeh M., *213*
Motarjem K., *213*
Muhtarov F.S., *158*
Mukhamadiev ., *84*
Mukhambetzhanov S., *238*
Mukhtarov O.Sh., *133*, *158*, *160*
Muminov Z., *141*
Muradoglu Z., *159*
Muratbekov M., *82, 83*
Muratbekov Ma., *83*
Murzabekov Z., *308, 309*
Mutallimov M., *300*
Mynbaev K., *85*

Nabi-zade F.N.., *306*
Naceri M., *86*
Naimanova A., *247*, *255*
Naimov ., *84*
Nalzhupbayeva G., *277*
Nasiri P., *155*
Nasirova T., *153*
Nauryz T., *239*
Nazhimov R.R., *59*
Nik Long N. M. A., *141*
Niyazova R., *249*
Nouri H., *253*
Novikov I., *87*
Nugaeva I., *270*
Nurakhmetov D., *282*
Nurmyradova N., *186*
Nurseitov D., *134*, *148*
Nurseitova A., *148*

Nursultanov E., *88*
Nyssanov E., *248*

Oinarov R., *89, 90*
Okur U., *132*
Olğar H., *133*, *158*, *160*
Omarova I., *192*
Omarova Sh., *196*
Oralbekova Zh., *212*
Oralsyn G., *91*
Orazov I., *52*, *92*
Orumbayeva N., *161*, *196*
Ospanov K., *93, 94*
Ozdemir Y., *140*, *162*, *165*
Ozesenli Tetikoglu F.S., *129*
Ozturk Beigmohammadi E., *163*

Pak A., *155*
Pak N., *194*
Panakhov E.S., *271*
Panov A., *164*
Pekonur E., *165*
Penkin O., *47*
Piskarev S., *95*
Plekhanova M., *96*
Polatbek A., *138*
Polyakova A., *257*
Popivanov N., *97*
Pskhu A., *98*
Pulkina L., *99*
Pushkaryeva T., *194*
Pya N., *219*
Pyatkov S., *166*
Pyrkova A., *249*

Rahimov A., *224*
Rajabov M.F., *227*
Rajabov N., *100*
Rakhimova R., *219*
Rakhymbek D., *195*
Ramazanov M., *264, 265*, *273*
Ramazanova G., *182*
Ranjbari A., *98*
Raskaliyev A., *299*
Rezaei S., *167*
Rogovoy A., *101*
Ruzhansky M., *283*
Rysbaiuly B., *250*

Sabitbek B., *274*, *283*
Sabitova G., *238*
Sadullaev A., *168*
Sadybekov M., *27*
Sadybekov M.A., *66*, *251, 252*, *272*, *284–287*
Safarova N.A., *227*
Sakabekov A., *145, 146*, *170*
Sakhaev Sh., *242*
Sakhit A., *238*
Salimov R., *127, 128*
Salimov S.M., *256*
Sapakov D., *65*
Saprygina M., *248*
Sarsenbi A., *130*, *251*, *281*, *286*, *288*
Sarsenbi Abdis., *268*
Sarsengeldin M., *241*, *253*
Sarybekova L., *47*
Savas E., *46*, *102, 103*
Sazaklioglu A.U., *56*
Seba D., *104*
Sengul H., *45*
Serikbaev J., *63, 64*
Serovajsky S., *134*, *197*
Sevryugin I., *298*
Shaidurov V., *151*, *206*, *254*
Shaikhova G., *312*
Shaimardan S., *105*
Shakenov K., *171*
Shakhan N., *255*
Shaldanbayev A., *289*
Shalginbayeva S., *43*, *67, 68*, *262*
Shamatayeva N., *221*
Sharipov Kad.S., *172*
Sharipov Kub.S., *172*
Sharipov S., *172*
Shaukenova K., *197*
Shayakhmetova B., *196, 197*
Shayakmetova B., *161*
Shikhlinskaya R.Y., *256*
Shkalikov A., *290*
Shomanbayeva M., *230*, *289*
Shpadi Yu.R., *106*
Shyrakbaev A., *82*
Skubachevskii A.L., *173*
Smailov E., *107*
Smatova G., *152*
Soldatov A.P., *108*
Somuncu E., *212*
Sonmez A., *233*
Starodumov I., *202*, *214*
Starodumov O., *202*
Stupina T., *194*
Suleimenov Zh., *310*
Suleymenova Za., *74*
Suleymenova Zo., *201*
Sultanbek T., *231*
Sultanov M., *234*
Sultanova M., *171*
Suragan D., *274*, *283*
Svetov I., *234*, *257*

Türk K., *177*
Taimanov I.A., *216*
Tapeeva S., *201*
Taskin A., *267*
Tasmambetov Zh., *108*
Temesheva S., *110*
Temirbekov A., *175*
Temirbekov N., *173*
Temirkhanova A., *89*, *111*
Tengayeva A., *251*, *286*, *291*
Testici A., *61*
Tetikoglu F.S., *33*
Thakur B.S., *111*
Tilleuov G., *228*
Tleubergenov M., *217*
Tleukhanova N., *88*, *119*
Tleulesova A., *112*
Tokibetov Zh., *113*
Tokmagambetov N., *276*
Toktorbaev A., *144*
Toleugazy Ye., *42*
Torebek B.T., *70*, *114*, *285*
Tuğ O., *115, 116*
Tulegenova B., *228*
Tulenov K., *258, 259*, *292, 293*
Turarov A., *173*
Turkcan K.T.., *302*
Turmetov B., *76*
Turmetov B.Kh., *114*, *136*, *147*
Turov M., *311*
Tyapkin V., *206*
Tyurekhojayev A.N., *176*

Ulbrikht O., *220*
Umbebaev K., *189*
Umirbaev ., *218*
Urun M., *178*

Usmanov K.I., *76*
Usubalieva S., *309*
Uygul F., *117*
Uzun M., *179, 180*

Vasilyev V., *118*
Vassilina G., *217*
Voinov V., *219*
Voinov Y., *219*
Vyas V.K., *117*
Vyatkin A., *254*

Yarmohamadi M., *155*
Ydyrys A., *119*
Yeshkeyev A., *220–222*
Yeskermessuly A., *120*
Yesmakhanova K., *312*
Yessirkegenov N., *121*, *252*, *287*, *294*
Yildirim O., *179, 180*

Zakir'yanova G. K., *181*
Zaman G., *150*
Zhakebayev D., *124*
Zhakenova M., *182*
Zhanbolova A., *197*
Zhanbusinova B., *197*
Zhapbasbayev U., *182*
Zhapsarbayeva L., *279*
Zhubatov Zh., *124*
Zhukabayeva T., *212*
Zhumali A., *260*
Zhumanova L., *262*
Zhumatov S., *183*
Ziaie A.H., *184*
Zireh A., *26*
Zulkarnain F.S., *141*
Zulkhazhav A., *94*