

First regularized trace of integro-differential Sturm-Liouville operator on a segment with punctured points at generalized conditions of bonding in deleted points

Abdisalam A. Sarsenbi and Lyazzat K. Zhumanova

Citation: **1789**, 040009 (2016); doi: 10.1063/1.4968462

View online: <http://dx.doi.org/10.1063/1.4968462>

View Table of Contents: <http://aip.scitation.org/toc/apc/1789/1>

Published by the [American Institute of Physics](#)

Articles you may be interested in

[On a Green's function and eigenvalues of a second-order differential operator with involution and Neumann boundary conditions](#)

1789, 040008 (2016); 10.1063/1.4968461

[Basis properties of eigenfunctions of the differential operator \$-u''\(-x\) + q\(x\)u\(x\)\$ with Cauchy data](#)

1789, 040002 (2016); 10.1063/1.4968455

First regularized trace of integro-differential Sturm-Liouville operator on a segment with punctured points at generalized conditions of bonding in deleted points

Abdisalam A. Sarsenbi^{1,2,a)} and Lyazzat K. Zhumanova^{1,3,b)}

¹*Institute of Mathematics and Mathematical Modeling, 125 Pushkin str., 050010 Almaty, Kazakhstan*

²*Auezov South Kazakhstan state University, 5 Tauke Khan ave., 160012 Shymkent, Kazakhstan*

³*al-Farabi Kazakh National University, 71 al-Farabi ave., 050040, Almaty, Kazakhstan*

a)Corresponding author: sarsenbi.a.a@math.kz

b)zhumanova@math.kz

Abstract. The present work is devoted to calculating a first regularized trace of one integro-differential operator with the main part of the Sturm-Liouville type on a segment with punctured points at integral perturbation of "transmission" conditions. The integro-differential Sturm-Liouville operator $-y''(x) + q(x)y(x) + \gamma \int_0^\pi y(t)dt = \lambda y(x)$ given on the segments $\frac{\pi}{n}(k-1) < x < \frac{\pi}{n}k$, $k = \overline{1, n}$; $n \geq 2$ is considered. Boundary conditions of the Dirichlet type: $y(0) = 0$, $y(\pi) = 0$ are given on the left-hand and right-hand ends of the segment $[0, \pi]$. The functions continuous on $[0, \pi]$, the first derivatives of which have jumps at the points $x = \frac{\pi}{n}k$, are solutions. The value of jumps is expressed by the formula $y'(\frac{\pi k}{n} - 0) = y'(\frac{\pi k}{n} + 0) - \beta_k \int_0^\pi y(t)dt$, $k = \overline{1, n-1}$. The basic result of the paper is the exact formula of the first regularized trace of the considered differential operator.

1 Introduction

At the present stage of its development, the spectral theory of ordinary differential operators is one of the important parts of the general spectral theory and is actively developing by different mathematical schools. One of the spectral theory directions is the trace theory of differential operators. Primarily, the trace theory has been developed by the Moscow school under the head of academician V.A. Sadovnichii [1, 2]. In recent years they have presented the formula of first regularized trace of discrete operators, regularized traces of singular operators, regularized traces of self-adjoint discrete operators with a non-nuclear resolvent, trace formula of M.G. Krein in case of perturbations such as Hilbert-Schmidt, regularized trace of the operator Sturm-Liouville equation on a finite interval, containing a delta-function, the trace formula for potential, formulas for regularized traces of operators with relatively compact perturbations, and so on.

However, a number of essential problems of spectral theory is still not resolved. Finding regularized traces of differential operators in areas with deleted points is regarded to these problems. This direction is closely related to the research of operators with potentials, containing a delta-function, however this direction also has its own characteristics. Nowadays, this direction is in the process of accumulating the primary knowledge. In order to accumulate the primary knowledge we have to obtain formulas of explicit form of calculating regularized traces of different specific differential operators in deleted points areas.

2 The concept of a regularized trace

The theory of regularized traces of linear operators originating from the fundamental fact of finite-dimensional theory is an invariance of the matrix trace of a linear operator and its coincidence with the spectral trace, and explores the question about spreading the concept of trace invariance on the unbounded operators.

Researching regularized traces of operators with discrete spectrum was started in the famous work of I.M. Gel'fand and B.M. Levitan [3], where the authors obtained a trace of the operator in Sturm-Liouville problem

$$-y''(x) + q(x)y(x) = \lambda y(x), \quad y'(0) = 0, \quad y'(\pi) = 0.$$

For $q(x) \in C^1[0, \pi]$, under fulfillment of condition $\int_0^\pi q(x)dx = 0$, the following formula

$$\sum_{n=0}^{\infty} (\lambda_n - \mu_n) = \frac{1}{4}(q(0) + q(\pi))$$

has been obtained, where λ_n are eigenvalues of the problem and $\mu_n = n^2$ are eigenvalues of the same problem with $q(x) = 0$. The sum of a series is called the first regularized trace of the operator. Formulas of this type are called the first regularized trace. One of the advantages of such formulas is that although the eigenvalues when $q(x) \neq 0$ can not be calculated in explicit form, the sum of the regularized trace is accurately determined and can always be calculated.

In the same year L.A. Dikiy came to similar results, he used a little bit different methods. The works of I.M. Gel'fand, M.G. Gasymova, R.F. Shevchenko, A.G. Kostyuchenko, V.A. Sadovnichii and many others were devoted to the obtained formulas for regularized traces of ordinary differential operators. The most common results for ordinary differential operators were obtained by V.B. Lidskii and V.A. Sadovnichii [4]. They found that the derivation of this type for a wide class of boundary value problems, generated by ordinary differential expressions on a finite interval with a complex spectral parameter, reduces to the study of regularized sums of roots of entire functions with a certain structure of the asymptotic.

3 Overview of similar category of works of Kazakh authors

Among the large number of published works on the formulas of regularized traces of differential operators it is necessary to note the works of Kazakh mathematicians. This is primarily the works of Professor B.E. Kanguzhin and his disciples and followers.

In [5] a mathematical model of forced oscillations package of flat plates with point elastic links was proposed. In [6, 7] a complete description of the correct solvability of boundary value problems for the biharmonic operator in a circle and ball was given, their correctly solvable finite perturbations arising in considering the problem in the deleted area were written. In [8] identities for eigenvalues of operator generated by ordinary differential expressions with internal boundary conditions were found. In [9] a complete description of correctly solvable boundary value problems for the Laplace operator in the circle and in the deleted circle was given, the formulas of resolvents of correct problems were shown. In [10] the Laplace operator in the pierced area was studied in the Hilbert space, an analogue of Green's formula was obtained and the class of self-adjoint extensions was described. In [11, 12] the class of correct problems for the m -Laplace operator in the deleted area was considered and formulas of the regularized trace were obtained.

[14] is the closest to the theme of this paper. In this paper the formula of the first regularized trace of a differential Sturm-Liouville type operator on a segment with deleted points under an integral perturbation of "jump" conditions of the first derivatives in the deleted points was obtained. We will give this result in more detail.

The problem of eigenvalues:

$$-y''(x) + q(x)y(x) = \lambda y(x), \quad \frac{\pi}{n}(k-1) < x < \frac{\pi}{n}k, \quad k = \overline{1, n}; \quad n \geq 2; \quad (1)$$

$$y(0) = 0, \quad y(\pi) = 0, \quad (2)$$

$$y\left(\frac{\pi k}{n} - 0\right) = y\left(\frac{\pi k}{n} + 0\right), \quad y'\left(\frac{\pi k}{n} - 0\right) = y'\left(\frac{\pi k}{n} + 0\right) - \beta_k \int_0^\pi y(t)dt, \quad k = \overline{1, n-1} \quad (3)$$

has been considered, where $q(x)$ are enough times differentiable real-valued function; β_k is a real constant, λ is a spectral parameter.

Note that in [13], [2, p. 112] formulas for the first regularized trace of the problem when $\beta_k = 0$, $k = \overline{1, n-1}$ for equation (1) with additional summands of the form $\sum_{k=1}^{n-1} \alpha_k y\left(\frac{\pi k}{n}\right)$ are written. These works as well as [14] are the most related to the theme of our problem under consideration.

The main result of [14] is:

THEOREM 1 [14] For the first regularized trace of problem (1) - (3) the following formula is true :

$$\begin{aligned} \sum_{m=0}^{\infty} \sum_{j=1}^{2n} \left(\lambda_{m,j} - (2nm + j)^2 - \left(1 + \frac{1}{2nm + j} \right) \frac{2}{\pi} \int_0^{\pi} q(t) dt \right) &= \\ &= -\frac{1}{4} (q(0) + q(\pi)) + \frac{1}{2\pi} \int_0^{\pi} q(t) dt, \end{aligned} \quad (4)$$

where $\lambda_{m,j}$ are eigenvalues of problem (1)-(3). Herewith, the eigenvalues have an asymptotic behavior $\lambda_{m,j} = s_{m,j}^2$, where

$$s_{m,j} = (2nm + j) + \frac{c_{1,j}}{2nm + j} + \frac{c_{2,j}}{(2nm + j)^2} + O\left(\frac{1}{(2nm + j)^3}\right), \quad (5)$$

$$c_{1,j} = \frac{1}{2\pi} \int_0^{\pi} q(t) dt, c_{2,j} = \frac{1 - (-1)^j}{2} \frac{i}{\pi} \sum_{k=1}^{n-1} (\beta_k + \beta_{n-k}) e^{\frac{ijk}{n}}, \quad (6)$$

$$j = \overline{1, 2n}, m = 0, 1, 2, \dots$$

Throughout this note we mainly use techniques from works [15-21].

4 Statement of the problem and formulation of the main result

Consider a problem on eigenvalues for an integral-differential operator:

$$-y''(x) + q(x)y(x) + \gamma \int_0^{\pi} y(t) dt = \lambda y(x), \quad \frac{\pi}{n}(k-1) < x < \frac{\pi}{n}k, \quad k = \overline{1, n}; \quad n \geq 2 \quad (7)$$

with the boundary conditions (2) and the generalized conditions of "bonding" (3) in the deleted points of the interval.

Our goal is to build formulas for the first regularized trace, similar to formula (4), for the spectral problem under consideration (7), (2) and (3).

The main result of this work is:

THEOREM 2 For the first regularized trace of problem (7), (2), (3) the following formula holds:

$$\begin{aligned} \sum_{m=0}^{\infty} \sum_{j=1}^{2n} \left(\lambda_{m,j} - (2nm + j)^2 - \left(1 + \frac{1}{2nm + j} \right) \frac{2}{\pi} \int_0^{\pi} q(t) dt \right) &= \\ &= -\frac{1}{4} (q(0) + q(\pi)) + \frac{1}{2\pi} \int_0^{\pi} q(t) dt - \gamma\pi, \end{aligned} \quad (8)$$

where $\lambda_{m,j}$ are eigenvalues of problem (7), (2), (3). In this case the eigenvalues at large m have the asymptotic behavior $\lambda_{m,j} = s_{m,j}^2$, $j = \overline{1, 2n}$, where the quantities $s_{m,j}$, $c_{1,j}$, $c_{2,j}$ are determined by formulas (5) and (6), respectively.

5 A short proof of the theorem

By standard calculations on each interval $I_k : \frac{\pi}{n}(k-1) < x < \frac{\pi}{n}k$, we can write the asymptotic behavior (when $|s| \rightarrow \infty$) of two linearly independent solutions of equation (1):

$$y_{1,k}(x, s) \sim e^{isx} \sum_{\nu=0}^{\infty} \frac{a_{\nu,k}(x)}{s^{\nu}}, \quad y_{2,k}(x, s) \sim e^{-isx} \sum_{\nu=0}^{\infty} (-1)^{\nu} \frac{a_{\nu,k}(x)}{s^{\nu}},$$

where $a_{0,k}(x) \equiv 1$,

$$a_{\nu,k}(x) = \frac{i}{2} \left\{ a'_{\nu-1,k}(x) - a'_{\nu-1,k} \left(\pi \frac{k-1}{n} \right) - \int_{\pi \frac{k-1}{n}}^x q(t) a_{\nu-1,k}(t) dt \right\}. \quad (9)$$

Here, as usual, it is assumed that the complex plane ($\lambda = s^2, s = \sqrt{\lambda}$) is divided into four sectors by rays $\arg s = 0$ and $\arg s = \frac{\pi}{2}$ and this asymptotic is available in each of the four sectors.

From the recurrence formula (7), we obtain

$$a_{1,k}(x) = -\frac{i}{2} \int_{\pi \frac{k-1}{n}}^x q(t) dt, \quad a_{2,k}(x) = \frac{1}{4} \left\{ q(x) - q\left(\pi \frac{k-1}{n}\right) - \frac{1}{2} \left(\int_{\pi \frac{k-1}{n}}^x q(t) dt \right)^2 \right\},$$

$$a_{v,k} \left(\pi \frac{k-1}{n} \right) = 0, \quad v = 1, 2, ;$$

$$y_{1,k} \left(\pi \frac{k-1}{n}, s \right) = e^{i \frac{\pi(k-1)s}{n}}, \quad y_{2,k} \left(\pi \frac{k-1}{n}, s \right) = e^{-i \frac{\pi(k-1)s}{n}}.$$

On each of the intervals I_k we build a general solution of the integro – differential equation (7) by the variation method of arbitrary constants. We are introducing the solution in the form

$$Y(x) = A_k(x)y_{1,k}(x, s) + B_k(x)y_{2,k}(x, s).$$

We get representation of the general solution of equation (7). This solution is a two-parameter family. We denote these constants by A_k^0, B_k^0 .

Satisfying the boundary conditions (2) and the generalized conditions of "bonding" (3), we obtain with respect to the constant A_k^0, B_k^0 a linear system of $2n$ equations, determinant $\Delta(s)$ of which will be characteristic determinant of the spectral problem (7), (2), (3).

The function $\Delta(s)$ is defined by the following asymptotic expression:

$$\Delta(s) = e^{i\pi s} \left\{ 1 + \frac{a_1}{s} + \frac{a_2 - (\beta_1 + \dots + \beta_{n-1}) + \gamma\pi}{s^2} + O\left(\frac{1}{s^3}\right) \right\} +$$

$$+ e^{-i\pi s} \left\{ -1 + \frac{a_1}{s} - \frac{a_2 - (\beta_1 + \dots + \beta_{n-1}) + \gamma\pi}{s^2} + O\left(\frac{1}{s^3}\right) \right\} +$$

$$+ \sum_{k=1}^{n-1} e^{i \frac{\pi k}{n} s} \left\{ \frac{\beta_k + \beta_{n-k}}{s^2} + O\left(\frac{1}{s^3}\right) \right\} - \sum_{k=1}^{n-1} e^{-i \frac{\pi k}{n} s} \left\{ \frac{\beta_k + \beta_{n-k}}{s^2} + O\left(\frac{1}{s^3}\right) \right\} + \left\{ \frac{4i\gamma}{s^3} + O\left(\frac{1}{s^4}\right) \right\}.$$

Here

$$a_1 = -\frac{i}{2} \int_0^\pi q(t) dt, \quad a_2 = \frac{1}{4} \left\{ q(\pi) - q(0) - \frac{1}{2} \left(\int_0^\pi q(t) dt \right)^2 \right\}.$$

Analyzing the equation $\Delta(s) = 0$, we obtain that problem (7), (2) (3) has a $4n$ series of eigenvalues with asymptotics (5) (6). Taking into account that the function $\Delta(s)$ is odd, we get that alongside with the eigenvalues $s_{m,j}$ from (5) the numbers $-s_{m,j}$ are also the roots of a characteristic polynomial. We denoted them by $s_{m,j+2n}, j = \overline{1, 2n}$. Hence we obtain the vanishing of the coefficients $c_{2,j}$ from (6) with the even numbers j . Thus, in terms of the spectral parameter λ we get the $2n$ series of eigenvalues.

The function $\Delta(s)$ belongs to the class K of entire functions of first order [4]. Therefore, for this function one can apply the method of calculating the regularized sum of roots of quasipolynomials based on the construction of the zeta function associated with the function $\Delta(s)$ and using the method of successive approximations of Horn (see. [4]).

Since this technique is well-established, due to cumbersome calculations, we shall not give them here and by this we complete the proof of the theorem.

REMARK 1 In the particular case for $\gamma = 0$, problem (7), (2), (3) coincides with problem (1) - (3) and Eq. (8) of Theorem 2 coincides with the main result of Theorem 1 (formula (4)).

REMARK 2 In the particular case for $\gamma = 0, \beta_k = 0, k = \overline{1, n-1}$, problem (7), (2), (3) coincides with the Dirichlet problem and the main result of Theorem 2 (the formula (8)) coincides with the classical result:

$$\sum_{m=0}^{\infty} \left(\lambda_m - m^2 - \frac{1}{\pi} \int_0^\pi q(t) dt \right) = -\frac{1}{4} (q(0) + q(\pi)) + \frac{1}{2\pi} \int_0^\pi q(t) dt.$$

ACKNOWLEDGMENTS

The authors express their gratitude to T.Sh. Kalmenov, B.E. Kanguzhin and M.A. Sadybekov, this paper has been prepared under their guidance. This research is financially supported by a grant from the Ministry of Science and Education of the Republic of Kazakhstan (Grants No. 0825/GF4 and No. 0824/GF4). This publication is supported by the target program 0085/PTSF-14 from the Ministry of Science and Education of the Republic of Kazakhstan.

REFERENCES

- [1] V.A. Sadovnichii, *Teoriya operatorov* (Drofa, Moscow, 2004). (in russian)
- [2] S.I. Mitrohin, *Spektral'naja teoriya operatorov: gladkie, razryvnye, summiruemye koefitsienty* (Internet-Universitet informacionnyh tehnologij, Moscow, 2009). (in russian)
- [3] I.M. Gel'fand, and B.M. Levitan, *Dokl. Akad. Nauk SSSR* 88:4, 593–596 (1953).
- [4] V.B. Lidskij, and V.A. Sadovnichij, *Funkcional'nyj analiz i ego prilozhenija* 1:2, 52–59 (1967).
- [5] V.V. Karachik, and M.A. Sadybekov, B.T. Torebek, *Electronic Journal of Differential Equations* 2015:244, 1–9 (2015).
- [6] T.Sh. Kal'menov, and D. Suragan, *Differential Equations* 48:3, 441–445 (2012).
- [7] T.Sh. Kal'menov, and D. Suragan, *Differential Equations* 48:4, 604–608 (2012).
- [8] B.E. Kanguzhin, D.B. Nurakhmetov, and N.E. Tokmagambetov, *Ufimsk. Mat. Zh.* 3:3, 80–92 (2010).
- [9] B.E. Kanguzhin, and A.A. Aniyarov, *Mathematical Notes* 89:5-6, 819–829 (2011)
- [10] B.E. Kanguzhin, D.B. Nurakhmetov, and N.E. Tokmagambetov, *Izv. Vyssh. Uchebn. Zaved. Mat.* 2, 9-16 (2014).
- [11] B.E. Kanguzhin, and N.E. Tokmagambetov, *Doklady Mathematics* 91:1, 1–4 (2015).
- [12] B.E. Kanguzhin, and N.E. Tokmagambetov, *Differential Equations* 51:12, 1583–1588 (2015).
- [13] M. Martinovich, *Differenc. uravnenija* 18:3, 537–540 (1982).
- [14] N.S. Imanbaev, and M.A. Sadybekov, ISSN 1563-0285. *Vestnik KazNU, ser. mat., meh., inf.* 2(81), 66–71 (2014).
- [15] T.Sh. Kal'menov, and D. Suragan, *Doklady Mathematics* 78:3, 913–915 (2008).
- [16] T.Sh. Kal'menov, and N.E. Tokmagambetov, *Siberian Mathematical Journal* 54:6, 1023–1028 (2013).
- [17] M.A. Sadybekov, and A.M. Sarsenbi, *Differential Equations* 48:8, 1112–1118 (2012).
- [18] T.Sh. Kal'menov, M.A. Sadybekov, and A.M. Sarsenbi, *Differential Equations* 47:1, 144–148 (2011).
- [19] M.A. Sadybekov, and A.M. Sarsenbi, *Differential Equations* 44:5, 685–691 (2008).
- [20] M.A. Sadybekov, and A.M. Sarsenbi, *Doklady Mathematics* 77:3, 398–400 (2008).
- [21] M.A. Sadybekov, and N.S. Imanbaev, *Differential Equations* 48:6, 896–900 (2012).