

ISSN 2518-1483 (Online),
ISSN 2224-5227 (Print)

2016 • 6

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
БАЯНДАМАЛАРЫ

ДОКЛАДЫ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН

REPORTS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN

ЖУРНАЛ 1944 ЖЫЛДАН ШЫҒА БАСТАҒАН
ЖУРНАЛ ИЗДАЕТСЯ С 1944 г.
PUBLISHED SINCE 1944



Бас редакторы
х.ғ.д., проф., ҚР ҰҒА академигі **М.Ж. Жұрынов**

Редакция алқасы:

Адекенов С.М. проф., академик (Қазақстан) (бас ред. орынбасары)
Боос Э.Г. проф., академик (Қазақстан)
Величкин В.И. проф., корр.-мүшесі (Ресей)
Вольдемар Вуйцик проф. (Польша)
Гончарук В.В. проф., академик (Украина)
Гордиенко А.И. проф., академик (Белорус)
Дука Г. проф., академик (Молдова)
Илолов М.И. проф., академик (Тәжікстан),
Леска Богуслава проф. (Польша),
Локшин В.Н. проф. чл.-корр. (Қазақстан)
Нараев В.Н. проф. (Ресей)
Неклюдов И.М. проф., академик (Украина)
Нур Изура Удзир проф. (Малайзия)
Перни Стефано проф. (Ұлыбритания)
Потапов В.А. проф. (Украина)
Прокопович Полина проф. (Ұлыбритания)
Омбаев А.М. проф. (Қазақстан)
Өтелбаев М.О. проф., академик (Қазақстан)
Садыбеков М.А. проф., корр.-мүшесі (Қазақстан)
Сатаев М.И. проф., корр.-мүшесі (Қазақстан)
Северский И.В. проф., академик (Қазақстан)
Сикорски Марек проф., (Польша)
Рамазанов Т.С. проф., корр.-мүшесі (Қазақстан)
Такибаев Н.Ж. проф., академик (Қазақстан), бас ред. орынбасары
Харин С.Н. проф., академик (Қазақстан)
Чечин Л.М. проф., корр.-мүшесі (Қазақстан)
Харун Парлар проф. (Германия)
Энджун Гао проф. (Қытай)
Эркебаев А.Э. проф., академик (Қырғыстан)

«Қазақстан Республикасы Ұлттық ғылым академиясының баяндамалары»

ISSN 2518-1483 (Online),

ISSN 2224-5227 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» Республикалық қоғамдық бірлестігі (Алматы қ.)
Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде 01.06.2006 ж.
берілген №5540-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.

Тиражы: 2000 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
<http://наука-нанрк.kz>, reports-science.kz

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2016

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Главный редактор
д.х.н., проф., академик НАН РК **М. Ж. Журинов**

Редакционная коллегия:

Адекенов С.М. проф., академик (Казахстан) (зам. гл. ред.)
Боос Э.Г. проф., академик (Казахстан)
Величкин В.И. проф., чл.-корр. (Россия)
Вольдемар Вуйцик проф. (Польша)
Гончарук В.В. проф., академик (Украина)
Гордиенко А.И. проф., академик (Беларусь)
Дука Г. проф., академик (Молдова)
Илолов М.И. проф., академик (Таджикистан),
Леска Богуслава проф. (Польша),
Локшин В.Н. проф. чл.-корр. (Казахстан)
Нараев В.Н. проф. (Россия)
Неклюдов И.М. проф., академик (Украина)
Нур Изура Удзир проф. (Малайзия)
Перни Стефано проф. (Великобритания)
Потапов В.А. проф. (Украина)
Прокопович Полина проф. (Великобритания)
Омбаев А.М. проф. (Казахстан)
Отелбаев М.О. проф., академик (Казахстан)
Садыбеков М.А. проф., чл.-корр. (Казахстан)
Сатаев М.И. проф., чл.-корр. (Казахстан)
Северский И.В. проф., академик (Казахстан)
Сикорски Марек проф., (Польша)
Рамазанов Т.С. проф., чл.-корр. (Казахстан)
Такибаев Н.Ж. проф., академик (Казахстан), зам. гл. ред.
Харин С.Н. проф., академик (Казахстан)
Чечин Л.М. проф., чл.-корр. (Казахстан)
Харун Парлар проф. (Германия)
Энджун Гао проф. (Китай)
Эркебаев А.Э. проф., академик (Кыргызстан)

Доклады Национальной академии наук Республики Казахстан»

ISSN 2518-1483 (Online),

ISSN 2224-5227 (Print)

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №5540-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 2000 экземпляров

Адрес редакции: 050010, г.Алматы, ул.Шевченко, 28, ком.218-220, тел. 272-13-19, 272-13-18

<http://nauka-nanrk.kz> reports-science.kz

©Национальная академия наук Республики Казахстан, 2016 г.

Адрес типографии: ИП «Аруна», г.Алматы, ул.Муратбаева, 75

E d i t o r i n c h i e fdoctor of chemistry, professor, academician of NAS RK **M.Zh. Zhurinov****E d i t o r i a l b o a r d :****Adekenov S.M.** prof., academician (Kazakhstan) (deputy editor in chief)**Boos E.G.** prof., academician (Kazakhstan)**Velichkin V.I.** prof., corr. member (Russia)**Voitsik Valdemar** prof. (Poland)**Goncharuk V.V.** prof., academician (Ukraine)**Gordiyenko A.I.** prof., academician (Belarus)**Duka G.** prof., academician (Moldova)**Ilov M.I.** prof., academician (Tadjikistan),**Leska Boguslava** prof. (Poland),**Lokshin V.N.** prof., corr. member. (Kazakhstan)**Narayev V.N.** prof. (Russia)**Nekludov I.M.** prof., academician (Ukraine)**Nur Izura Udzir** prof. (Malaysia)**Perni Stephano** prof. (Great Britain)**Potapov V.A.** prof. (Ukraine)**Prokopovich Polina** prof. (Great Britain)**Ombayev A.M.** prof. (Kazakhstan)**Otelbayv M.O.** prof., academician (Kazakhstan)**Sadybekov M.A.** prof., corr. member. (Kazakhstan)**Satayev M.I.** prof., corr. member. (Kazakhstan)**Severskiy I.V.** prof., academician (Kazakhstan)**Sikorski Marek** prof., (Poland)**Ramazanov T.S.** prof., corr. member. (Kazakhstan)**Takibayev N.Zh.** prof., academician (Kazakhstan), deputy editor in chief**Kharin S.N.** prof., academician (Kazakhstan)**Chechin L.M.** prof., corr. member. (Kazakhstan)**Kharun Parlar** prof. (Germany)**Endzhun Gao** prof. (China)**Erkebayev A.Ye.** prof., academician (Kyrgyzstan)**Reports of the National Academy of Sciences of the Republic of Kazakhstan.****ISSN 2224-5227****ISSN 2518-1483 (Online),****ISSN 2224-5227 (Print)**

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 5540-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 2000 copies

Editorial address: 28, Shevchenko str., of.219-220, Almaty, 050010, tel. 272-13-19, 272-13-18,

<http://nauka-nanrk.kz/> reports-science.kz

© National Academy of Sciences of the Republic of Kazakhstan, 2016

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

T.S. Ramazanov¹, S.K. Kodanova¹, M.K. Issanova¹, A.Tikhonov², M. Kaikanov²

¹IETP, Al-Farabi Kazakh National University, Almaty, Kazakhstan;

²National Laboratory Astana, Nazarbayev University, Astana, Kazakhstan
isanova_moldir@mail.ru

TRANSPORT PROPERTIES OF INERTIAL CONFINEMENT FUSION PLASMAS

Abstract. In this paper the transport properties of non-isothermal dense deuterium-tritium plasmas were studied. Based on the effective interaction of potentials between particles, the Coulomb logarithm for a two-temperature nonisothermal dense plasma was obtained. These potentials take into consideration long-range multi-particle screening effects and short-range quantum-mechanical effects in two-temperature plasmas. Transport processes in such plasmas were studied using the Coulomb logarithm. The obtained results were compared with the theoretical works of other authors and with the results of molecular dynamics simulations.

Keywords: dense plasma, Coulomb logarithm, inertial confinement fusion, transport properties.

Introduction. Nowadays, much attention is paid to the high density energy of the substance and matters at high pressures and temperatures. Research in the field of fusion power with inertial confinement (ITS) in the heavy ion beams have a special significance among the works devoted to various aspects of this problem. Basically, these heavy ion accelerators are well known as the main tool in experimental studies of nuclear physics, elementary particle physics and dense plasma physics [1-3]. However, at present time, the lack of new theoretical and experimental data on the transport properties of deuterium-tritium (DT) plasma produced under compression of target with heavy ions beam, requires an adequate qualitative description of the interaction of heavy ions with a dense plasma in a wide range of parameters and provides an additional impetus to the research in this area. Understanding and control of high-pressure behavior of DT fuel are crucial for the success of the experiments with the ignition. Accurate knowledge of transfer coefficients in dense DT plasma is essential for the correct description of the processes occurring in the inertial confinement. This issue has been the subject of many theoretical and experimental studies [4-7]. In some researches [8], the transport properties of dense plasmas, such as diffusion and viscosity were studied on the basis of molecular dynamics simulation (MD) using density functional theory to describe the electron component of the plasma (DFTMD). Kress and others [9] obtained values of viscosity and diffusion of DT plasma, using a density functional theory at finite temperatures on the basis of Kohn-Sham theorem on molecular dynamics and DFTMD.

One of the most promising approaches for the study of transport properties of dense DT plasma is a binary collision approximation, and we can use two approaches. One of them is to calculate the transport coefficients determined on the basis of the particles scattering which interact via potential. The second approach is producing kinetic equation, which collision integral has logarithmically divergent integral on impact parameters, which can be replaced by the Coulomb logarithm.

In this paper, previously proposed model [17, 18, 19] for description of the dense plasma properties on the basis of effective interaction potentials [20, 21] is extended for calculation of the ion transport properties and thermal conductivity for deuterium and deuterium-tritium plasma of inertial confinement. Below we demonstrate a brief description of the model and the results of calculation of plasma transport

properties. In order to show the correctness of the model, its results are compared with the results of quantum molecular dynamics of KMD and TFTMD modelling.

Coulomb logarithm on the basis of the effective potential. Transport properties are obtained from on the basis of the Coulomb logarithm using the effective potential for inertial confinement plasma. Coulomb logarithm is determined by the scattering angle in the center-of-mass system with the pair Coulomb collisions [15-17]:

$$\lambda_{ei} = \frac{1}{b_{\perp}^2} \int_0^{b_{\max}} \sin^2\left(\frac{\theta_c}{2}\right) b db, \quad (1)$$

The scattering angle in the center-of-mass system θ_c is defined as [15]:

$$\theta_c = \pi - 2b \int_{r_0}^{\infty} \frac{dr}{r^2} \left(1 - \frac{\Phi_{\alpha\beta}(r)}{E_c} - \frac{b^2}{r^2} \right)^{1/2}, \quad (2)$$

where $E_c = \frac{1}{2} m_{\alpha\beta} v^2$ - is energy of center-of-mass system, $m_{\alpha\beta} = m_{\alpha} m_{\beta} / (m_{\alpha} + m_{\beta})$ - resulted mass of the particles of α and β (ion or electron); $b_{\perp} = Z_{\alpha} Z_{\beta} / (m_{\alpha\beta} v^2)$. As a minimum impact parameter there was $b_{\min} = \max\{b_{\perp}, \lambda_{\alpha\beta}\}$, where $\lambda_{\alpha\beta} = \hbar / \sqrt{2\pi m_{\alpha\beta} k_B T}$ - thermal wavelength de Broglie.

The following dimensionless variables, such as the connection parameter are used:

$$\Gamma_{ee} = \frac{e^2}{ak_B T_e}, \quad \Gamma_{ii} = \frac{Z_i^2 e^2}{ak_B T_i} \left(\frac{n_i}{n_e} \right)^{1/3}, \quad \Gamma_{ei} = \frac{Z_i e^2}{ak_B T_{ei}}, \quad (3)$$

where e - electron charge, $a = (3/4\pi n_e)^{1/3}$ - the average interparticle distance between the particles, k_B - Boltzmann constant. In the formula (2) $\Phi_{\alpha\beta}(r)$ - the potential of interaction between the particles and the distance of closest approach r_0 for a given impact parameter b is determined from the equation:

$$1 - \frac{\Phi_{\alpha\beta}(r_0)}{E_c} - \frac{b^2}{r_0^2} = 0. \quad (4)$$

As it is well known, calculation of collective screening effects in the interaction of plasma particles is essential for the correct description of the static and dynamic properties of the plasma. In this paper the dense plasma is considered, which is also important for the quantum effects at small interparticle distances. In addition, we will use electron-ion effective potential, which takes into account both the quantum effects at small distances, and the effect of the screening - at large distances [18-19]:

$$\Phi_{\alpha\beta}(r) = \frac{Z_{\alpha} Z_{\beta}}{r} \frac{1}{\gamma^2 \sqrt{1 - (2k_D / \lambda_{ee} \gamma^2)^2}} \left(\left(\frac{1/\lambda_{ee}^2 - B^2}{1 - B^2 \lambda_{\alpha\beta}^2} \right) \exp(-Br) - \left(\frac{1/\lambda_{ee}^2 - A^2}{1 - A^2 \lambda_{\alpha\beta}^2} \right) \exp(-Ar) \right) - \frac{Z_{\alpha} Z_{\beta} e^2 (1 - \delta_{\alpha\beta})}{r (1 + C_{\alpha\beta})} \exp(-r/\lambda_{\alpha\beta}), \quad (5)$$

here

$$A^2 = \frac{\gamma^2}{2} \left(1 + \sqrt{1 - \left(\frac{2k_D}{\lambda_{ee} \gamma^2} \right)^2} \right), \quad B^2 = \frac{\gamma^2}{2} \left(1 - \sqrt{1 - \left(\frac{2k_D}{\lambda_{ee} \gamma^2} \right)^2} \right),$$

$$C_{\alpha\beta} = \frac{k_D^2 \tilde{\lambda}_{\alpha\beta}^2 - k_i^2 \tilde{\lambda}_{ee}^2}{\tilde{\lambda}_{ee}^2 / \tilde{\lambda}_{\alpha\beta}^2 - 1},$$

where $2k_D / (\tilde{\lambda}_{ee} \gamma^2) < 1$, $k_D^2 = k_e^2 + k_i^2$ - screening parameter that takes into account the electrons and ions, $\gamma^2 = k_i^2 + 1/\tilde{\lambda}_{ee}^2$. For non-isothermal plasma we use electron-ion characteristic temperature T_{ei} [22-23]. In the research [22] it is shown, that for the correct description of the properties of the plasma electron-ion temperature should be expressed in the form of: $T_{ei} = \sqrt{T_e T_i}$. These effective potentials can be used for non-isothermal and isothermal plasma.




Figure 1 – Coulomb logarithm of the two-temperature DT plasma depending (a) on of the coupling parameters (Γ) and (b) on the parameter of degeneration (Θ).

In the study of the transport properties of dense high-temperature plasma, the values of the Coulomb logarithm are very vital. This paper presents the study of the transport properties of dense plasma on the basis of Coulomb logarithm using the effective potential [5].

Figure 1 shows the calculated values of the Coulomb logarithm, depending on the coupling parameter Γ and the degeneracy parameter $\Theta = k_B T / E_F$ (E_F Fermi energy) at different ratios of the electron and ion temperatures in the dense DT two-temperature plasma. We considered the loosely coupled plasma, in Fig. 1 a) the values of the Coulomb logarithm for $\Gamma_{ee} < 1$ and $\Gamma_{ii} < 1$ were shown. The value of the Coulomb logarithm decreases with increasing of T_e / T_i ratio. At a given electrons temperature T_e , the lower values of the Coulomb logarithm at higher values T_e / T_i are the result of strong screening of the ionic component of the plasma. Fig. 1 b) shows that the increase in the degeneracy parameter leads to increase of the value of the Coulomb logarithm. At constant density we have the lowest values of the screening length for higher temperatures. This leads to higher values of the Coulomb logarithm.

Transport properties of inertial confinement dense plasma. The phenomenon of transfer in a dense plasma is of considerable interest in the various areas of science and technology (plasma physics, inertial confinement, physics of hot dense matter, etc.) [24-25]. In particular, intense studies of fusion power require more reliable information on the transfer of coefficients, i.e, thermal conductivity coefficient, diffusion and viscosity. We will consider the dense DT plasma particles interacting through the effective potential (5).

The coefficient of diffusion, viscosity, and thermal conductivity of the plasma are connected with the effective collision rate using equations:

$$D = \frac{k_B T}{m_e \nu_{eff}}, \tag{6}$$

$$\eta = \frac{5}{4} \sqrt{\frac{m}{\pi}} \frac{(k_B T)^{5/2}}{e^4 \lambda}, \tag{7}$$

$$\kappa = \frac{5n_e k_B^2 T}{m_e \nu_{eff}}, \tag{8}$$

where e - electron charge, m_e - electron mass, n - plasma particle density, and

$$\nu_{eff} = (4/3) \sqrt{2\pi} e^4 \lambda / \sqrt{m_e} (k_B T)^{3/2} \tag{9}$$

effective collision frequency is directly proportional to the logarithm of the Coulomb. The diffusion coefficients of D and viscosity η are given in the following form: $D^* = D/\omega_p a^2$ and $\eta^* = \eta/n_i M \omega_p a^2$, $\kappa^* = \kappa/(m_e \omega_p/a)$, where $\omega_p = (4\pi n_i/M)^{1/2} Z e$ - the plasma frequency for the mass of M ions. In this paper, we use $M = (2 + 3)/2 = 2.5 \text{ amu}$ [26] for considering DT.




Figure 2 – The diffusion coefficient, depending on the coupling parameter (Γ), $T_e = T_i$: (a) for different density parameter values (r_s), (b) for different ratios of the electron and ion temperatures at $r_s = 1$.




Figure 3 – The coefficients (a) of thermal conductivity and (b) viscosity depending on the coupling parameter (Γ) for different values of the parameter of density

$$(r_s), T_e = T_i.$$

Figures 2 a), 2 b) and 3 a), 3 b) show the results of diffusion, thermal conductivity and viscosity coefficient of dense plasma depending on the coupling parameter at $r_s = 0.5$, $r_s = 1$ and $r_s = 1.5$. In order to have a more detailed physical description of processes, obtained by numerical methods, the results will be calculated on the basis of the effective interaction potential (5), which takes into account both the quantum and the collective effects. Figures 2 a) and 3 a), 3 b) shows that at lower values of coupling parameter, the coefficients of viscosity and diffusion have higher values. For higher density, the transfer coefficients have lower values.

The coefficients of diffusion, thermal conductivity and viscosity at different temperatures and densities on the basis of the Coulomb logarithm using an effective potential (5) are obtained. Figures 4 a) and 4 b) show the coefficients of diffusion and viscosity of the dense plasma DT calculated on the basis of the Coulomb logarithm depending on the coupling parameter (Γ) at the plasma density, $\rho = 6.135 \text{ g/cm}^3$, $\rho = 13.45 \text{ g/cm}^3$, $\rho = 26.3 \text{ g/cm}^3$, $\rho = 108 \text{ g/cm}^3$, respectively. It is evident that coefficients of diffusion and viscosity increase with increasing temperature.




Figure 4 – Transport coefficients of dense DT plasma depending on the temperature for different densities: (a) diffusion, (b) the viscosity

Figures 5 a) and 5 b) show the results of thermal conductivity of deuterium plasma from temperature for different values of the density $\rho = 43.105 \text{ g/cm}^3$ and $\rho = 199.561 \text{ g/cm}^3$. Solid red line is the thermal conductivity, obtained on the basis of the effective interaction potential (5), the black triangles are KMD modeling results [27]. Blue dot-dashed line is the total Coulomb logarithm $\lambda = \ln \Lambda$. In the paper [27], it was estimated KMD modeling of thermal conductivity of deuterium plasma in a wide range of densities and temperatures.. S.X. Hu and others [27] used the following function to describe the results of calculations of KMD modeling of deuterium thermal conductivity on inertial confinement explosions:

$$\kappa_{QMD} = \frac{20(2/\pi)^{3/2} k_B^{7/2} T^{5/2}}{\sqrt{m_e} Z_{eff} e^4} \frac{0.095(Z_{eff} + 0.24)}{1 + 0.24Z_{eff}} \frac{1}{\ln \Lambda_{QMD}}. \quad (10)$$




Figure 5 – Thermal conductivity of deuterium plasma for effective interaction potential (5) and KMD modeling depending on the temperature at $\rho = 43.105 \text{ g/cm}^3$ and $\rho = 199.561 \text{ g/cm}^3$.

Figures 5 a) and 5 b) clearly show, that the more temperature increases the more thermal conductivity increases. We should note that at high values of density, the result obtained on the basis of the effective potential at low temperatures approaches to the result of quantum molecular dynamics method.




Figure 6 – Diffusion (a) and the viscosity (b) of dense plasma depending on coupling parameter (Γ), $T_e = T_i$.

Figures 6 a) and 6 b) show the dependence of the diffusion coefficient and the viscosity of the coupling parameter in comparison with the results obtained on the basis of the hyperchaed approximation (HCA) [28], the method of molecular dynamics (MD) [28-30], in the framework of the kinetic theory [33-34], as well as the Spitzer-Landau theory [31]. Daligalta-Balruda theory is based on approximation of pair scattering taking into account the correlation effects on the basis of the use of effective interaction potential [28-30, 32]. The effective potential of Daligalta-Balruda is connected with the mean-field potential, which is included in the pair interaction potential. Wallenborn-Baus used the renormalized kinetic theory and the generalized kinetic theory of correlation functions in phase space [34]. Figures 6 a) and 6 b) show that the results obtained on the basis of the effective potential (5) are well correspond with the results of other studies in the weakly bound limit $\Gamma_{ee} < 1$, but vary at $\Gamma_{ee} \geq 1$. The difference in weakly connected case $\Gamma_{ee} \sim 1$ is caused by non-ideal and quantum effects.

We calculated the diffusion and viscosity of DT plasma for density $\rho = 5 \text{ g/cm}^3$ and temperature in the range of $2 - 10 \text{ eV}$ using the Coulomb logarithm on the basis of the effective potential, taking into account the quantum diffraction effect at short distances and the effect of screening at large distances. Figures 7 a) and 7 b) show the comparison of calculation data on diffusion and viscosity in the DT plasma with the theoretical results of other authors [9], calculated on the basis of the density functional theory at finite temperatures using Kohn-Sham theorem in combination with molecular dynamics and functional theory density without the exchange term to describe the electron component of the plasma (DFTMD). The results are well correspond with the results of KMD and DFTMD modeling at higher temperatures, and therefore, we conclude that our method can be used in this way. At below 3 eV temperatures, comparison with KMD and DFTMD results shows deterioration of correspondence, since at these temperatures, the ideality effect becomes important. Compared with the results of KMD, obtained viscosity data are not as good as for the diffusion, where the temperature dependence differs significantly, while the results obtained for the viscosity on the basis of the effective potential corresponds to the DFTMD modeling results.




Figure 7 – Diffusion (a) and viscosity (b) of dense DT plasma depending on the temperature at $\rho = 5.0 \text{ g/cm}^3$.

Conclusion. The study of transport properties in dense DT plasma on the basis of two-temperature effective interaction potential, which takes into account the quantum effects of diffraction at small distances and screening at large distances has been conducted. The results obtained using Coulomb logarithm and transfer coefficients for various plasma parameters are correspond with the theoretical and experimental results of other authors, and with the results of MD modeling. According to the results it is clear that the transport properties of dense plasma can be adequately expressed in terms of the Coulomb logarithm based on the effective potentials. Thus, the knowledge of coefficient values of transfer of heavy, charged particles in the plasma will allow obtaining more accurate calculation of the construction of a thermonuclear target.

This work was supported by the Ministry of Education and Science of the Republic of Kazakhstan under the grant №0115RK03029 (2016).

REFERENCES

- [1] V.E. Fortov (2009) Extreme states of matter on Earth and in the Cosmos. Springer. Germany. ISBN: 978-3-642-16463-7 (in English)
- [2] V.E. Fortov, O.F. Petrov, O.S. Vaulina, R.A. Timirkhanov (2012) Viscosity of a Strongly Coupled Dust Component in a Weakly Ionized Plasma, Phys. Rev. Lett. 109: 055002. DOI: 10.1103/PhysRevLett.109.055002 (in English)

- [3] V. E. Fortov, A.V. Ivlev, S. A. Khrapak, A. G. Khrapak and G. E. Morfill (2005) Complex (dusty) plasmas: Current status, open issues, perspectives. *Phys. Rep.* 1: 424. DOI: 10.1016/j.physrep.2005.08.007 (in English)
- [4] D. H. H. Hoffmann, A. Blazevic, P. Ni et al (2005) Present and future perspectives for high energy density physics with intense heavy ion and laser beams. *Laser and Particle beams.* 23: 47. DOI: 10.1017/S026303460505010X (in English)
- [5] D. H. H. Hoffmann, K. Weyrich, H. Wahl et al (1990) Energy loss of heavy ions in a plasma target. *Phys. Rev. A.* 42: 2313. DOI: 10.1103/PhysRevA.42.2313 (in English)
- [6] D. H. H. Hoffmann, J. Jacoby, W. Laux et al (1995) Stopping of Heavy Ions in a Hydrogen Plasma. *Phys. Rev. Lett.* 74: 1550. DOI: 10.1103/PhysRevLett.74.1550 (in English)
- [7] C.-V. Meister, G. Ropke (1982) Electrical conductivity of non-ideal plasmas and the ion distribution function. *Annalen der Physik.* 39: 133. DOI: 10.1002/andp.19824940208 (in English)
- [8] L. Burakovsky, C. Ticknor, J.D. Kress, L.A. Collins, F. Lambert (2013) Transport properties of lithium hydride at extreme conditions from orbital-free molecular dynamics. *Phys. Rev. E.* 87: 023104. DOI: 10.1103/PhysRevE.87.023104 (in English)
- [9] J.D. Kress, J.S.Cohen, D.A. Horner, F. Lambert, L.A. Collins (2010) Viscosity and mutual diffusion of deuterium-tritium mixtures in the warm-dense-matter regime. *Phys. Rev. E.* 82: 036404. DOI: 10.1103/PhysRevE.82.036404 (in English)
- [10] R. E. Phillips and C. A. Ordonez (2013) Accelerator-based neutron source using a cold deuterium target with degenerate electrons. *AIP Advances.* 3: 072115. DOI: 10.1063/1.4816407 (in English)
- [11] Y. Chang, C. A. Ordonez (2000) Velocity space scattering coefficients with applications in antihydrogen recombination studies. *Physical Review E.* 62: 8564. DOI: 10.1103/PhysRevE.62.8564 (in English)
- [12] D.O. Gericke, M.S. Murillo, M. Schlages (2002) Dense plasma temperature equilibration in the binary collision approximation. *Phys. Rev. E.* 65: 036418. DOI: 10.1103/PhysRevE.65.036418 (in English)
- [13] K.Chen, S. T. Sullivan, W.G. Rellergert, E.R. Hudson (2013) Measurement of the Coulomb Logarithm in a Radio-Frequency Paul Trap. *Phys. Rev. Lett.* 110: 173003. DOI: 10.1103/PhysRevLett.110.173003 (in English)
- [14] J. Vorberger, D.O. Gericke (2014) Comparison of electron–ion energy transfer in dense plasmas obtained from numerical simulations and quantum kinetic theory. *High Energy Density Physics.* 10: 1. DOI: 10.1016/j.hedp.2013.10.006 (in English)
- [15] C.A. Ordonez, M.I. Molina (1994) Evaluation of the Coulomb logarithm using cutoff and screened Coulomb interaction potentials. *Phys. Plasmas.* 1: 2515. DOI: 10.1063/1.870578 (in English)
- [16] T.S. Ramazanov, S.K. Kodanova (2001) Coulomb logarithm of a nonideal plasma. *Phys. Plasmas.* 8: 5049. DOI: 10.1063/1.1407820 (in English)
- [17] T. S. Ramazanov, S. K. Kodanova, Zh. A. Moldabekov, M. K. Issanova (2013) Dynamical properties of non-ideal plasma on the basis of effective potentials. *Phys. Plasmas.* 20:112702. DOI: 10.1063/1.4829042 (in English)
- [18] T. S. Ramazanov, Zh. A. Moldabekov, M. T. Gabdullin (2015) Effective potentials of interactions and thermodynamic properties of a nonideal two-temperature dense plasma. *Phys.Rev. E.* 92: 023104. DOI: 10.1103/PhysRevE.92.023104 (in English)
- [19] S.K. Kodanova, T.S. Ramazanov, M.K. Issanova, Zh.A. Moldabekov, G. Nigmatova (2015) Investigation of Coulomb Logarithm and Relaxation Processes in Dense Plasma on the Basis of Effective Potentials. *Contrib. Plasma Phys.*55, 2-3: 271 – 276. DOI: 10.1002/ctpp.201400094 (in English)
- [20] Zhandos Moldabekov, Tim Schoof, Patrick Ludwig, Michael Bonitz, and Tlekkabul Ramazanov (2015) Statically screened ion potential and Bohm potential in a quantum plasma. *Phys. Plasmas.* 22:102104. DOI: 10.1063/1.4932051 (in English)
- [21] T. S. Ramazanov, Zh. A. Moldabekov, M. T. Gabdullin (2016) Interaction between ions in hot dense plasma via screened Cornell potential. *Phys. Plasmas.* 23: 042703. DOI: 10.1063/1.4945648 (in English)
- [22] P. Seufferling, J. Vogel, and C. Toepffer (1989) Correlations in a two-temperature plasma. *Phys. Rev. A.* 40: 323-329. DOI: 10.1103/PhysRevA.40.323 (in English)
- [23] R. Bredow, Th. Bornath, W.D. Kraeft, R. Redmer (2013) Hypernetted Chain Calculations for Multi-Component and NonEquilibrium Plasmas. *Contrib. Plasma Phys.*53: 276-284. DOI: 10.1002/ctpp.201200117 (in English)
- [24] S. Atzeni and J. Meyer-ter-Vehn (2004) *The Physics of Inertial Fusion: Beam Plasma Interaction, Hydrodynamics, Hot Dense Matter*, International Series of Monographs on Physics. Clarendon, Oxford. ISBN: 978-0-19-956801-7 (in English)
- [25] J. D. Lindl (1998) *Inertial Confinement Fusion: The Quest for Ignition and Energy Gain Using Indirect Drive*. Springer, Verlag, New York. ISBN: 978-3-642-16463-7 (in English)
- [26] C. Wang, Y. Long, X.-T. He, J.-F. Wu, W.-H. Ye, and P. Zhang (2013) Transport properties of dense deuterium-tritium plasmas. *Phys. Rev. E.* 88: 013106. DOI: 10.1103/PhysRevE.88.013106 (in English)
- [27] S.X. Hu, L.A. Collins, T.R. Boehly, J.D. Kress et al (2014) First-principles thermal conductivity of warm-dense deuterium plasmas for inertial confinement fusion applications. *Phys. Rev. E.* 89:043105. DOI: 10.1103/PhysRevE.89.043105 (in English)
- [28] J. Daligault and S.D. Baalrud (2013) Effective Potential Theory for Transport Coefficients across Coupling Regimes. *Phys. Rev. Lett.* 110:235001. DOI: 10.1103/PhysRevLett.110.235001 (in English)
- [29] J. Daligault (2012) Diffusion in Ionic Mixtures across Coupling Regimes. *Phys. Rev. Lett.*108:225004. DOI: 10.1103/PhysRevLett.108.225004 (in English)
- [30] J. Daligault (2012) Practical model for the self-diffusion coefficient in Yukawa one-component plasmas. *Phys. Rev. E.* 86: 047401. DOI: 10.1103/PhysRevE.86.047401 (in English)
- [31] L. Spitzer, Jr. (1962) *Physics of Fully Ionized Gases*, 2nd ed. Interscience, New York. ISBN: 978-0-486-44982-1 (in English)
- [32] J. Daligault and K. Rasmussen, S.D. Baalrud (2014) Determination of the shear viscosity of the one-component plasma. *Phys. Rev. E.* 90: 033105. DOI: 10.1103/PhysRevE.90.033105 (in English)

[33] S. Tanaka and S. Ichimaru (1986) Theory of interparticle correlations in dense, high-temperature plasmas. VIII. Shear viscosity. Phys. Rev. A. 34: 4163. DOI: 10.1103/PhysRevA.34.4163 (in English)

[34] J. Wallenborn and M. Baus (1978) Kinetic theory of the shear viscosity of a strongly coupled classical one-component plasma. Phys. Rev. A. 18: 1737. DOI: 10.1103/PhysRevA.18.1737 (in English)

Т.С. Рамазанов¹, С.К. Коданова¹, М.К. Исанова¹, А.Тихонов², М. Кайканов²

¹ЭТФГЗИ, әл-Фараби ат. Қазақ Ұлттық университеті, Алматы, Қазақстан;

²Астана Ұлттық Лабораториясы, Назарбаев Университеті, Астана, Қазақстан
isanova_moldir@mail.ru

ИНЕРЦИЯЛЫҚ ТЕРМОЯДРОЛЫҚ СИНТЕЗ ПЛАЗМАСЫНЫҢ ТРАНСПОРТТЫҚ ҚАСИЕТТЕРІ

Аннотация. Бұл жұмыста изотермиялық емес, тығыз дейтерий-третий плазмасының транспорттық қасиеттері зерттелді. Бөлшектердің әсерлесуінің эффективті потенциалы негізінде екі температуралы, изотермиялық емес, тығыз плазма үшін Кулон логарифмі алынды. Бұл потенциал екі температуралы плазмада кіші ара-қашықтықта квантты-механикалық дифракция әсерін, үлкен ара-қашықтықта экрандау әсерін ескереді. Кулон логарифмі көмегімен изотермиялық емес тығыз плазманың тасымалдау коэффициенттері зерттелді. Алынған нәтижелер басқа авторлардың теориялық жұмыстарымен, молекулалық динамика модельдеу нәтижелерімен салыстырылған.

Түйін сөздер: тығыз плазма, Кулон логарифмі, инерциялық термоядролық синтез, транспорттық қасиеттер.

УДК 533.93

Т.С. Рамазанов¹, С.К. Коданова¹, М.К. Исанова¹, А.Тихонов², М. Кайканов²

¹НИИЭТФ, Казахский национальный университет им. Аль-Фараби, Алматы, Казахстан;

²Национальная Лаборатория Астана, Назарбаев Университет, Астана, Казахстан
isanova_moldir@mail.ru

ТРАНСПОРТНЫЕ СВОЙСТВА ПЛАЗМЫ ИНЕРЦИОННОГО ТЕРМОЯДЕРНОГО СИНТЕЗА

Аннотация. В данной работе исследованы транспортные свойства неизотермической, плотной дейтерий-третиевой плазмы. На основе эффективного потенциала взаимодействия частиц для двухтемпературной, неизотермической, плотной плазмы был получен кулоновский логарифм. Данный потенциал учитывает квантово-механические эффекты дифракции на малых расстояниях и эффекты экранировки - на больших в двухтемпературной плазме. С помощью кулоновского логарифма исследованы коэффициенты переноса в неизотермической плотной плазме. Полученные результаты сравнены с теоретическими работами других авторов и результатами моделирования молекулярной динамики.

Ключевые слова: плотная плазма, кулоновский логарифм, инерционный термоядерный синтез, транспортные свойства

RAMAZANOV T.S., chief researcher, Doctor of science, Professor, Member - correspondent NAN RK, IETP, Al-Farabi Kazakh National University, Almaty, Kazakhstan

KODANOVA S.K. leading researcher, Ph.D., Professor, IETP, Al-Farabi Kazakh National University, Almaty, Kazakhstan

ISSANOVA M.K. researcher, IETP, Al-Farabi Kazakh National University, Almaty, Kazakhstan

TIKHONOV A. leading researcher, Ph.D., National Laboratory Astana, Nazarbayev University, Astana, Kazakhstan

KAIKANOV M. leading researcher, Ph.D., National Laboratory Astana, Nazarbayev University, Astana, Kazakhstan

МАЗМҰНЫ
Химия

Шадин Н.А., Anderson J. A., Закарина Н.А., Волкова Л.Д. Ауыр вакуумдық газойль крекингіндегі монтмориллонитте алюминиймен пилларирленген цеолитқұрамды (HY+HZSM-5) катализатор..... 5

Әлеуметтік ғылымдар

Құрманов Н.А., Сатбаева А.Ж., Рахимбекова А.Е., Махатова А.Б. Адами потенциалының даму индексі: заманауи әлемдегі Қазақстанның орны..... 14

Панзабекова А.Ж., Турабаев Г.К. Система оплаты и стимулирования труда на предприятиях реального сектора Казахстана..... 20

Турабаев Г.К., Несіпбеков Е.Н. Білім беру ұйымдарының даму процесінің құрамдасы ретіндегі кадрлық резервпен жұмыс жасау.....27

Физика

Рамазанов Т.С., Коданова С.К., Исанова М.К., Тихонов А., Кайканов М. Инерциялық термоядролық синтез плазмасының транспорттық қасиеттері 34

Рамазанов Т.С., Коданова С.К., Исанова М.К., Тихонов А., Кайканов М. Инерциялық термоядролық синтез плазмасының транспорттық қасиеттері..... 43

Химия

Малышев В.П., Зубрина Ю.С., Макашева А.М. Өзіндік ұйымдастыру үрдістерінің түсінігінде Больцман-Шеннон энтропиясының рөлі..... 53

Комекова Н.М., Козлов В.А., Жұрынов М.Ж. Қара сланецтен ванадийді күкүрт қышқылды атмосферлі-автоклавты шаймалау 62

Баешов А.Б., Адайбекова А.А., Абдувалиева У.А. Молибден электродының натрий гидроксиді ерітіндісіндегі электрохимиялық қасиеті..... 70

Жер туралы ғылымдар

Метакса Г.П., Чекушина Т.В., Молдабаева Г.Ж., Метакса А.С. Байкал көлі – көмірсутектердің табиғи реакторы.. 77

Биология

Хакімжанов А.А., Мамытова Н.С., Бескемпірова Ж.Д., Тілеген Б., Дәлелханқызы А., Кузовлев В.А., Айтхожина Н.Ө. Бидайдың хитиназалық кешені және оның кейбір қасиеттері 85

Техникалық ғылымдар

Жирнова О.В., Тойгожинова А.Ж., Жакипов Ж., Туриканов Т.С., Оразалин А., Матенов Н. Парниктік газдар шығарындыларын азайту автоматтандырылған басқару жүйесін жану процесі биогаз математикалық модельдерін әзірлеу..... 94

Калдыбаева Б.М., Хусанов А.Е., Дмитриев Е.А., Сабырханов Д.С., Абылмағжанов А.З. Хемосорбциялық аппаратта гидродинамикалық жағдайды және конструкциялық ерекшеліктерін ескере отырып ағындардың құрамын есептеу.....106

Қоғамдық ғылымдар

Аюпова З.К., Құсайынов Д.Ө. Қазақ мәдениетіндегі білім беру жүйесінің бастаулары мәселесіне 115

Даулетбақов Б.Д., Примжарова К.К., Қонырбеков М.Ж. Қазақстан республикасы өңірлерінде интеллектуалды әлеуетпен қамтамасыз ету саласындағы инновациялық қызмет деңгейін модельдеу және бағалау..... 122

Қалдыбай Қ.Қ., Абдрасилов Т.Қ., Насимов М.Ө. Заманауи қазақ ойшылдарының дінтанулық мұраларындағы адам мәселесі..... 131

Магай Т.П. Трансформация бизнес-білім беру: инновациялық тәсіл..... 141

Мырзағалиева А.Б., Туктасинова А.А., Самарханова Т.Н., Акзамбек А.М. Алтай қасқыржидегін (*Darphnealtaicarpall.*) *In vitro* мәдениетіне енгізу..... 151

Ордабаева М. Қазақстандағы емдік-сауықтыру туризм үрдістерінің мәселелер мен ағымдары.....161

Саткалиева Т.С. Қазақстан энергетикалық секторының даму үрдістері 167

Берік А.Б. Психикалық дамуы тежелген балаларды психологиялық қолдау..... 176

СОДЕРЖАНИЕ

Химия

Шадин Н.А., Anderson J. A., Закарина Н.А., Волкова Л.Д. Цеолитсодержащий (HY+HZSM-5) катализатор на пилларированном Al - монтмориллоните в крекинге утяжеленного вакуумного газойля..... 5

Социальные науки

Курманов Н.А., Сатбаева А.Ж., Рахимбекова А.Е., Махатова А.Б. Индекс развития человеческого потенциала: место Казахстана в современном мире..... 14

Панзабекова А.Ж., Турабаев Г.К. Система оплаты и стимулирования труда на предприятиях реального сектора Казахстана..... 20

Турабаев Г.К., Несипбеков Е.Н. Работа с кадровым резервом как составляющая процесса развития образовательных организаций..... 27

Физика

Рамазанов Т.С., Коданова С.К., Исанова М.К., Тихонов А., Кайканов М. Транспортные свойства плазмы инерционного термоядерного синтеза..... 34

Рамазанов Т.С., Коданова С.К., Исанова М.К., Тихонов А., Кайканов М. Транспортные свойства плазмы инерционного термоядерного синтеза..... 43

Химия

Малышев В.П., Зубрина Ю.С., Макашева А.М. Роль энтропии Больцмана-Шеннона в понимании процессов самоорганизации..... 53

Комекова Н.М., Козлов В.А., Журинов М.Ж. Серноокислотное атмосферно-автоклавное выщелачивание ванадия из черных сланцев..... 62

Баешов А.Б., Адайбекова А.А., Абдувалиева У.А. Электрохимическое поведение молибденового электрода в растворе гидроксида натрия..... 70

Науки о Земле

Метакса Г.П., Чекушина Т.В., Молдабаева Г.Ж., Метакса А.С. Озеро Байкал – природный реактор углеводородов..... 77

Биология

Хакимжанов А.А., Мамытова Н.С., Бескемпирова Ж.Д., Тилеген Б., Далелханкызы А., Кузовлев В.А., Айтхожина Н.А. Хитиновый комплекс пшеницы и некоторые его свойства..... 85

Технические науки

Жирнова О.В., Тойгожинова А.Ж., Жакипов Ж., Туриканов Т.С., Оразалин А., Матенов Н. Разработка математической модели для автоматизированной системы управления процессом сжигания биогаза для снижения выброса парниковых газов..... 94

Калдыбаева Б.М., Хусанов А.Е., Дмитриев Е.А., Сабырханов Д.С., Абиьмагжанов А.З. Расчет структуры потоков с учетом гидродинамической обстановки и конструктивных особенностей в хемосорбционном аппарате..... 106

Общественные науки

Аюпова З.К., Кусаинов Д.У. К вопросу об истоках системы образования в казахской культуре..... 115

Даулетбаков Б.Д., Примжарова К.К., Конарбеков М.Ж. Моделирование и оценка уровня инновационной деятельности в сфере обеспечения интеллектуальным потенциалом регионов республики Казахстан..... 122

Калдыбай К. К., Абдрасилов Т.К., Насимов М.О. Проблема человека в религиозном наследии современных казахских мыслителей..... 131

Магай Т.П. Трансформация бизнес-образования: инновационный подход..... 141

Мырзагалиева А.Б., Туктасинова А.А., Самарханов Т.Н., Акзамбек А.М. Введение в культуру *In vitro* волчегодника Алтайского (*Daphnealtaicapall*)..... 151

Ордабаева М. Современные проблемы и тенденции развития лечебно-оздоровительного туризма в Казахстане..... 161

Саткалиева Т.С. Тенденции развития энергетического сектора РК..... 167

Берик А.Б. Психологическое сопровождение детей с задержкой психического развития..... 176

CONTENT

Chemistry

- Shadin N.A., Anderson J. A., Zakarina N. A., Volkova L.D.* Zeolite containing (HY+HZSM-5) catalyst on pillared Al-montmorillonite for cracking of weighted vacuum gas oil..... 5

Social sciences

- Kurmanov N., Satbayeva A., Rakhimbekova A., Makhatova A.* Human development index: place of Kazakhstan in the modern world..... 14
- Panzabekova A.Zh., Tyurabayev G.K.* The system of payment and stimulation of labor at real sector enterprises of Kazakhstan 20
- Tyurabayev G.K., Nesipbekov Ye. N.* Work with personnel pool as a constituent of educational organizations development process27

Physics

- Ramazanov T.S., Kodanova S.K., Issanova M.K., Tikhonov A., Kaikanov M.* Transport properties of inertial confinement fusion plasmas..... 34
- Ramazanov T.S., Kodanova S.K., Issanova M.K., Tikhonov A., Kaikanov M.* Transport properties of inertial confinement fusion plasmas..... 43

Chemistry

- Malyshev V.P., Zubrina Y.S., Makasheva A.M.* The role of the boltzmann-Shannon entropy in understanding the processes of self-organization..... 53
- Komekova N.M., Kozlov V.A., Zhurinov M.Zh.* Sulfuric acid atmospheric pressure leaching of vanadium black shale..... 62
- Bayeshov A.B., Adaybekova A.A., Abduvaliyeva U.A.* Electrochemical behavior of electrodes of molybdenum in sodium hydroxide solution..... 70

Earth sciences

- Metaksa G.P., Chekushina T.V., Moldabaeva G.Zh., Metaksa A.S.* Lake Baikal - natural reactor of hydrocarbons..... 77

Biologiya

- Khakimzhanov A.A., Mamytova N.S., Beskempirova Zh.D., Tilegen B., Dalelhankhyzy A., Kuzovlev V.A., Aitkhozhina N.A.* Wheat chitinase complex and some of its properties..... 85

Technical sciences

- Zhirnova O.V., Toigozhinova A.Zh., Zhakipov Zh., Turikanov T.S., Orazalin A., Matenov N.* Development of mathematical models for automated control system combustion process biogas to reduce greenhouse gas emissions..... 94
- Kaldybaeva B.M., Khusanov A. E., Dmitriev E.A., Sabyrkhanov D.S., Abilmagzhanov A.Zh.* Calculation of the bubble diameter, taking into account the hydrodynamic conditions and structural features in the chemisorption apparatus..... 106

Social Sciences

- Ayupova Z.K., Kussainov D.U.* To the question about the sources of the system of education in kazakh culture..... 115
- Dauletbaev B.D., Primzharova K.K., Konyrbekov M. Zh.* Modeling and assessment of the level of innovative activity in the sphere of intellectual potential of regions of the republic of Kazakhstan.....122
- Kaldybay K.K., Abdrassilov T.K., Nassimov M.O.* The problem of human in the religious heritage of modern kazakh thinkers..... 131
- Magay T.P.* Transformation of business education: an innovative approach..... 141
- Myrzagaliev A.B., Tuktassinova A.A., Samarkhanov T.N., Akzambek A.M.* *In vitro* introduction of *daphne* *Altaica pall*..... 151
- Ordabayeva M.* Current issues and trends of the medical and health tourism in Kazakhstan 161
- Satkalieva T.* Trends in energy sector of Kazakhstan..... 167
- Berik A.B.* Psychological support of children with mental retardation..... 176

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the work described has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the originality detection service Cross Check <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)

ISSN 2518-1483 (Online), ISSN 2224-5227 (Print)

<http://www.reports-science.kz/index.php/ru/>

Редакторы *М. С. Ахметова, Д. С. Аленов, Т.А. Апендиев*
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 09.12.2016.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
5,5 п.л. Тираж 2000. Заказ 6.